

REFERANCES

- Abdel-Hafez, A.I.I and El-sharouny, H.M.M. (1990):** Keratinophilic and Saprophytic fungi isolated from student's nails in Egypt. J.Basic. Microbiol. Vol. 30, No. 1, pp. 3-11.
- Ahmad, M.S.(1990):** Production of some antibiotics by certain microorganisms .Ph.D Thesis .Fac. Sci. Al-Azhar Univ.
- Alexander, M. (1961):** Introduction to soil Microbiology. John Wiley, New york.472 pp.
- Ali-Shtayeh, M.S; Arda, H.M. and Abu- Ghdeib, S.I. (1998):** Epidermiological study of tinea capitis in school children in the Nablus area (West Bank).Mycoses 41; 243-248.
- Al- Zarban, S.; I.H. Abbas; Al-Musallam, A., Steiner, U.; Stackebrandt, E. and Kroppenstedt, R.M. (2002):** *Nocardiosis halotolerans* Sp, nov., isolated from salt march soil in Kuwait. Int. J. of Syst. and Evol. Microbiol. 52: 525-529.
- Al-Sogair, S.M.; Moawad, M.K. and Al-Humaidan, Y.M. (1990):** Fungal infection as a cause of skin disease in the east province of Saudia Arabia, Tinea pedis and Tinea manuum .Mycoses, Vol.34 No. (7-8) pp: 339-344.
- Ajello, L and Gerog, L.K (1957):** In vitro hair cultures for differentiating between typical isolates of *Trichophyton mentagrophytes* and *Trichophyton rubrum*. Mycopathol. Mycol. Appl., 8:3.
- Ajello L, (1968):** A taxonomic review of the dermatophytes and related species. Sabouraudia. Feb; 6(2):147–159.
- Aly, R.(1994):** Ecology and epidermiology of dermatophyte infections. J. Amm. Acad. Dermatol. 31 (3pt 2): S 21-5.
- Anstey, A; Lucke, T.V and Philpot,C.(1996):** Tinea capitis caused by *Trichophyton rubrum*. Br.J. Dermatol. Jul. 135(1): 5-113.

- Augustine, S.K., Bhavsar, S.P. and apadnis, B.P. (2005):** Isolation of *Streptomyces albidoflavous* with antagonistic activity against pathogenic fungi. J. Biosci. 30 (20):201-211.
- Badji, B, Mostefaous, A, Saboous, N.; Lebrihi, A. and Mathieu, F. (2007):** Isolation and partial characterization of antimicrobial compounds from new strain *Nonomuraea Sp* .NM 94. J .Int. Microbiology Biotechnology 34(6): 403-12.
- Balajee, S.A; Ranganthan, S. and Menon, T. (1997):** Soil dermatophytes in Madras, India, Nov; 40 (7-8): 317-20.
- Baldacci, E., Spalia, C., and Grein, A (1954).**The classification of the Actinomycetes species (*Streptomyces*). Arch. Mirobiol.20, 347-357.
- Bargman, H.; Kane, J.; Lou Baxter, M. and Summerbell, R.C. (1995):** Tinea capitis due to *Trichophyton rubrum* in adult women. Mycoses, 38: 231-234.
- Bergus, G.R and Johnson, J.S. (1993):** Superficial tinea infection, Am.Fam.Pgysician. 48: 259-86.
- Bergy's (1974):** Bergyey's Manual of Determinative Bacteriology (Williams and Wilkins Co.) Baltimore.London.
- Block, R.J; Durrun, G., and Zweig, G. (1955):** A Manual of paper chromatograoh and paper Electrophoresis, New York. Acad-Press.
- Bocobo, F.C and Benham, R. W. (1949):** Pigment production in the differentiation of *T. mentagrophytes* and *T. rubrum*. Mycologia, 41: 291.
- Burkholder, P.R., Sun., S.H., Ehrlich, J., and Andrson, L. (1954):** Criteria of speciation in the genus *Streptomyces*. Ann. N.Y. Acad. Sci.60: 102-123.
- Caprilli F, Mercantini R, Marsella R, Farotti E. (1980):** Etiology of ringworm of the scalp, beard and body in Rome, Italy. Sabouraudia. , Jun; 18(2):129–135.

- Christofidou, M.; Panagiotopoulou, E.; Iacoidou, E. and Dimitracopoulos, G. (1996):** Dermatophyte species among patients with presumptive diagnosis of superficial mycoses. *Acta- Microbiologica – Hellenica*, 41 (1): 49-53.
- Cooper, R; Horan, A., Genittle, F.; Gullo, V., Loebenberg, D.; Marquez, J; Patel, M.; Puar, M. and Truumees, T. (1988):** A novel antifungal compound produced by a *Micromonospora* Sp. Taxonomy, fermentation, Isolation , Structure and biological reports .*J Antibiotic* Vol.41 No.1, PP. 13-19.
- Cooper, R; Truumees, I.; Gunnarsson, I.; Loebenberg, D.; Horan, A.; Marquez, J.; Partel, M.; Gullov, V.; Puar, M.; Das ,P. and Mittelman, S. (1992) :** A novel antifungal antibiotic from An *Actinoplanes* Sp. *J Antibiotic* Vol. 54 No. 4, pp : 444-453.
- Conant, N.F (1971):** Manual of clinical mycology. 3rd.ed. Philadelphia, Saunders.
- Cross, T., Maciver, A and Laoy, J. (1968):** The thermophilic Actinomycetes in mouldy hay *Micropolyspora foeni* Sp. *J. Gen. Microbiol.*, 50: 351.
- Cutler, J.E (1991):** Putative virulence factors of *Candida albicans*, *Ann. Rev. Microbiol*, 454: 187-218.
- Egorov, N. (1995):** Microorganisms antagonists and Biological methods for evaluation of antibiotic activity. Moskva. *Vissha Shkola* 200.
- El-Doraidy, E.H. (1998):** Potentiality of microbial isolates against dermatophytes fungi. M.Sc. Benha, Faculty of Science, Univ. of Zagazig.
- Elewski, B E. (1996):** Cutaneous mycoses in children. *Br.J.Dermatol.* 134 (Suppl 46): 7-11.
- El-Gammal, Amira, A.(1979):** Studies on neomycin produced by *Streptomyces fradiae*. *Egyptian J. of Microbiol.* 14 (1): 103-105.

- El-Mehalawy, A.A., Abd-Allah, N.A., Mohamed, R.M. (2005):** Actinomycetes antagonizing plant and human pathogenic. Int.J. of Agriculture and Biology vol. 7 No. 2:188-196.
- Emmons, C.W. (1934):** Dermatophytes: Natural grouping based on the form of spores and accessory organ. Arch. Dermatol. Syphilol, 30: 337-62.
- Emmons, C. W; Binford, C.H. and Kwonchung. K.J (1977):** Medical mycology, Lea and Febiger. 3th ed. Philadelphia.
- Enweani, I.B; Ozan, C.C; Agbonlahor, D.E. and Nadip, N. (1996):** Dermatophytosis in school children in Ekpoma. Nigeria. Mycoses, 39; 303-305.
- Espinel, A. Ingrff and S. Shadomy (2005):** In vitro evaluation of antifungal agents. European. J.of clinical Microbiol. & infection diseases vol. 8 No.4: 352-361.
- Fahmy, S.H. (1986):** Biological studies on grey *Streptomyces* isolates from Egyptian soil. M.Sc. Benha, Faculty of science, University of Zagazig.
- Feng N., Yew.Wup, Hung, Y., Xieh.Weix. (2007):** Two new antifungal alkaloids produced by *Streptoverticillium morookaense*. J. Antibiotic 60(3): 83-179.
- Flaig, W.; Beutelspeacher, H. Kuster, E and Segier Holzweissing, G. (1952):** Beitrage Zur physiologie and morphology der *Streptomycete*. Pl. Soil 4, 118.
- Flaig, W., and Kutzner, H.J (1954):** Zur Systematik der Gattung *Streptomyces*. Naturwissenschaften, 41, 287.
- Flaig, W.and Kutzner, H.J (1960):** Beitrag zur Systematik der Gattung *Streptomyces*, Waksman et Henrici., Arch. Mikrobiol. 35, 105-138.
- Flaig, W.; Kuster, E; and Beutelspacher, H. (1955):** Elektronenmikroskopische untersuchungen a sporen verschied ener *Streptomyceten*. Zenter. Backteriol. Parasitenk. Abt. 11, 108-376.

- Frey, D.; Oldfield, R.J. and Bundger, R.C. (1979):** a color atlas of pathogenic fungi, London, Wolf medical Publishers.
- Gauze, G.F.; Preobrazhenskaya, T.P.; Kudrina, E.S.; Blinov, N.o.; Ryabova, I.D.; and Sveshnikova, M.A. (1957):** Problems in the classification of antagonistic Actinomycetes, State Publishing House for medical literature (In Russian) Medzig, Moscow.
- Gauze,G.(1983):** Manual for determination of actinomycetes, Moskva, Nuka (in Russian).
- Germain, G. and Sumerbell, R. (1996):** Identification filamentous fungi.A clinical Lab Handbook 1st Ed. Belmont. California.
- Georgl k, (1954):** The relationship between the downy and granular forms of *Trichophyton mentagrophytes*. J Invest Dermatol. Aug; 23(2):123–141.
- Glazebrook, M.A; Vining, L.C; White, R.L; Smith, K.C. and Chedrawy, E.G. (1993):** Nutrient effects on growth and the production of 5- hydroxyl-oxonorvaline by *Streptomyces akiyoshiensis*. Can.J.Microbiol.39 (5): 536-542.
- Golt, P.K; Bhalta, H.E and Andeson, H.W. (1951):** Endomycin, a new antibiotic. Phytopathology 41 (5): 393-400.
- Gordon, (1966):** Some criteria for recognition of *Nocardia madura* (Vincent) Blanchard. J. Gen. Microbiol. 45 = 355 - 364.
- Goodfellow, M; and Orchard, V.A. (1974):** Antibiotic sensitivity of some nocardio-form bacteria and its values as a criterion for taxonomy. J. Gen. Microbiol. 83, 375-387.
- Gordon, N.A. (1976):** Characteristics of *Dermatophilus angolensis*, its affinities with the actinomycetes and differentiation from *Geodermatophilus*. In: *Dermatophilus infection in Animal and Man* (D.E. Liloyd and K.D. Sellars, eds.) pp = 187 – 201. Acad. Press. London.

- Gupta, R.C. and Tandon, R.N (1977):** Growth inhibition of fungi by volatiles from *Streptomyces*. Trans. Brit. Mycol. Soc.68: 438-439.
- Gupta, A.K; Kauder, D.N; Shear, N.H (1994):** Antifungal agents: an overview. Part 1. J.Am,Acad. Dermatol. 30 (5 pt): 677-98.
- Hacene, H. and Lefebvre, G. (1995):** A new non-polyenic antifungal antibiotic AH17 produced by a strain of *Spirillospora*. Microbiol. 83(336): 199-205.
- Hay, R.J; Reid, S.; Talwat, E; Macnamara, K (1983):** Immune response of patients with tinea imbricate. Brj. Dermatolo. May, 108 (5): 581-586.
- Hay, R.J (1992):** Clinical manifestation and management of superficial fungal infection in the compromised patient in fungal infection the compromised patient, edited by D.W, Wamock and M.D, Richaradson. John Wiley & sons.
- Hazen, E. and Brown, R. (1951):** Fungicidin, an antibiotic produced by soil actinomycetes. Proc. Soc. Expt.Biol and Med. 76 (1): 93-97.
- Hasegawa, S., Kondo, N., and Kodama.F (1990):** Suppression of *Fusarium* with of adzuki bean by rhizosphere microorganisms. Pest. Alter. 477-486.
- Hesseltine, C.W., Benedict, R.G. and Pridham, T.G. (1954):** Useful criteria for species differentiation in the genus *Streptomyces*. Ann. N.Y Acad.Sc. 60, 136-151.
- Hopwood, D.A. and Merric, M.J. (1977):** Genetics of antibiotic production. Bacteriol.Rev. 41, 595-635.
- Igarashi, M.;Kinoshite, N.; Ikeda, T.; Nakagawa, E.; Hamada, M. and Takeuchi, T.(1997):** Formamycin, a novel antifungal antibiotic produced by a strain of *Saccharothrix sp.* I. Taxonomy, production, isolation and biological properties J. Ant.Nov: 50(11): 926-931.

- Isoa, T. (1957):** An antifungal substances Flavacid studies on the antibiotic substances from actinomycetes, J Antibiotic (Tokgs), 6 (3): 117-121.
- I.S.C.P (1964):** Methods manual of International Cooperative Project for Description and Deposition of type cultures of *Streptomyces*. Ohio, Weselyan.
- Jain, P.K and Jain P.C. (2005):** Antifungal activity of some actinomycetes isolated from various habitats .Hinustan Antibiotic Bull. 54-46(1-4):5-10.
- Jain, P.K and Jain P.C. (2007):** Isolation, characterization and antifungal activity of *Streptomyces sampsonii* GS 1322. Indian Exp.Biol.45 (2): 6-203.
- Jeuniaux, C. (1966):** Chitinases methods. Enzymol. 8: 644-650
- Jones, K (1949):** Fresh isolates of actinomycetes in which the presence of sporogenous aerial mycelia is a fluctuating characaristics. J. Bacteriolo. 57: 141-145.
- Johnson, M.J (1955):** Recent advances in Penicillin fermentation. Bull. World Health Org.6: 99-121.
- Johnson, L.F., Curl, E.A, Bond, J.H. and Fribourg, H.A. (1960):** Methods for studying soil microflora- plant disease relationships.2nd printing. Burgess Publishing Company. Minnea, Polis 1960.
- Kane J, Fischer JB. (1971):** The differentiation of *Trichophyton rubrum* and *T. mentagrophytes* by use of Christensen's urea broth. Can J Microbiol. Jul; 17(7):911- 913.
- Kaneda, M.; Naid, T.; Kitahara, T.M.; Nakamura, S.; Hirata, T. and suga, T (1988):** Carbazomycin G and H, novel carbazomycin congeners containing a quinol moiety J. Ant. 41(5): 606-608.
- Ketoki, B. and Majumdar, S.K. (1973):** Utilization of carbon and nitrogen sources by *Streptomyces kanamyceticus* for kanamycin production. Antimicrobial agent's che. 4(1): 6-10.

- Knight AG. (1972):** A review of experimental human fungus infections. J. Invest Dermatol. Oct; **59**(4):354–358.
- Kotake, C; Yamasaki, T.; Moriyama, T.; Shinoda, M.; Komiyama, N.; Furamai, T.; Konishi, M. and Oki. (1992):** Butyrolactols A and B, new antifungal antibiotics. Taxonomy, isolation, physicochemical properties, structure and biological activity. J. Antibiotic 45(9): 1442-1450.
- Krassilinkov, N.A (1950):** Actinomycetes antagonistic and antibiotic substances Jzol.Akad. Nauk. SSSR. Moscow. USSR.
- Krassilinkov, N.A (1960):** Taxonomic principles in the actinomycetes J.Bacteriol.79, 65-71.
- Krassilinkov, N.A (1970 a):** Pigmentation of actinomycetes and its significance taxonomy. The Actinomycetes (Prauser Ed,) 123-131.
- Krassilinkov, N.A (1970 b):** Ray fungi Higher forms. Nauka. Moscow, USSR.
- Kriss, A.E.; Rukina, E.A.; and Issaiev, B.M (1945):** Electron microscopic studies on the structure of actinomycetes. Mikrobiolgiya, 14, 127.
- Kutzner, H.J (1981):** The family *Streptomycetaceae* in the prokaryotes, A handbook of habitats, Isolation and Identification of bacteria. Vol.II (M. Pstarr, H. Stolp, H.G. Truper A. Balows and H.G.Schlegel Eds) 2028-2090.
- Kuster, S and Williams S.T (1964):** Production of hydrogen sulphide by *Streptomyces* and methods for it detection. App. Microbiol.12: 46-52.
- Lackner, T.E. and Clissold, S.P. (1989):** Bifonazde, A review of its antimicrobial activity and therapeutic use in superficial mycoses. Drugs. Agu. 38(2): 204-225.

- Lechevalier, M. P. and Lechevalier, H.A. (1970):** chemical composition as a criterion in the classification of aerobic Actinomycetes. *Int. J. Syst. Bacteriol.* 20, 435-443.
- Lechevalier, H.A; Lechevalier, M.P; and Gerber, N.N. (1971):** chemical composition as a criterion in the classification of Actinomycetes. *Adv. Appl. Microbiol.* 14, 47-72.
- Lenhart K. (1970):** Grisofulvin resistance in dermatophytes. *Experientia.* 15; 26(1):109–110.
- Lloyd, A.B. (1969):** Dispersal of *Streptomyces* in air. *J. Gen. Microbiol.* 57, 35-40.
- Locci, R. and Sharples, G.P (1984):** Morphology. In Goodfellow, Mordarski and Williams (Editors). In *Biology of the actinomycetes*, Acad. Press.London, pp: 165-199.
- Mac ginnies, M.R. (1980):** Laboratory hand book of medical mycology. Academic Press, New York pp: 661.
- Mahgoub, E.S and Baxter, M. (1983):** Superficial mycoses at King Abd-Al- aziz teaching hospital in Riyadh. Proceeding of the eightin Con. of Int. Soc. for human and animal mycology 397-400.
- Marchisio, V.F; Preve, L. and Tullio, V. (1996):** Fungi responsible for skin mycoses in Turin (Italy) *Mycoses*, 39; 141-150.
- Maruyama, R; Katoh, T. and Nishioka, K. (1998):** Demonstration of dermatophyte dissemination from the infected soles using the foot-press method. *Mycoses*, 41; 145-151.
- Martin, J.F. and Demian, A.L. (1980):** Control of antibiotic synthesis. *Microbiol. Rev*, 44: 230-251.
- Maslen, M.M and Andrew, P.J (1997):** Tinea due to *Trichophyton violaceum* in Victoria. Australia, *Australas J. Dermatol. Agu*; 38 (3): 8-124.

- Miguens, M.P; Pereiro, M. and Pereiro, M.I (1991):** Review of dermatophytosis in Galicia from 1951 to 1987 and comparison with other areas of Spain. *Mycopathologia*. Vol. 113, No.2, pp. 65-78.
- Misrask and Sahuks (1977):** screening of some antifungal agents against dermatophytes. *Indian j. of pharmacology* 9(3):195-198.
- Monk.J.P; Brogden, R.N. (1991):** Naftifine, A review of its antimicrobial activity and therapeutic use in superficial dermatomycosis, *Drugs*. Oct.42 (4): 659-672.
- Mostafa, M.A; Osman, H.G. and Abo-Zeid, Z.K. (1968):** The production of tetracycline by *Streptomyces aureofaciens*. *J. Microbiol. USSR* 3(1): 1-9.
- Naik, S.R, Desaisk, Nanda, R.K; Narayanan, M.S. (2007):** Fermentation, isolation, purification and biological activity of SJA-95, a heptane polyene macrolide antibiotic produced by the *Streptomyces Sp.* Strain S24. *Arzneimittelforschung* 57(3):9-171.
- Nair, M.G; Putnam, A.P; Mishra, S.K; Mulks, M.H; Taft, W.H; Keller, J.R; Zhu, P.P; Meinhart, J.D. and Lyum, D.G.(1989):** Faerie fungi, a new broad spectrum antibiotic from *Streptomyces griseus* Var. autotrophicus. *J. Na.Prod.* 52 (4): 797-809.
- Ninomiya, J, Ide, M.; Ito, Y. and Takiuchi, I. (1998):** Experimental penetration of *Trichophyton mentagrophytes* into human stratum corneum. *Mycopathologia*, 141: 153-157.
- Norris, H.A; Elewskib.E. And Ghannoum, M.A (1999):** Optimum growth conditions for the determination of the antifungal susceptibility of three species of dermatophytes with the use of a microdilution method. *J.Am. Acad. Dermatol.* Vol.40 (2): pp: 9- 13.
- Nowicki, R (1996):** Dermatophytosis in the Gdansk area, Poland; a 12-years Survey. *Mycoses*, 39; 399-402.

- Nouguchi, H., Hiruma, M.; Kawada, A.; Ishibashi, and Kono, S. (1995):** Tinea pedis in members of the Japanese self-defence forces: relationships of its prevalence and its severity with length of military service and with of interdital spaces. *Mycoses*, 38; 495-494.
- Nuesh, J (1965):** Isolierung and selektionierung von A ktinomyceten in: Symposium anreicherung skulturund Mu-t enauslse. Gottunen April 1964, Zentralblantt fur Bakteriologie. Parasitenkunde, Infektion skrnk Heiten und hygiene, Abt.1 supp, 1:234-252.
- Nurettin F. (2002):** Investigation of the antimicrobial Activity of some *Streptomyces* isolates. *Tuck. J. Biol.* 27: 79-84.
- Nwobu, R.A.V and Odugbemi, T (1990):** Fungi causing dermatophytosis in Lagos, Nigeria. *East.African-Medical –J.*, 67 (4): 246-249.
- Oxoid, L. (1982):** The oxid manual of culture media, ingredient and other Laboratory sevies. Turnergraphic Ltd England 4th Ed.
- Padhya, A.A; Young, C.N. and Ajello.L. (1980):** Hair perforation as a diagnostic criterion in the identification of *Epidermophyton*, *Microsporum* and *Trichophyton* species. PAHO Sci. Pub No.396, pp.115.
- Patel, M.; Hegda, V.; Horan, A.C.; Gullo, V.P; Loebenberg, D.; Marquez, J.A.; Miller, G.H; Puar, M.S. and Waitz, J.A (1984):** A novel phenazine antifungal antibiotic, 1,6-dihydroxy-2-chlorophenazine. Fermentation, isolation, structure and biological properties. *J. Antibiotic* 37 (9): 943-948.
- Paveia MH. (1975):** Culture medium alkalization by dermatophytes. (Influence of time and temperature of incubation). *Mycopathologia*. Feb 28; 55(1):35–40.
- Philpot, C. (1967):** The differentiation of *Trichophyton mentagrophytes* from *Trichophyton rubrum* by a simple urease test. *Sabouraudia* 5: 189.

- Praveen Kumar Jain, P.C Jan (2002):** Production of heptane antifungal antibiotic by *Streptomyces purpeofuscus* CM 1216. Indian J. Exp. Biol. 43 (4): 342-515.
- Praveen Kumar Jain, P.C Jan (2005):** Antifungal activity of some actinomycetes isolated from various habitats. Hindustan Antibiot. Bull 45-46 (1-4).
- Praveen Kumar Jain, P.C Jan (2007):** Isolation, characterization and antifungal activity of *Streptomyces sampsonii*. Indian J. Exp. Biol. 45 (2):203-206.
- Pridham, T.G; Felser, L.A; Shotwell, O.L.; Stodola, F.A.; Benedict, R.G.; Foley, C. and Mitchell, P.W(1956):** Antibiotic against plat disease.(1) Lab. And green house Survey Phytopathology. 46: 568-575.
- Pridham, T.G; Hesseltine, C.W and Benedict R.G. (1958):** A guide for the classification of *Streptomyces* according to selected groups. Placement of strains in morphological section. Appl. Microbiol. 6; 52-79.
- Ponnighous, J.M; Yvonne, C. and Warndorf.F. (1996):** The spectrum of dermatophytes in Northern Malawi (Africa, Mycoses; 39: 293-97.
- Ray tapadar, S. and Paul, A.K. (2001):** Production of an antifungal antibiotic by *Streptomyces aburavrensis* IDA-28. Microbiol. Res. 155(4): 315-323.
- Rebell, G. and Taplin, D. (1970):** Dermatophytes, their recognition and identification. University of Miami Press, Coral Gables. Florida.
- Raddi, G.S., and Rao, A.S (1971):** Antagonism of soil actinomycetes to some soil –borne plant pathogenic fungi, Indian phytopath. 24, 649-657.
- Richardson, M. and wornock D (1997):** Fungal infection Diagnosis and Management. 2nd.ed Oxford England.

- Rippon, J.W. (1974):** Ascomycetes, the dermatophytes in fungi pathogenic for human and animal>Vol. 3 Mareel Dekker, Inc. New York and Basal pp: 113.
- Rippon, J.W. (1988):** Medical Mycology 3rd Edi.W.B Saunders Co.Philadelphia.
- Roberts, F.B. (1995):** Basic medical Microbiology the pathogenic fungi, Little Brown and company 5th ed. 33: 481.
- Sabota, J., Brodell, R., Rutecki, G.W. and Hoppes, W.L. (1996):** Severe tinea barbae due to *Trichophyton verrucosum* infection in dairy farmers. Clin. Infect. Dis. Dec.; 23(6): 10-1308.
- Shah PC, Krajden S, Kane J, Summerbell RC (1988):** Tinea corporis caused by *Microsporum canis*: report of a nosocomial outbreak. Eur J Epidemiol. Mar; 4(1): 33–38.
- Shirling, E.B. and Gottlieb, D. (1966):** Methods for characterization of *Streptomyces* species. Int. J. Syst. Bacteriol. 16: 313-340.
- Shirling, E.B. and Gottlieb, D. (1968):** Cooperative description of type cultures of *Streptomyces* II Species description from first study. Int.J. Bacteriol. 18: 69-189.
- Staib F, Mishra, S.K; Formtling, R.A; Bilsse, A; Kleinkauf, F.H; Mahmutoglu, I.; Muller, K.D.; Salnkow; J.; Holnne, G. and Schwarz, H. (1983):** A new polyene with antimycotic activity produced by *Streptomyces* from rhizosphere of an indoor plant. Arzneimette. L- forschung 33(1): 84- 87.
- Stanely, W., Chapman, M.D and Ralph Danid, C. (1994):** Cutaneous manifestation of fungal infection. Infection disease clinic of North America 8 (4): 879-910
- Takiguchi, Y.H; Muramatsu, M.S; Mishina, I.J.H. and Mibencin, M.(1981):** A new family of macrolides antibiotics, fermentation, isolation and physico-chemical properties of Mibeny. Cins, D.E.F and H.J. antibiotics, 36: 502-508.
- Thirumalachar, M.J; Menon, M.V. and Bhatt, V.V (1961):** Polyene antibiotic- hamycin. Hind. Antibiotic Bull, 3: 136-138.

- Thirumalachar, M.J; Menon, S.K. (1962):** Polyene antibiotic-dermostatin. Hind Antibiotic Bull, 4: 106-108.
- Thirumalachar, M.J; Rahalkar, P.W. and Gopalkrishnan, K.S(1964):** Polyene antibiotic – aureofungin. Hind. Antibiotic Bull.6: 108-111.
- Thomas, C. and Raper, K.B (1945):** An Annual of the Aspergilli. Williams and Williams. Co. Baltimore 1md.
- Thomas,P.; Korting, H.C; Strassl, W and Ruzicka, T.(1994):** *Microsporium canis* infection in a 5-year old boy: Transmission from the interior of a second hand car. Mycoses, 37: 14-142.
- Tresner, H.D., Davies, M.C. and Backus, E.L (1961):** Electron microscopy of *Streptomyces* spore morphology and its role in species differentiation. J. of Bacteriol. 81: 70-80.
- Tresner, H.D., and Backus, K.J (1963):** System of color wheels for *Streptomyces* taxonomy. Appl. Microbiol. 11, 335-338.
- Ueki, M.; Kusumoto, A.; Hanafi, M.; Shibata, K; Tanaka, T. and Tangiuchi, M.(1997):** A novel antifungal antibiotic UK-3A from *Streptomyces* Sp. 517-02. Fermentation, isolation, structural elucidation and biological properties J. Antibiotic 50 (7): 551-555.
- Waksman, S.A (1919):** Cultural studies of species of Actinomycetes. Soil Sci. 8,77-215.
- Waksman, S.A (1959):** Strain specificity and production of antibiotic substances. X. Characterization and classification of species within the *Streptomyces griseus* group. Proc. Nat. Acad. Sci. U.S.A.
- Waksman, S.A (1961):** The actinomycetes, classification, identification and descriptions of genera and species, Vol.2, The Williams and Wilkins Co. Baltimore. pp. 1-363.
- Waksman, S.A, and Lechevalier, H.A (1962):** The Actinomycetes. vol. 111. Antibiotics, Baltimore, the Williams and Wilkins company U.S.A, 430

- Williams, David., Marten, R.H. & Sarkany, I. (1959):** Griseofulvin, British Journal of Dermatology, 71(12), 434-443.
- Williams, S.T, and Cross, T. (1971):** Isolation, purification, cultivation and preservation of actinomycetes. Methods Microbiol.4, 295-334.
- Williams, S.T. and Khan, M.R. (1974):** Antibiotics-a soil microbiologist's view point. Postepy Higieny; Medycyny Toswiadenzalij. 28, 259-408.
- Wishart, J.M. (1994):** A double blind study of itraconazole vs. griseofulvin in patients with tinea pedis and tinea manus. N.Z. Med.J.107: 8-126.
- Yano, K.; Yokoi, K.; Sato, J.; Oono, J.; Kouda, T.; Ogawa, Y. and Nakashima, T. (1986):** Actinopyrons A, B and C, new physiologically active substances. J. Antibiotic 39 (1): 32-37.
- Yano, K.; Oono, J.; Mogi, K.; Asaoka, T. and Nakashima, T. (1987):** Pyrroxamycin, a new antibiotic taxonomy, fermentation, isolation structure. Determination and biological properties. J. Antibiotic 40 (7): 961-969.
- Yassin, M.H. and Galal, A.M. (1998):** Antiphytoviral potentialities of some fungi and actinomycetes isolates against kidney bean plants infected with T.N.V. 1st conf. Protec. Egypt.pp: 156-161.
- Vertesy, L.; Aretz, W.; Ehlers, E.,; Hawser, S.; Isert, D.; Knauf, M; Kurz, M.; Schiell, M.; Vogel, M. and Wink, J. (1998):** A novel antifungal, heptane antibiotics produced by *Streptomyces* Sp. H A.G 003874. J. Antibiotic 51(10): 921-928.
- Yu, R. J, Harmons, S.R; Wacter, P.E and Blank, F. (1969):** Hair digestion by keratinase of *Trichophyton mentagrophytes*. J.Irest.Dermatol.53:166-171.