

The background features a light green and white polka-dot pattern. A large, stylized white flower with a pink center is prominent in the upper left. A white ribbon with a pink circular detail is draped across the middle. In the lower right, there are decorative swirls and a small white flower.

REFERENCES

REFERENCE

- Anwer, S, Mustafa, S. and Ashraf, S.:* Rapid identification of neonatal sepsis. J. Pak. Med. Assoc. 45: 94, 2000.
- Assicot, M., Gendral, D. and Carsin, H.:* High serum procalcitonin concentration in patients with sepsis and infection. Lancet. Infect. Dis. 341: 515, 2005.
- Athhan, F., Genel, F. and Bak, M.:* PCT: a marker of neonatal sepsis. J. Trop. Pediatr. 48: 10, 2004.
- Atli, D., and Petur, L. :* Neonatal sepsis: diagnosis and initial antibiotic therapy. Pediatr . clin. North Am. 34: 219, 2005
- Auriti, C., Maccallini, A. and Diliso, D.:* Risk factors for nosocomial infection in a neonatal intensive care unit. J. Hosp. Infect. 53: 30, 2003.
- Backer, C.:* Group- B streptococcal infection. Crit. Care med. 27: 120, 2005.
- Balci, G., sungurtekin, H. and Gurses, E.:* Usefulness of procalcitonin for diagnosis of sepsis in the intensive care unite. Crit. Care Med. 7: 85, 2003.
- Bataineh, H., and Al Rashed, K. :* C-reactive protein in neonates with suspected septicemia. Rawal medical journal. 32:24,2007.
- Baungart, S., Halls, E. and Campos, J.:* Sepsis with coagulase negative staph in critically ill newborn. Am. J. Dis. Child. 137: 661, 2004.

-
- Benaor, N., Siegrist, C. and Gendrel, D.:* Procalcitonin is a marker of severity of renal lesions in pyleonephritis. *Pediatrics*. 102: 1425, 2005.
- Berger, M.:* Complement deficiency and neutrophil dysfunction as a risk factors for bacterial infections in newborn. *Rev. Inf. Dis.* 12: 405, 2000.
- Boo, N.Y., Azlina, A.A., Rohana, J.:* Usefulness of semi- quantitative procalcitonin test kit for early diagnosis of neonatal sepsis. *Singapore Med. J.* 49: 204, 2008.
- Bracq, S.:* Calcitonin gene expression in human liver. *FEBS*. 331 : 18, 1993.
- Brain, S., Tippins, J. and Morris, H.:* Potent vasodilator activity of calcitonin – gene related peptide in human skin. *J. Invest. Dermatol.* 87: 533, 1996.
- Brood, P., Symes, A. and Craig, R.:* Structure and methylation of the human calcitonin ALPHA – CGRP gene. *Nucleic Acid Res.* 17: 7011, 1989.
- Brunkhorst, F., Heinz, U. and Forycki, F.:* Kinetics of procalcitonin in iatrogenic sepsis. *Intensive Care Med.* 24: 892, 2001.
- Brunkhorst, F., Wagner, J. and Forckin, Z.:* Discrimination of bacterial and non infectious etiology of ARDS (adult respiratory distress syndrome) with procalcitonin immnoreactivity. *Intensive Care Med.* 382 : 372, 2003.
- Caplan, E., Oed, L. and Kassel, R.:* Cytokines serum level in neonatal sepsis. *Ann Surg.* 15: 210, 2005.

Castelli, G., Pognani, C. and Cita, M.: Procalcitonin, C- reactive protein, white blood cells and sepsis score in ICU: diagnosis and monitoring of sepsis. *Minerva Anestesiologica.* 72:69, 2006.

Castillo, M., Sacheen, A. and Lefebuse, P.: Amylin/ Islet amyloid polypeptide.: biochemistry, physiology, pathophysiology. *Diabet Metab.* 21: 25, 1995.

Chiesa, C., Panero, A. and Rossi, N.: Reliability of procalcitonin concentration for the diagnosis of sepsis in critically ill neonates. *Clin. Infec. Dis.* 26: 672, 2004.

Chiesa, C., Carsin, H. and Buck, C.: Procalcitonin: a new marker for bacterial infection. *Ann. Bisl.clin.* 58: 149, 1998.

Cole, F.: Infection: an immunologic defense mechanism. *J. Clin. Endocrinol. Metab.* 79: 1605,1999.

D'Angio, C., MCGowon, K and Baungart, S: Surface colonization with coagulase negative Staphylococci in premature neonates. *J. Pediatr.* 114: 1034, 1999.

Delma, S., Monneret, G. and Lapillonne, A.: Procalcitonin & C- reactive protein kinetics in neonates with group B-streptococcal infection. *Ann Biol. Clin.* 58: 208, 2003.

Eberhard, O., Haubitz, M. and Brunkhorst, F.: Usefulness of procalcitonin for differentiation between activity of systemic autoimmune disease and invasive bacterial infection. *Arthritis Rheum.* 40: 1256, 2005.

Edward, M., Jackson, C. and Baker, C.: Twin pregnancies, incidence of group- B streptococcal colonization and disease. *J. Pediat.* 97: 635, 2001.

Engle, W., Rosenfield, C. and Mouzinho, A.: Circulating neutrophil in septic preterm neonates. *Pediatrics.* 99: 10, 2005.

Fanaroff, A., Korores, S. and Wright, L.: Incidence, presenting features, risk factors and significance of late onset septicemia in very low birth weight infants. *Pediatr. Infect. Dis.* 17: 598, 2005.

Frausts, M.S., Pittet, D. and Costigan, M.: The natural history of the systemic inflammatory response syndrome (SIRS). *JAMA.* 273: 117, 1995.

Feigin, R.D, Abramson, S.L. and Edwards, M.S.: Infection in the fetus and neonate. *Pediat. Infect. Dis. J.* 4: 763, 2000.

Ferrieri, P.: Neonatal susceptibility & immunity to major bacterial pathogens. *Rev. Infect. Dis.* 4: 5394, 2006.

Fioretto, N., Rossano, C. and Sandra, M.: Procalcitonin in children with sepsis and septic shock. *J. paediatr. (Rio J).* 83: 323, 2007.

Fischer, G.: Use of IV IG in newborn. *Clin. Exp. Immunol.* 5: 325, 2004.

Gasser, L.: Usefulness of body surface cultures in newborn with early infection risk. *An Esp. Pediat.* 45: 40, 2006.

Gendrel, D., Assicot, M. and Raymond: Measurement of procalcitonin levels in children with bacterial and viral meningitis. *J. Pediatrics.* 128: 573, 2006.

Ghosh, A., Mittal, M and Jaganthen, G: Early diagnosis of neonatal sepsis. *Inian J. Med. Sci.* 55: 500, 2004.

Gottof, S.: Infections of the neonatal infant: Epidemiology, immunity and pathogenesis. *Arch Dis. Child.* 15: 525, 1996

Gramm, H., Beier, W. and Oedran, A: Procalcitonin a biological marker of the inflammatory response with prognostic properties. *Clin. Intensive Care.* 1: 3, 2005.

Guida, J., Runing, A. and Leaf, K: Platelet count and sepsis, is there an organism specific response. *Pediatrics.* 11: 315, 2003.

Hague, K: Infection & immunity in the newborn. *Lancet. Infect. Dis.* 12: 316, 2002.

Hanson, Z., Karlsson, B. and Jalil, F.: Antiviral and antibacterial factors in human milk: Biology of human milk. *J. Egypt. Soc. Obstet. Gyneco.* 4: 157, 1999.

Hasanc, C., Bodu, U. and Cipic, D: Procalcitonin: a sensitive marker of severe infection in immunosuppressed children with cancer. *Intensive Care Med.* 2: 211, 2004.

Hebra, A., Diello, C. and Russil, A: Necrotizing enterocolitis. *JAMA.* 5: 874, 2006.

Hoovers, F., Redeker, E. and Speleman, E: High resolution chromosomal localization of the human calcitonin/ CGRP/ IAPP gene family members. *Genomic.* 15: 529, 1993.

Jacobson, J.A., Chester, T.J. and Fraser, D.W.: Bacterial infection of newborn. *Clin. Inf. Dis.* 26: 187, 2006.

-
- Jerome, O:* Bacterial sepsis and meningitis. J. Pediatr. 5: 966, 2001.
- Joyce, C:* CGRP levels are elevated in patients with sepsis. Surgery. 108: 1101, 2006.
- Kaufman, D. and Fairchild., K.D.:* Clinical microbiology of bacterial and fungal sepsis in very-low birth weight infants. Clin. Microbiol. rev. 17: 638, 2004.
- Kemp, A. and Campbell, D. :* The neonatal immune system. J. Pediatr. 101: 1025, 2006.
- Kennon, C., Overtur, G. and Sierra, M.:* Granulocyte colony stimulating factor as a marker for bacterial infection in neonates. J. Pediatr. 128: 769, 1996.
- Klein, J.:* Bacterial sepsis and meningitis. Med. Clin. N. Am. 4: 90, 2005.
- Kliegman, R.:* Prenatal risk factors in neonatal infections. Ind. J. Pediat. 59: 335, 1998.
- Kieft, B. M., Overtur, G. and Sierra, M.:* Early diagnosis of septicemia in newborns. Clinical studies and sepsis score. Eur. J. Pediatr.45:56,2006.
- Kocherlakota, P:* Preliminary report: GM- CSF may reduce the incidence of neonatal sepsis in pre-eclampsia associated neutropenia. Pediatrics. 102 : 1107, 2003.
- Lacour, A, Gervaix, Z. and Zamara, S:* Procalcitonin, IL 6, IL 8, IL 1 and C – Reactive protein as identifiers of serious bacterial infections in children with fever without localizing signs. Eur. J. Pediat. 160 : 100, 2006.

-
- Laforgia, N., Coppola, B. and Carbone, A. :* Rapid detection of neonatal sepsis using polymerase chain reaction. *Acta pediat.* 80 : 1097, 2004.
- Lee, C., Tang, R. and Ching, R:* Evaluation of different blood culture media in neonatal sepsis. *J. Microbiol. Immunol. Infect.* 33: 165, 2005.
- Leeming, H.D., Labaune, J. and Issac, C.:* Laboratory aids for diagnosis of neonatal sepsis. *Acta paediatrica.* 47:451, 2006.
- Lerman, V.F. :* Lack of specificity of blood culture for sepsis diagnosis in premature infants . *Lancet.* 312:1211, 2007.
- Leon, A., Lepouse, C. and Assicot, M:* Procalcitonin concentrations in pleural effusion as a marker of septic response. *Shock.* 14: 102, 2005.
- Lewis, D.:* Developmental immunology and role of host defense in neonatal susceptibility to infection. *J. pediatr.* 108: 11, 2005.
- Linda, B.:* Neonatal sepsis, the scientific basis for practice. *Neonatal network.* 10: 40, 2002.
- Linder, N., Ohel, G. and Gazit, G. :* Neonatal sepsis after prolonged premature rupture of membranes. *J. Perinatol.* 1: 36, 1995.
- Lock, G.:* Infection, antibiotics and preterm delivery. *Clin. Infect. Dis.* 25: 309, 2001.
- Lott, W.:* Neonatal bacterial sepsis. *Crit. Care. Nurse. Clin. North Am.* 15: 65, 2006.
- Maisels, M., Khare, S. and Kring, P.:* Risk of sepsis in newborn with severe hyperbilirubinemia. *Peditrics.* 90:741, 2002.

Manusha, V., Rusia, U. and Skik, M: Utility of hematological parameters and CRP in detection of neonatal sepsis. *Pediat. Infect. Dis. J.* 125: 1201, 2004.

Maruna, P., Nedelnikova, K. and Gurlich, A: Physiology & genetics of procalcitonin. *Physiol. Res.*4: 60, 2003.

Meisner, M., Palmers, T. and Schmidt: Comparison of procalcitonin and C- reactive protein plasma concentrations at different SOFA scores during the course of sepsis and MODS. *Critical Care.* 3 : 55, 2004.

Meisner, M., Schmidt, J. and Huttner, H.: The natural elimination rate of procalcitonin in patients with normal and impaired renal function. *Intensive Care Med.* 2 : 216, 2005.

Meisner, M., Schmidt, J. and Rausch, M.: Postoperative plasma concentration of procalcitonin after different types of surgery. *Intensive Care Med.* 24 : 684, 2002.

Meisner, M., Schmidt, J. and Schnabels: Procalcitonin influence of temperature, storage, anticoagulation and arterial or venous asservation of blood samples on procalcitonin concentration. *Eur. J. Clin. Chem .Biochem.* 35: 601, 2003.

Meisner, M., Tschikowsky, K. and Scuttler, J.: Procalcitonion: a marker or modulator of the acute immune response. *Intensive Care Med.* 22: 14, 1996.

Monneret, G., Labaune, J. and Issac, C Procalcitonin and C – reactive protein levels in neonatal infections. *Acta Pediat.* 82: 209, 2004.

Moullec, J., Jullienne, A. and Gulian, J.: The complete sequence of human preprocalcitonin. FEBS. 176: 97, 1984.

Nelson, W.E., Vaughan, V. C. and Behrman, R. E.: Infections of the neonatal infant. In: Nelson Textbook of Pediatrics. B. Saunders company: 381, 17th edition, 2007.

Noya, F: Prevention of GBS infection. Infect. Disease. Clin. North Am. 6 : 41, 2002.

Nylen, E., Muller, B. and Snider, R: Pathophysiological significance of calcitonin precursors in sepsis and systemic inflammation. Shock. 12:14, 1999.

Nylen, E., Snider, R. and Thmpson, K: Pneumonitis associated hyperprocalcitoninemia. Am. J. Med. Sci. 18 : 3212, 1996.

Oberhoffer, M., Bgel, D. and Meier – Hellmann, A: Procalcitonin is higher in non – survivors during the clinical course of sepsis, severe sepsis and septic shock. Intensive Care Med. 3:22, 2005.

Oberhoffer, M., Stonans, I. and Russwurm, S: Procalcitonin expression in human peripheral blood mononuclear cells and its modulation by lipopolysaccharides and sepsis related cytokines in vitro. J. Lab. Clin. Med. 55: 134, 1999.

Ottolini, M., Lungren, K. and Cason, S.: Utility of complete blood count and blood culture screening in the diagnosis of neonatal sepsis. Pedat. Infect. Dis. J. 17: 235, 2003.

Persson, E., Trollfors, B. and Brendberg L.: Septicemia and meningitis in neonates and during infancy. Acta, Pediatr. 10: 92, 2002.

Petitjean, S., Assicat, M. and Bohuou, C. : Procalcitonin is a new biological marker of human septic response. Clin. Intensive Care. 2: 88, 1994.

Petzold, L., marchard, M. and Marie, F: Determination of procalcitonin in the diagnosis of maternal – fetal infections. Ann Clin. Bioch. 56: 602, 2004.

Polin, R: The "Ins and Outs" of neonatal sepsis. J. Pediat. 143: 8, 2003.

Powel, K.: Laboratory aids for diagnosis of neonatal sepsis. J. Clin. Invest. 4: 1240, 2005.

Ragharen, M., Mandal, G. and Bhat, B. : Prenatal risk factors in neonatal infections. Ind. J. Pediat. 3: 335, 2002.

Reinhart, K., Karzai, W. and Meisner, M: Procalcitonin as a marker of the systemic inflammatory response to infection. Intensive Care Med. 26 :1193, 2004.

Reith, H., Debus, E. and Thieda, A: Procalcitonin in early detection of post operative complications. Dig. Surg. 15: 200, 2006.

Russwurm, S., Wiederhold, M. and Oberhoffer, M: Molecular aspects and natural source of procalcitonin. Clin. Chem. Lab. Med. 37: 789, 1999.

Saez, X.: Clinical pharmacology of antibacterial agents. Pediatr. Infect. Dis. 102: 1107, 2005.

Saizou, C., Famoux, C. and Bingen, E: Severe neonatal bacterial infections. Arch Pediat.4 : 725, 2001.

Seaward, P., Hannah, M. and Myhr, T.: International multicenter term PROM study: Evaluation of infection born to patients with premature rupture of membranes at term Premature rupture of membranes. *Obstet. Gynecol.* 635 : 9, 2006.

Sheperd, C., Fisher, M. and Rosenstein, N.: Neonatal meningococcal disease in the United States. *Pediatr. Infect. Dis. J.* 20: 31, 2003.

Soman, M., Green, B. and Daling, J.: Risk factors for early neonatal sepsis. *Am. J. Epidemiol.* 121: 712, 2005.

Staehler, M., Hammer, C. and Whang, K.: Procalcitonin: a new marker for differential diagnosis of acute rejection and bacterial infection. *Clin. Intensive Care.* 5 : 82, 2005.

Steinwald, P., Becker, K. and Nylen, E.: Hyperprocalcitoninemia of E-coli sepsis associated with hypocalcaemia and hyperphosphatemia. *J. Lab. Clin. Med.* 95: 362, 1996.

Stoll, B.J., Gordon, T. and korones, S.B.: Early-onset sepsis in very low birth weight neonates. *J. Pediatr.* 129: 720, 1996.

Surpine, E., Camus, C. and Gacouin, A.: Procalcitonin: a valuable indicator of infection in a medical ICU. *Intensive Care Med.* 2 : 89, 2005.

Tausch, W., Bellard, R. and Avery, L.: Neonatal sepsis: diagnosis, treatment and management. *Pediatr. Infect. Dis. J.* 5: 438, 1998.

Tollner, U.: Early diagnosis of septicemia in newborn: clinical studies and sepsis score. *Eur. J. Pediatr.* 33:456, 2004

Turner,D., Hammerman,C. and Rudensky,B.: The role of procalcitonin as a predictor of nosocomial sepsis in preterm infants. *Acta paediatrica*.95: 1571, 2006.

Van Rossum, A.M., Wulkan, R.W. and Oudesluys- Murphy, A.M.: Procalcitonin as early marker of infection in neonates and children. *Lancet. infect. Dis.* 4: 620, 2004.

Vazzalwar, R., Pina- Rodrigues, E. and Puppala, B. : Procalcitonin as a screening test for late- Onset sepsis in preterm very low birth weight infants. *J. perinatol.* 25: 397, 2005.

Vervloet, M.G., Thijs, L.G. and Hack, C.E.: Derangements of coagulation and fibrinolysis in critically ill patients with sepsis and septic shock. *Eur. J. Pediatr.* 24: 33, 2003.

Vesikari, T., Janas, M. and Gronoos, P: Neonatal septicaemia. *Arch Dis. Child.* 60: 542, 1986.

Wilding, E., Push, F. and Zimpfer, A.: Procalcitonin is elevated in patients after severe injury. *Intensive Care Med.* 11: 562, 2005.

Yoder, M.: Clinical Gram-positive sepsis: does it fundamentally differ from Gram-negative bacterial sepsis. *J. Pediatr.* 18 : 340, 2001.

Zeni, F., Viallon, A. and Assicot, M: Procalcitonin serum concentrations and severity of sepsis. *Intensive Care Med.*21: 994, 2004.