

LITERATURE CITED

- Abd El- Sattar, A.A. (1992). Breeding studies on maize (*Zea may* L.).
Ph.D. Thesis, Fac. Agirc., Moshtohor, Zagazig Univ. Egypt.
- Abo- Dheef, M.H. (1987). Physiological genetic studies on maize
(*Zea may* L.).
M. Sc. Thesis, Fac. Agric. Ain Shams, Univ., Egypt.
- Abo- El- Fadhl, M.A. (1978). Heritability estimates of some
characteristics in maize (*Zea may* L.).
Ph.D. Thesis, Fac. Agirc., Ain Shams Univ., Egypt.
- Akhtar, S.A.; T.P. Singh, (1981). Heterosis in varietal cross of
maize.
Moderas Agric. J. 68: 27- 51 (c.f. Plant Breed Abst. 25: 9169).
- Allard, R.W. (1960). Principles of Plant Breeding .
John, Wiley & Sons, New York.
- Altinbas, M. (1996). Heterosis and combining ability in maize for
grain yield and some plant characters.
(C.f. Plant Breed Abst. 66 No. 5. 4716).
- Badr, M.M. (1989). Breeding studies on some divergent maize
populations.
M. Sc. Thesis, Fac. Agric. Moshtohor, Zagazig Univ., Egypt.
-

- Beck, D. I. ; S.K. Vasal and J. Crossa (1991). Heterosis and combining ability among subtropical and temperate intermediate- maturity maize germplasm. *Crop Sci.*, 31: 68: 73.
- Borov, M. (1966). General combining ability of inbred maize lines. Omsk Agric. Inst. 61. Pp. 14- 20, 1966. (c.f. *Plant Breed Abst.* 2, 4924, 1967).
- Dhillon, B.S. and J. Singh (1976). Diallel analysis of yield and other traits of maize varieties .
SABRAO J. 8, (2), 147- 152.
- Dhillon, B.S. and J. Singh (1977). Combining ability and heterosis in diallel crosses of maize.
Theoretical and Applied Genetics, 49 (3): 117- 122.
- Duncan, W.G. (1971). Leaf angle, leaf area and canopy photosynthesis.
Crop Sci., 11: 482- 485.
- El- Absawy, E.A. (1990). Comparison between some experimental genetic designs in two populations of maize.
Ph.D. Thesis, Fac. Agric., Moshtohor, Zagazig Univ., Egypt.
- El- Hennaway, M.A. (1980). Morphological and physiological studies on some inbred lines and their hybrids of corn (*Zea mays* L.).
M. Sc. Thesis, Fac. Agric. Al- Azhar Univ., Egypt.
-

- El- Hosary, A.A. (1985). Study of combining ability in some top crosses in maize. Egypt. J. Agron., 10(1-2): 39- 47.
- El- Hosary, A.A (1986). Estimation of genetic variance in open pollinated variety with reference to expected gain from selection.
Egypt. J. Agron. 11 (1-2): 119- 128.
- El- Hosary, (1987). Genetic studies in a synthetic variety of maize.
Egypt. J. Agron. 12 (1-2): 57- 64.
- El- Hosary, A.A. (1988). Heterosis and combining ability of ten maize inbred lines as determined by diallel crossing over two planting dates.
Egypt. J. Agron. 13 (1-2): 13- 25.
- El- Hosary, A.A. (1988a). An analysis of the combining ability of inbred lines of maize ((*Zea may* L.) in diallel cross system.
Egypt. J. Agron. 13 (1-2): 27- 29.
- El- Hosary, A.A (1988b). Evaluation of twenty new inbred lines by top crosses in corn (*Zea may* L.).
Proc. 3rd Egyptian Conf. Agron. Kafr El- Sheikh, Vol. 1: 48- 56.
- El- Hosary, A.A (1989). Heterosis and combining ability of six inbred lines of maize in diallel crosses over two years.
Egypt. J. Agron. 14 (1-2): 47- 58.
-

- El- Hosary, A.A; A.A. Sedhom and S.A. Mohamed (1989).
Correlation and path coefficient studies in maize.
Annals of Agric. Sc., Moshtohor, 27 (3):
- El- Hosary, A.A. and S.A. Sedhom (1990). Diallel analysis of yield
and other agronomic characters in maize (*Zea may* L.).
Ann. of Agric. Sc. Moshtohor 28(4):
- El- Hosary, A.A; G.A. Sary and A.A. Abd El- Sattar (1990). Studies
on combining ability and heterosis in maize (*Zea may* L.). II-
Yield and yield components.
Egypt. J. Agron. 15 (1-2): 9- 22.
- El- Hosary, A.A; G.A. Sary and A.A. Abd El- Sattar (1990a). Studies
on combining ability and heterosis in maize (*Zea may* L.). I-
Growth attributes.
Egypt. J. Agron. 15 (1-2): 23- 34.
- El- Hosary, A.A.; M.K. Mohamed and S.A. Sedhom (1994).
Performance and combining ability in diallel crosses of maize.
Annals of Agric. Sc., Moshtohor, 32(1): 203- 215.
- El- Ítriby, H.A. (1986). Genotyping of parental inbred lines: Its
utilization in the quality control of hybrid seed production.
17 th Annual Meeting of the Egyptian Society of Gene. And
Cyto. 26- 27, March.
-

El- Marakby, A.M. (1964). A study of correlation between yield and some morphological characters in maize.

M. Sc. Thesis, Fac. Agric. Ain Shams Univ., Egypt.

El- Rouby, M.M. and Galal (1972). Heterosis and combining ability in variety crosses of maize and their implications breeding schemes.

Egypt. J. Genet. Cytol. 1: 270- 279.

El- Rouby, M.M; A.M. Gad and R.M. Abdallah (1979). Components of genetic variance and their interactions with locations and years in maize.

Egypt. J. Genet. Cytol. 8: 107- 121.

El- Zeir, F.A.A. (1984). Combining ability and heterosis of some inbred lines of maize.

M. Sc. Thesis, Fac. Agric. Zagazig Univ., Egypt.

Erakey, A.G.; A.H. Salem and A.S. Gouda (1983). The interrelationship between growth and yield attributes in maize. 2- Path coefficient and regression for yield attributes in maize .

Egypt. Soc. Of Crop Sci. Proceeding of the first conference of Agron 1 (A) Cereal Crops 95, 126.

Galal, A.A.; F.A. El- Zeir; M.A. Younis and F.M. Omar (1990). Heterosis and combining ability in two sets variety cross diallel in maize.

J. Agric. Res., Tanta Univ., 16 (3): 412- 421.

Gerrish, E.E. (1983). Indication from a diallel study for intervarital maize hybridization in the corn Belt.

Crop Sci., 23: 1082- 1084.

Girffing, B. (1956). Concept of general and specific combining ability in relation to diallel crossing systems.

Aust. J. Bio. Sci., 9: 463- 493.

Gomes, E. (1996). Combining ability among fifteen early cycle maize population in Brazil.

(c.f. Plant Breed Abst. Vol. 66 N. 5, 4717, 1996).

Hajkham, A. (1972). Combining ability in maize (*Zea may* L.).

In Proceeding of the 21 th . 22 nd Pakistan Sci. Conf. Rajshahi, Part III (Abst.)(c.f. Plant Breed Abst. 42 (3), 4777, 1972).

Hassaballa, E.S. ; M.A. El- Morshidy; M.A. Khalifa and M.M. Abdallah, (1980). Genetical analysis of diallel crosses in maize. II- Plant and ear height.

Res. Bull., Fac. Agric. Ain Shams Univ., No. 1294.

Kalsy, H.S. and D. Sharma (1970). Study of genetic parameters and heterotic effects in crosses of maize (*Zea may* L.). varieties with varying chromosome knob numbers .

Euphytica, 19: 522- 530.

Krolikowski, Z. (1973). Assessment of general and specific combining ability in a diallel system.

Binlety instytutu Hodowli Aklimaty Zacji Roslin (314), 39- 47.
(c.f. Plant Breed Abst. 44, 8, 5114, 1974).

Kumar, P. (1994). Heterosis in varietal crosses of maize (*Zea may* L.).

Neo Botanica 2(1): 15- 19.(c.f. Plant Breed Abst. 65, 2510, 350, 1995).

Luchsinger, L.A.; M.A. Violic and B.R. Barnier (1971). General and specific combining ability for yield and its components in ten lines of maize (*Zea may* L.).

Agric. Tencnia, 31,(3), 144- 150. (c.f. Plant Breed Abst. 43 (4), 2633)

Mahajan, V. and A.S. Khehra (1992). Inheritance of quantitative traits in maize (*Zea may* L.). in winter and monsoon season, Indian J. of Genet. & Plant Breed 51 (3); 292- 300. (c.f. Plant Breed Abst. 64, 333, 144, 1994).

Mahmoud, A.A. (1989). Genetic studies through diallel of maize inbred lines.

M. Sc. Thesis, Fac. Agric. Cairo Univ.

Makee, G.W. (1964). A coefficient for computing leaf area in hybrid corn.

Agron. J. 56: 240- 241.

- Manoliu, M. (1969). Studies on the heritability of some quantitative characters in maize.
- Lucrari. Stuntifice, Institute Agronomic "N. Balcescu" 12: 35-43.(c.f. Plant Breed Abst. 43, 9, 6808, 1973)
- Mariani, G. and E. Desiderio (1975). Grain yield and prolificacy in maize . III- A diallel analysis of eight inbreds from the open population variety. Nostrano delle soland two densities. Maydica, 20 (2): 95- 110.(c.f. Plant Breed Abst. 46, 5147, 1976).
- Mason, L. and S.Z. Marcus (1976). Diallel analysis of maize for leaf angle, leaf area, yield and yield components.
- Crop Sci., 16: 693- 696.
- Merdith, W.R. (1984). Quantitative genetics. P. 131- 150. In R.J. Kohel and C.F. Lewised. Colton Agronomy Monar. A.S. CSSA and SSA Madison, WI.
- Mohamed, A.A. and M. Yasien (1987). Evaluation of thirty one inbred lines of maize (*Zea may* L.).
- Annals of Agric. Sc., Moshtohor, 25(1): 105- 117.
- Mohamed, S.A.S. (1979). Genetic studies on corn (*Zea may* L.). .
- M. Sc. Thesis, Fac. Agric. Al- Azhar Univ., Egypt.
- Mohamed, S.A.S. (1984). Studies on the genetic basis for heterosis in corn (*Zea may* L.).
- Ph.D. Thesis, Fac. Agirc., Al- Azhar Univ., Egypt.
-

Mohamed, M.K. and S.A. Sedhom, (1994). A comparison between four statistical procedures of relating yield components in a set of corn varieties.

Annals of Agric. Sc., Moshtohor, 31(4); 1855- 1866.

Mokbel, H.A.A. (1988). Genetical studies on maize. Estimation of the genetic variation for nine maize quantitative characters by diallel analysis .

M. Sc. Thesis, Fac. Agric. Alex. Univ., Egypt.

Moll, R.H.; W.S. Salhuana, and H.F. Robinson (1962). Heterosis and genetic diversity in variety crosses of maize.

Crop Sci., 2: 197- 198.

Morshed G.A.; N.N. El- Hefnawy, and A.A. Galal (1990). Estimation of combining ability and heterosis effects under different environmental in maize (*Zea may* L.).

Minofiya J. Agric. Res. 15(2): 1509- 1517.

Motawea, M.H. (1997). Genetic studies on maize (*Zea may* L.).

Ph.D. Thesis, Fac. Agric., Moshtohor, Zagazig Univ., Egypt.

Mourad, S.S.B. (1978). Study of combining ability in some lines of maize (*Zea may* L.).

M. Sc. Thesis, Fac. Agric., Al- Azhar Univ. Egypt.

Mourthy, A.R. and A. Ramamurthy (1981). Diallel analysis of yield and maturity components in maize.

Indian J. Genet. Plant Breed., 41(1), 30- 33.

Mufti, M.N. and H.R. Rao, (1995). Quantitative studies on heterosis in *Zea may* L. diallel crosses .

Sarhad J. Agric. 11(1); 71- 77. (c.f. Plant Breed Abst. 66, 3668, 497, 1996).

Naspolini, F.; V.E.E. Gama; R.T. Vianna, and J.R. More (1981). General and specific combining ability for yield in a diallel cross among 18 maize populations (*Zea may* L.).

Revista Brasileirde Gent. 4 (4): 571- 577. (c.f. Plant Breed Abst. 52, 6495, 1982).

Nawar, A.A. (1985). Diallel analysis of the combining ability of inbred lines and utilization inbreeding maize hybrids. Minofyia J. Agric., Res. 10 (4): 2015- 2026.

Nawar, A.A.; M.E. Gomaa and M.S. Rady (1980). Heterosis and combining ability in maize.

Egypt. J. Genetic. Cytol., 9: 255- 267.

Nawar, A.A. and A.A. El- Hosary (1985a). A comparison between two experimental diallel crosses designs .

Minofya J. Agric. Res. 10(4): 2029- 2039.

Nawar, A.A. and A.A. El- Hosary (1985). General vs. specific combining ability in top crosses of maize (*Zea may* L.).

2nd Agric. Conf. Bot. Sci., 221- 232.

- Nawar, A.A.; F.A. Hendawy; H.A. Dawwam and A.A. El-Hosary (1988). Study of the reciprocal effects on the performance of double cross hybrid under two nitrogen levels in maize. Proc. 3rd Egypt. Conf. Agron. 1: 1-13.
- Nawar, A.A.; F.A. Hendawy; S.A. Shamarka; A.M. Shehata and M.M.M. Asfour (1995a). Estimation of some genetical parameters in a composite cultivar. Menofiya J. of Agric. Res., 20: 463- 482.
- Nawar, A.A.; F.A. Hendawy; S.A. Shamarka; A.M. Shehata and M.M.M. Asfour (1995b). Genetic variance components in Giza 2 maize populations. Menofiya J. of Agric. Res., 20: 423- 439.
- Paterniani, E. and J.H. Lonnquist (1963). Heterosis in intervarietal crosses of corn (*Zea mays* L.). Crop Sci., 3: 504- 507.
- Pena O.M.G. and V.J.L. Arellano (1991). Effect of inbreeding depression on grain yield and its components in the maize hybrid H 131, in two environments. Revista Chapingo, 15: 71- 72, 12- 16. (c.f. Plant Breed. Abst.63, 12844, 1656, 1993).
- Ragheb, M.M. (1985). Physiological explanation of heterosis in maize (*Zea mays* L.). Ph.D. Thesis, Fac. Agric., Al Azhar Univ., Egypt.
-

- Ragheb M.M; A.A. Bedeer; A.S.H. Goda; A. Abo- El-Saad, and S.H.F. Abdel Aziz (1993). Phenotypic stability parameters of white maize hybrids evaluated under different environmental conditions.
Zagazig J. Agric. Res. 5: 1447- 141.
- Ramamurthy, A. (1980). Genetic analysis of some quantitative characters in maize (*Zea may* L.).
Mansoura J. Agric. Sci., 14, 1, 126.
- Rashed, M.A. (1977). Genetic studies on corn (*Zea may* L.).
M. Sc. Thesis, Fac. Agric. Ain Shams Univ., Egypt.
- Reddy, K.H.P. and B.D. Agrawal (1992). Estimation of genetic variation in an improved population of maize (*Zea may* L.).
Madras, Agric. J. 79: 714- 719.(c.f. Maize Abst. 1994, 10 (6) 3347, p. 444).
- Salama, F.A.; H. Gado; M. El- Goda and S.E. Sadek (1994). Correlation and path coefficient analysis in eight white maize (*Zea may* L.). hybrid characters .
Minofyia J. Agric. Res., 19(6) 3009- 3020.
- Salem, A.H.; A.A. Galal and F.A. El- Zeir (1986). Diallel analysis of combining ability in maize under different environments.
Egypt. J. Genet. Cytol. 15: 231- 139.
- Sanghi, A.K.; K.N. Agrwal and M.I. Qadri, (1982). Gene effects and heterosis for grain yield and ear traits in miaze.
-

- Indian J. Gene. and Plant Breed., 42(3); 360- 363.
- Sary, G.A.; A.A. El- Hosary, S.A. Mohamed and A.A. Abdel- Sattar (1990). Studies on combining ability and heterosis in maize (*Zea may* L.). III- Association studies. Egypt. J. Agron., 15 (1-2): 1-8.
- Sedhom, S.A. (1994). Estimation of general and specific combining ability in maize under different planting dates. Annals of Agric. Sc., Moshtohor, 32(1): 119- 130.
- Shaher, A.G. (1982). Effect of planting and nitrogen fertilization on growth and yield of some varieties. M. Sc. Thesis, Fac. Agric. Zagazig Univ., Egypt.
- Shieh, G.J. (1994). The effects of crop season on the heterosis of agronomic characteristics in maize . J.Agric. Res. Of Ehina, 43 (4): 345- 353 . (c.f. Plant Breed Abst. 66, 5900, 804, 1996).
- Singh, I.S. and V.L. Asnani (1979). Combining ability analysis for yield and some yield components in maize. Indian j. Genet. Plant Breed., 31: 322- 332.
- Snedecor, and W.G. Cockran (1967). Statistical methods. 6th Edition. The Iowa State Univ. Press, Ames Iowa, U.S.A.
- Sotcenko, V.S. (1970). Influence of external conditions on variation in general and specific combining ability in maize lines. Doklvses Akad, Sel' Shohoz, Nauk Lenin, Kepatt. Mnlenin. Acad Agric. Sci., 2: 14- 16. Russian.(c.f. Plant Breed Abst. 40, 7388).

- Sprague, G.F. and W.I. Thomas, (1967). Further evidence of epistasis in single and three way cross yields of (*Zea mays* L.). Crop Sci., 7: 355- 356.
- Stubber, C.W.; R.H. Moll and W.D. Hanson (1966). Genetic variance and interrelationship of six traits in a hybrid population of (*Zea mays* L.). Crop Sci., 6: 455- 458.
- Surinder, K.; S. Ganesan, C. Jose and B. David (1992). Heterosis and combining ability of CIMMYT'S subtropical and temperate early maturity maize Germplasm. Crop Sci., 32 (4) 884- 890.
- Tkachenko, I.K. and A.A. Goncharenko, (1973). Comparative study of general combining ability of maize lines on basis of reproductive and vegetative characters. Selskohozyaistvennaya. Biologiya 816: 823- 829. (c.f. Plant Breed Abst. 44, 7, 4357, 1974).
- Turgut, I.; S. Yuce and M. Altinbas (1995). Inheritance of some agronomic traits in a diallel cross of maize inbreds. II- Grain yield and its components. Anadolu, 5 (1): 74- 92.(c.f. Plant Breed Abst. 65, 12612, 1769, 1996).
- Verma, R.K. and T.P. Singh (1980). Heterosis and combining ability studies for certain quantitative traits in pop corn. Mansoura J. Agric., Sci. 14: 15- 17.
- Wright, S. (1921). Correlation and causation. J. Agric. Res. 20: 557- 585.

Wright, S. (1923). The theory of path coefficient .

Genetics, 8: 239- 285.

Wright, S. (1934). The method of path coefficients.

Ann. Math. Stat. 5: 161- 215.

Yasien, M. (1977). Morphological and physiological manifestation of combining ability .

M. Sc. Thesis, Fac. Agric. Ain Shams Univ. Egypt.

Yasien, M. (1982). A genetic study of some physiological and morphological attributes in maize.

Ph.D. Thesis, Fac. Agric., Ain Shams, Univ.