

REFERENCES

- Abd El- Sattar, A.A. (1986). Studies on combining ability and heterosis in maize (*Zea mays* L.) M.Sc. Thesis, Fac. of Agric., Moshtohor, Zagazig Univ., Egypt.
- Abd El- Sattar, A.A. (1992). Breeding studies on maize (*Zea mays*, L.). Ph. D. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ. Egypt.
- Abo- Dheaf, M.H. (1987). Physiological genetic studies on maize (*Zea mays*, L.) M.Sc. Thesis, Fac. of Agric. Ain Shams Univ., Egypt.
- Allard, R.W. (1960). Principles of Plant Breeding. John Wiley & Sons, New York.
- Altinbas, M. (1995). Heterosis and combining ability in maize for grain yield and some plant characters. *Anadolu*, 5 (2): 35 - 51. (c. f. Plant Breed. Abst. 66, 4716, 643, 1996).
- Alvarez, A.; G. Garay; J. Gimenez and J.I. Ruiz DE Galarreta (1993). Heterosis in two synthetic varieties of maize based on morphological and reproductive traits. *Investigation Agraria, Production y Proteccion Vegetales*, 8 (3): 333- 340. (c. f. Plant Breed. Abst., 65, 2564, 355, 1995).
-

- Atia, A.E.M. (1986). Breeding studies on maize. M.Sc. Thesis, Fac. of Agric. Menoufia Univ., Egypt.
- Badr, M.M. (1989). Breeding studies on some divergent maize populations. M.Sc. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ., Egypt.
- Beck, D.L.; S.K. Vasal and J. Crossa (1991). Heterosis and combining ability among subtropical and temperate intermediate maturity maize germplasm. *Crop sci.*, 31: 68- 73.
- Borov, M. (1966). General combining ability of inbred maize lines. *Omsk Agric, Inst.* 61: pp. 14- 20, 1966 (c. f. *Plant Breed. Abst.*, 2, 4929, 1967).
- Brenner, D.; F.I. DE. Carvalho and L.C. Federizzi (1991). Estimates of combining ability for grain yield in maize (*Zea mays*, L.) populations. *Revista Brasileira de Genetica.* 14(4): 1001 - 1009 (c. f. *Plant Breed. Abst.*, 63, 2584, 314 - 315, 1993).
- Buhinicek, I.; D. Vasilj; R. Pavlina and B. Palaversic (1994). Estimates of GCA and SCA based on Griffing's methods 2 and 4 for maize lines and hybrids in vitro and in situ. *Poljoprivredna Znanstvena Smotra.* 59 (4): 357- 367. (c. f. *Plant Breed. Abst.*, 65 (5), 4903, 685, 1995).
- Cao, L.J.; J.H. Liang; F.C. Wang; Y.F. Zhang and S. Liu (1994). Heritability analysis of major characters in silage maize. *Grassland of China*, 2: 36 - 40. (c.f. *Plant Breed. Abst.*, 65 (5), 4937, 689, 1995).

- Choi, Keunjin, C. Moonup; P.K. Yong; L. Hongseok; S. Jongho and S.D. Young (1995). Maize Genetics Cooperation Newsletter. 69, 122- 123. (c.f. Plant Breed. Abst., 66, 10162, 1401, 1996).
- Damborsky, M.; O. Chloupek and J. Ehrenbergerova (1994). Variability of maize lines and diallel cross hybrids. Genetika a Slechteni. 30 (4); 297- 303. (c.f. Plant Breed. Abst., 65 (5) 4938, 689, 1995).
- Danial, L. (1969). Heterosis and mean degree of dominance in the inheritance of plant height in maize (*Zea mays*, L.) Z. Pflanzenz, 61: 366- 380. (c.f. Plant Breed. Abst. 40: 2911, 372, 1970).
- Dawod, K.M. and A.S.A. Mohamed (1989). Estimates of heritability and path coefficient analysis for some characters in maize (*Zea mays*, L.) Mesopotamia J. Agric., 12 (4): 243- 254.
- Debnath, S.C. (1984). Heterosis in maize : 1- Grain yield and some of its attributes. Bangladesh J. Agric. Res., 9: 78 - 83 (c. f. Plant Breed. Abst., 55: 6780, 1985).
- Dhillon, B.S. ; U.K. Bansal and W.R. Kapoor (1994). Variability for shelling in maize. Crop Improvement, 12 (1- 2): 87- 88. (c. f. Plant Breed. Abst., 66, 2562, 348, 1996).
- El- Enany, M.A.M. (1991). Biochemical genetic studies on salt tolerance in maize. M.Sc. Thesis, Fac. of Agric. Ain Shams Univ., Egypt.
-

- El- Hosary, A.A. (1985), Study of combining ability in some top crosses in maize. Egypt. J. Agron., 10 (1-2): 39- 47.
- El- Hosary, A.A. (1986). Estimation of genetic variance in open pollinated variety with reference to expected gain from selection. Egypt. J. Agron., 11 (1- 2): 119- 128.
- El- Hosary, A.A. (1987). Genetic studies in a synthetic variety of maize. Egypt. J. Agron., 12 (1- 2): 57- 64.
- El- Hosary, A.A. (1988a). Heterosis and combining ability of ten maize inbred lines as determined by diallel crossing over two plant dates. Egypt. J. Agron., 13 (1- 2): 13 - 25.
- El- Hosary, A.A. (1988b). An analysis of the combining ability of inbred lines of maize (*Zea mays*, L.) in diallel cross system. Egypt. J. Agron., 13 (1-2): 27- 39.
- El- Hosary, A.A. (1988c). Evaluation of twenty new inbred lines by top crosses in corn (*Zea mays*, L.). Proc. 3 th Egyptian Conf. Agron., Kafr El- sheikh, Vol. 1: 48- 56.
- El- Hosary, A.A. (1989). Heterosis and combining ability of six inbred lines of maize in diallel crosses over two years. Egypt. J. Agron., 14 (1- 2): 47 - 58.
- El- Hosary, A.A.; G.A. Sary and A.A. Abd El- Sattar (1990a). Studies on combining ability and heterosis in maize (*Zea mays*, L.). II. Yield and yield components. Egypt. J. Agron., 15 (1- 2): 9 - 22.
-

- Griffing, B. (1956). Concept of general and specific combining ability in relation to diallel crossing system. *Australian J. Biol. Sci.*, 9: 463- 493.
- Han, G.C. and A.R. Hallauer (1989). Estimates of genetic variability in F_2 maize population. *J. Iowa Acad. Sci.*, 96: 14 - 19.
- Hassaballa, E.S.; M.A. El- Morshidy; M.A. Khalifa and M.M. Abdalla (1980). Genetical analysis of diallel crosses in maize. II. Plant and ear height. *Res. Bull., Fac. Agric. Ain Shams Univ.*, No. 1294.
- Hayman, B.I. (1954 a). The analysis of variance of diallel tables. *Biometrics*, 10: 235 - 244.
- Hayman, B.I. (1954 b) The theory and analysis of diallel crosses. *Genetics*, 39: 789- 809.
- Hayman, B.I. and K. Mather (1955). The dissection of gene interaction in continuous variation. *Biometrics*, 10: 69- 82.
- Ivanov, S. and N. Tomov (1995). Combining ability for yield components in high- lysine maize lines homozygous for the opaque-2 gene. *Rasteniev "dni Nauk*; 32: 120 - 123. (c. f. *Plant Breed. Abst.*, 66, 1475, 203, 1996).
- Jenkins, M.T. (1934). Methods of estimation the performance of double crosses in corn. *J. Amer. Soc., Agron.*, 26: 199 - 204.
- Jinks, J.L. (1954). The analysis of continuous variation in a diallel cross of *Nicotiana rustica*. *Genetics*, 39: 767- 88.
- Jinks, J.L. (1955). A survey of the genetical basis of heterosis in a variety of diallel crosses. *Heredity*, 9: 223- 238.
-

Jinks, J.L. and B.I. Hayman (1953). The analysis of diallel crosses. Maize genetics Coop. Newsl. 27: 48- 54.

kalsy, H.S. and D. Sharma (1970). Study of genetic parameters and heterotic effects in crosses of maize (*Zea mays*, L.) varieties with varying chromosome Knob numbers. Euphytica 19: 522-530.

Kassem, E.S.; M.Z. El- Hifny; M.A. El- Morshidy and K.A. Kheiralla (1979). Genetic analysis of maize grain yield and its components by diallel crossing. Egypt. J. Agron., 4(1): 125-132.

Kravenchko, V.A.; M.C. Korolchak and G.A. Morshed (1971). Heterosis in maize in respect of some economically valuable characters of after interline crossing . Nauk Pratisukr. Sel Skogopod, Akad., 40: 47 - 49. (c. f. Plant Breed. Abst., 45 (8): 6311, 1975).

Krolikowski, Z. (1969). Heterosis and combining ability of four inbred lines of maize in diallel crosses. Genet. Polon, 10: 104-109.

Krolikowski, Z. (1973). Assessment of general and specific combining ability of maize inbred lines crosses in a diallel system. Biuletyn Instytutu Hodowli i Aklimatyzacji Roslin 39 47. (c. f. Plant Breed. Abst., 44, 8, 5114, 1974).

- Kumar, P. (1994). Heterosis in varietal crosses of maize (*Zea mays*, L.). Neo Botanica 2(1): 15 - 19. (c. f. Plant Breed. Abst. 65, 2510, 350, 1995).
- Landi, P.; A. Camussi; A. Verderio and C.Lorenzoni (1986). Combining ability of early maize (*Zea mays*, L.) inbred lines and evaluated under conditions of double cropping. Maydica, 37: 369 - 377 (c. f. Plant Breed. Abst. 57, 4870).
- Lee, W. and B. Choe (1995). Characteristics and combining ability of Korean local waxy maize inbreds and hybrids. Korean J. of Crop Sci., 40 (2): 175 - 184. (c. f. Plant Breed. Abst. 66, 1471, 203, 1995).
- Mahajan, V. and A.S. Khehra (1991). Inheritance of quantitative traits in maize (*Zea mays*, L.) in winter and monsoon season, Indian J of Genet. & Plat Breed. 51 (3): 292- 300. (c. f. Plant Breed. Abst. 64, 333, 144, 1994).
- Mahajan, V.; A.S. Kherhra; S.S. Pal and A.S. Gupta (1992). Combining ability of plant height and leaves gained per day in maize. Crop Res. (Hisar), 5 (2): 391 - 394. (c. f. Plant Breed. Abst. 64, 10267, 1422, 1994).
- Mahmoud, A.A. (1989). Genetic studies through diallel of maize inbred lines . M.Sc. Thesis Fac. of Agric. Cairo Univ. Egypt.
- Major, D.J.; R.B. Hunter; L.W. Kannenberg; T.B. Daynard and J.W. Tanner (1972). Comparison of inbred and hybrid corn grain
-

yield measured to equal leaf area index. *Canad. J. Pl. Sci.*, 52: 315 - 319.

Mather, K. and J.L. Jinks (1971). *Biometrical genetics* (2nd edition). Chapman and Hall, Ltd., London.

Mohamed, A.A. and M. Yasien (1987). Evaluation of thirty- one inbred lines of maize (*Zea mays*, L.). *Ann. of Agric. Sci.*, Moshtohor, 25 (1): 105 - 117.

Mohamed, S.A.S. (1979). Genetic studies on corn (*Zea mays*, L.). M.Sc. Thesis, Fac. of Agric. Al- Azhar Univ, Egypt.

Mohamed, S.A.S. (1984). Studies on the genetic basis for heterosis in corn (*Zea mays*, L.). Ph. D. Thesis, Fac. of Agric. Al- Azhar Univ. Egypt.

Mokbel, H.A.A. (1988). Genetical studies on maize. Estimation of the genetic variation components for nine maize quantitative characters by diallel analysis. M.Sc. Thesis, Fac. of Agric. Alex. Univ. Egypt.

Mourad, S.S.B. (1978). Study of combining ability in some lines of maize (*Zea mays*, L.). M.Sc. Thesis, Fac. of Agric. Al- Azhar Univ. Egypt.

Mufti, M.U. and H.R. Rao (1995). Quantitative studies on heterosis in *Zea mays*, L. diallel crosses. *Sarhad J. of Agric.*, 11 (1): 71 - 77 (c. f. *Plant Breed. Abst.*, 66, 3668, 497, 1996).

- Mukherjee, B.K.; N. L. Dhawan and N.N. Sing (1971). Heterosis in intervarietal crosses in maize (*Zea mays*, L.). Indian J. Agric. Sci., 41: 489- 494.
- Nawar, A.A. (1985a). Diallel analysis of the combining ability of inbred lines and utilization in breeding maize hybrids. Menoufia J. of Agric. Res., 10 (4): 2015 - 2026.
- Nawar, A.A. (1985b). Genetic variance components and their implications in yield improvement in maize. Communications in Science and Development Res., 12: 83- 93.
- Nawar, A.A. and A.A. El- Hosary (1985a). A comparison between two experimental diallel crosses designs. Menoufia J. of Agric. Res., 10 (4): 2029 - 2039.
- Nawar, A.A. and A.A. El- Hosary (1985b). General vs. specific combining ability in top crosses of maize (*Zea mays*, L.) 2nd Agric. Cond. Bot. Sci., 221- 232.
- Nawar, A.A.; A.A. Abul- Naus and M.E. Gomaa (1981). Heterosis and general vs specific combining ability inbred lines of corn. Egypt. J. Genet. Cytol., 10: 19 - 29.
- Nawar, A.A.; M.E. Gomaa and M.S. Rady (1980). Heterosis and combining ability in maize. Egypt. J. Genet. Cytol., 9: 255 - 267.
- Nawar, A.A.; M.S. Rady and A.A. El- Hosary (1984). Genotypic variability and correlation coefficients of some quantitative
-

- characters in Egyptian maize (*Zea mays*, L.). Menoufia J. of Agric. Res., (8): 123- 139.
- Nawar, A.A.; F.A. Hendawy; S.A. Shamarka; A.M. Shehata and M.M.M. Asfour (1995a). Estimation of some genetical parameters in a composite cultivar. Menoufia J. of Agric. Res., 20: 463- 482.
- Nawar, A.A.; F.A. Hendawy; S.A. Shamarka; A.M. Shehata and M.M.M. Asfour (1995b). Genetic variacne components in Giza 2 maize populations. Menoufia J. of Agric. Res., 20: 423- 439.
- Odiemah, M.A. (1973). Estimation of general and specific combining ability in maize under different environments. M.Sc. Thesis, Fac. of Agric. Alex. Univ. Egypt.
- Odiemah, M. A. (1992). Estimates of genetic parameters for yield of six maize inbreds and their crosses in low and high plant densities. Acta Aronomica Hungarica, 41 (1- 2): 85 - 92. (c.f. Plant Breed. Abst. 63, 11765, 1505, 1993).
- Paterniani, E. (1980). Heterosis in intervarietal crosses of maize (*Zea mays*, L.) and their advanced generations. Revista Brasileira de Genetica, 235 - 249. (c. f. Plant Breed. Abst. 51, 7040).
- Pena O.M.G. and V.J.L. Arellano (1991). Effect of inbreeding depression on grain yield and its components in the maize hybrid H- 131, in two environments. Revista Chapingo, 15, 71 - 72, 12 - 16. (c. f. Plant Breed. Abst. 63, 12844, 1656, 1993).
-

- Qin, T.C. and Z.G, Li (1991). Studies of the inheritance of kernel growth characters and their relation to yield characters in maize (*Zea mays*, L.) . Acta Agronomica Sinica, 17 (3): 183 - 191. (c. f. Plant Breed. Abst., 63, 2579, 314, 1993).
- Rashed, M.A. (1977). Genetic studies on corn (*Zea mays*, L.). M.Sc. Thesis, Fac. of Agric. Ain Shams Univ., Egypt.
- Reddy, K.H.P. and B.D. Agrawal (1992). Estimation of genetic variation in an improved population of maize (*Zea mays*, L.) Madras Agric. J. 79: 714- 719. (c. f. Maize Abst. 1994 Vol. 10, No. 6, 3347, p. 444).
- Salem, A.H.; A.A. Galal and F.A. El- Zeir (1986). Diallel analysis of combining ability in maize under different environments. Egypt. J. Genet. Cytol., 15: 231- 239.
- Sanghi, A.K.; K.N. Agrwal and M.I. Qadri (1982). Gene effects and heterosis for grain yield and ear traits in maize. Indian J. Genet. and Plant Breed., 42 (3): 360- 363.
- Sedhom, S.A. (1984). Study on late wilt disease resistance in maize. Ph. D. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ., Egypt.
- Sfakianakis, J.; N. Fotiadis; G. Evgenidis and N. Katranis (1996). Genetic analysis of maize variety diallel crosses and related populations. Maydica, 41 (2): 113- 117. (c. f. Plant Breed. Abst. 66, 11323, 1547, 1996).
-

- Shieh, G.J. (1994). The effects of crop season on the heterosis of agronomic characteristics in maize. *J. Agric. Res. of China*, 43 (4): 345- 353. (c. f. *Plant Breed.*, Abst. 66, 5900, 804- 805, 1996).
- Singh, I.S. and V.L.Asnani (1979). Combining ability analysis for yield and some yield components in maize. *Indian J. Genet. Plant. Breed.*, 31: 322- 332.
- Singh, R.K. and B.D. Chaudhary (1977). Biometrical method in quantitative genetic analysis. India Published by USHA. Rajkumar for Kalyani Publishers.
- Singh, S.N.; K.N. Singh and H.G. Singh (1989). Combining ability in varying conditions of soil in maize. *Crop Improvement*, 16 (2): 150 - 153 (c. f. *Plant Breed. Abst.*, 62(2), 1259, 167, 1992).
- Sotcenko, V.S. (1970). Influence of external conditions on variation in general and specific combining ability in maize lines. *Dokl. Vses Akad. Sel' Shohoz. Nauk. Lenin, Kep All. Un. Lenin. Acad Agric. Sci.*, 2: 14 - 16. Russian. (c. f. *Plant Breed. Abst.* 40, 7388).
- Sprague, G.F. (1964). Estimates of genetic variations of two open-pollinated varieties of maize and their reciprocal F₁ hybrids. *Crop sci.*, 4: 332- 334.
- Turgut, I.; S. Yuce and M. Altinbas (1995). Inheritance of some agronomic traits in a diallel cross of maize inbreds. II. Grain
-