

LITERATURE CITED

Abdel - Sabour M.S, Al - Koddoussi A.R, and E.E. hassan (1990) Genetic behaviour of some yield components in barley. Zagazig J. Agric. Res. Vol. 17 (1) 23:30.

Abul - Nass, A.A; A.A. El - Hosary and M.M. Sakr (1986) : Genetic studies in common wheat (*Triticum aestivum* L.) Egypt. J. Agron. Vol. 11.93-103.

.....(1991): Genetical studies on durum wheat (*Triticum durum* L.) . Egypt. J. Agron. Vol.16.81-94.

Ahmed, I.A. (1981) : Inheritance studies of some economic characters in some barley crosses. M.s.c Thesis. Fac. of Agric. Moshtohor, Zagazig, university.

Anwar, A.R. and A.R. Chowdhry (1969): Heritability and inheritance of plant height, heading date, grain yield in four spring wheat crosses. Crop Sci. 9: 760 - 761.

Bashir - Ahmed, M.A Chowdhry; Muhammod Sharif, A.S. Khan and M.L. Shaha (1984): Combining ability for grain yield and other related traits in spring wheat. J. Agric. Res. Pakistan, 2: 1-5.

Bhatti, M.S; N.U.I.Khan; M.A. Bajwa and A.G. Khan (1982) : Heterosis in sp spring wheat. J. Agric. Res. Pakistan, 2: 1-5. Bhatyia. V.J; B.S. Jadon and M.S. Pithia (1986): Genetic of quantitative characters in macaroni wheat. Indian J. Agric Sci., 56:878- 880 .

Bitzer, M.J.(1969): Hybrid vigor, combining ability and gene action in crosses

- of (*T. aestivum* L.)Diss.Abst.29:2869-91(C.F.Pl.Breed.Abst.,40:4654)
; F.L. Patterson and W.E. Nyquist (1971): Hybrid vigor and gene action
in six parent diallel crosses of soft winter.Can J. Genet. cytol.13:131-137.
-, (1972): Heterosis and combining ability in southern soft red winter
wheats. Crop Sci., 12: 35- 37.
- Blanco,A.; O.A. Tanzarella; F. Cairo and G.T.Scarscia Mugnozza (1982).
Inheritance of plant height in durum wheat, *Triticum durum*
Desf.Genetic. Agraria, 36:393- 406(C.F. Plant Breed.Abst.,53:6122.
(1983.)
- Brown, C.M;R.O Wible and R.D. Seif (1966): Heterosis and combining ability
in common winter wheat. Crop Sci., 6: 382-383.
- Crumpacker,D.W.and N.S. Urguharts (1962): Relationship of gene action and
combining ability in a set of wheat parents. Abstracts of the annual
meeting of the American Society of Agronomy held at Cornell Univ.,
Ithaca, N.Y. on August 20- Pp. 114 (C.F.Pl. Breed. Abst., 34: 1706)
- Darwish, I.H.I (1992): Breeding studies on wheat. M.Sc. Thesis Fac. of Agric.
Menofiya Univ. Egypt .
- Eisea,R.K. (1976):Effect of heterosis on some chemical components in hybrid
wheat. M.Sc. Thesis Fac. Agric. Shebin El- Kom Tanta Uni., Egypt.
- El- Khatib, M.K. (1990): Relationships between some morpho- physiological c
characters and grain yield of wheat Ph.D. Thesis Fac.of Agric.at
Moshtohor,Zagazig Univ. (Banha- branch).
- El- Shamarka, S.A. (1980): Studies on heterosis and combining ability in

- common wheat M. Sc. Thesis Fac. of Agric. Menofiya univ. Egypt.
- Fonseca, S. and F.L. Patterson (1968): Hybrid vigor in seven parental diallel cross in common wheat, *Triticum aestivum* L. Crop Sci., 8: 85- 88.
- Freeman, G.F (1919): Heredity of quantitative characters in wheat. Genetics 4 : 1 -93.
- Gandhi, S.M.; T.U. Menon; P.D. Bhavgava and M.P. Bhatnager (1961): Studies on hybrid vigor in wheat. Indian J. Genetic Pl. Breed., 21: 1- 10.
- Griffing , J.B. (1956) Concept of general and specific combining ability in relation to diallel crossing systems. Australian J. Biol., Sci., 9:463-493.
- Gyawali, K.K; C.O. Qualset and W.T. Yamazaki (1968): Estimates of heterosis and combining ability in winter wheat. crop Sci., 8: 322-324.
- Hamada, A.A. (1988): Studies on some selection criteria in wheat. M.Sc. Thesis, Fac. of Agric. Zagazig Univ.
- Hamdy, M.N.A. (1978): Genetic studies on gene action and interaction in hybrid wheat (*Triticum aestivum* L.). M.Sc. Thesis Fac. Agric. Shebin El- Kom Tanta Uni., Egypt.
- Harrington, J.B. (1940): Yield Capacity of wheat crosses as indicated by bulk hybrid testes. Canad. J Res. 18: 578-484.
- Hassaballa, E.A.; M.A. Khalifa; M.A. El - Morshidy and A.A. Ismail (1984).

- Inheritance of some agronomic characteristics in wheat (*Triticum aestivum* L.) . Assiut J. of Agric. Scie., 15 (8): 235 - 247.
- Hassan, E.E and A.M. Abd El - Moniem (1991) . combining ability, heterosis and gene action of some quantitative characters in wheat. Zagazig, J. Agric. Res., Vol . 18 (5) 1369: 1381.
- Hayman, B.I. (1954a) .The analysis of variance of diallel tables *Biometrics*, 10 : 235 - 244 .
-(1954b) : The Theory and analysis of diallel crosses .
Genetic, 39 : 789 - 809 .
-and K . Mather (1955) : The description of gene interaction in continuous variation.*Biometrics*, 10: 69 - 82.
- Hendawy, F.A (1989): Influence of nitrogen levels on heterosis and combining ability in common wheat . *Menofiya J. Agric Res.* Vol. 14 No.1:135-152
- Hendawy, F.A (1994): General and specific combining ability estimates in a diallel cross of seven bread wheat varieties. *Menofiya J. Agric. Res.*, Vol Vol. 19 No. 1: 75- 93.
-H. I. (1990) Breeding for yield and its components in wheat.M.Sc. Thesis Fac. Agric. Menofiya Univ. Egypt .
- Ikram Uti Hag and Tanach laila (1991). Diallel analysis of grain yield and other agronomic traits in durum wheat. *Rachis*, Vol. 10, No. 1.8-13.
- Johnson; K.J. Biever; A. Hounold and J.W. Schmidt (1966): Inheritance of plant height, yield of grain and other plant and seed characteristics in a cross of hard red winter wheat,(*Triticum aestivum* L) *Crop Sci.*, 6: 336-338.

-, V.A. and J.W. Schmidt (1968): hybrid wheat. *Advances of Agron.*, 20:199-232.
- Jinks, J.L. (1954): The analysis of continuous variation in a diallel cross of *nicotiana rustica* varieties *Genetics*, 39: 767 - 788.
- (1955): A survey of the genetical basis of heterosis in a variety of diall cross. *Heredity* 9 : 233 - 238.
- Jinks, J.L. and B. L. Hayman (1953). The analysis of diallel crosses. *Maize Genetic coop. News*, 27: 48- 54.
- Kaltsikes, P.J and J. Inheritance of plant height, yield of grain and other plant and seed Lee (1971): Quantitative inheritance in durum wheat. *Can. J. Genet. Cytol.* 13: 210- 218.
- Karrar, A.M.A. (1980): Performance and combining ability of durum wheat in a diallel cross. M.Sc. Thesis Fac. Agric. El - Minia Univ. Egypt.
- Knott, D.R. (1965): Heterosis in seven wheat hybrids. *Can.J. Plant Sci.*, 45:499- 501.
- Kronstad, W.E. and W.H. Foote (1964): General and specific combining ability estimates in winter wheat (*Triticum aestivum* L.). *Crop Sci.*, 4: 616 - 619.
- Kumar, H., G.S. Sharma and R.B. Singh (1983): Genotype x environment interaction in relation to combining ability in spring wheat. *Indian J. Genet Pl. Breed.* 43 : 232 - 238.
- Larrea, P.E. (1966): Agronomic milling and backing characters of the hybrid

- spring wheat (*T. aestivum* L.).M.Sc. Thesis, North Dakota state univ., Fargo.
- Lee, Y.W. (1974): Heterosis, inbreeding depression and combining ability estimates for diallel crosses in hard red winter wheat. Dissertation Abstracts International, 34(10): 4788 B. (C.F. Pl. Breed. Abst.46:7882).
-, and E.L. Smith (1969): Heterosis and combining ability in crosses among seven hard red winter wheat varieties. Agron, Abst., Madison Pp.12 (C.F. Pl. Breed. Abst. 41: 4833).
- Lonc, W. (1987) . Gene action for agronomic characters in winter wheat. Referativnyi zhurnal 1: 65 -238
(C.F. Wheat, Barley and Triticum, Abst. April.1988, 918)
-and D. Zalewski (1991). Diallel analysis of quantitative characters in F1 hybrids of winter wheat. Hodowla Roslin, Aklimatyzacja Nasiennictwo 35 (3-4) 101-113(C.F. Plant Breed. Abst. Vol.62No.6,4954,1992.).
- Mahdy, E.E. (1988): The relative importance of the types of gene action in the some agronomic traits in (*T. aestivum* L.) Assiut J. Agric. Sci., 19(13) : 215-224.
- Maloo, S.R. (1987): Combining ability for grain yield and its contributory characters in durum wheat. Indian J.Agric. Sci.57(8):535-538.
- Mani, S.C.and M.V.Rao (1975): Heterosis in winter x spring wheat crosses. Indian J. Genet. Pl. Breed. 35: 330 - 335.

- Mather, K. and J.L. Jinks, (1971). Biometrical genetics (2nd edition).
Chapman and Hall, Ltd., london.
- Maya De Leon, J.L. (1975): combining ability and association of agronomic traits involving three sources of dwarfism in wheat, (*T. aestivum* L.)
Dissertation Abstracts International, 35(7): 3138 B. (C.F. Pl. Breed Abst. 47: 2137).
- Mc- Neal, F.H; D.F. Bladridge; M.A. Berg and C.A. Walton (1965): Evaluation of three hard red spring wheat crosses for heterosis Crop Sci.,5:399- 400
- Mekhamer, K.I. (1995) Breeding for some quantitative traits in common wheat.
M.Sc. Thesis Fac. Agric. Menofiya Univ. Egypt.
- Mitkess, R.A . (1981): Inheritance of some morphological and technological characters in wheat crosses. Ph. D. Thesis. Cairo, Univ. Egypt .
- El - Rassas (1986). Gene action in the inheritance of some characters of bread wheat. Egypt J. Genet Cytol. ,15: 135 - 146, J.,1986.
- Mosaad, M.G. (1981). Genetical and technological studies on durum wheat (*T. durum*). Ph. D. Thesis Fac. Minia Univ. Egypt
-, M. A. El - Morshidy, B.R. Bakheit, and A.M. Tamam (1990)
Genetical studies of some morpho-physiological traits in durum wheat crosses. (Assiut J. Agric. Res., Vol. 21-1,78: 94.)
- Narula, P.N. (1987): A diallel cross analysis of grain number and grain weight under different environments in wheat. Wheat information service, No. 64:37- 43(C.F.Pl.Breed. Abst.58: 2940) .

Palmer, T.P.(1952): Population and selection studies in a triticum cross.

Heredity, 6: 171-185.

Patwary, A.K.; M.U. Ghani and M.M. Rahuman (1986): Heterosis in wheat.

Indian J Agric. Sci., 56 : 382 - 384 .

Peterson, C.J.; W.E. Krnostad, and O.A. Voged (1969): Maximizing heterosis in winter wheat. Agron Abst. Madison, Pp. 15 (Abst) (C.F. Pl. Breed. Abst. 41: 4837).

Ranvir - Singh, G.S. Bhullar and K.S. Gill (1983): Combining ability over environment in durum wheat. Indian J. Genet. Pl.Breed., 43: 152-155.

Rodriguez, R., m.a. Quinones, N.E. Borlaug, and I. Narvaez (1967): Hybrid wheat. Their development and food potential. Central Interaction DeMejoramiento Demaizy Trigo Res, Bull. 3: 1-37.

Sharma, S.K.and R.K. Singh (1983): Combining ability for harvest index and its components in wheat. Indian J. Genet. Pl. Breed. 43: 229 - 231 .

....., and, I. Singh and K. P. Singh (1986): Heterosis and combining ability in wheat (T. aestivum L.) Crop Improvement, 13: 101 - 103.

Sajnani, D.N. (1968): Studies on hybrid vigor and combining ability in wheat diallel crosses. Dissertation Abstracts International, 28(68):4641B. (C.F Pl. Breed. Abst. 39: 3335).

Sikka, S.M.K.; K.B.L. Jain and K.S. Parmer (1959): Evaluation of the potentialities of wheat crosses based on mean parental and early

generation values. Indian J. Genet. Pl. Breed . 19: 150 - 170.

Singh, ss. (1990). Bias caused by episasis in the estimates of additive and dominance components and their interactions with environment in wheat
Indian J. Genetics and Plant Breed . 50 (2) 157 - 160.

....., R.K, and B.D. Chaudhary (1977). Biometrical Methods in quantitative Genetic Analysis. Kalyani Publishers, Ludhiana, New Delhi, India.

....., K.B. and V.P. Gupta (1969): Combining ability in wheat . Indian J. Genet. Pl. Breed. 29: 227 - 232.

.....; D. Sharma and P.D. Mehndiratta (1969): Study of combining and genetic parameters for yield and its components in wheat. Idengku Zasshi/ Jap. J. Genet. 44: 367 - 377. (C.F. Breed. Abst. 41: 770).

.....; P.V.; M.B. Singh; Z. Ahmed, and A.N. Khana (1973): Combining ability study on some developmental traits related to yield productivity in breed wheat. Indian J. of Fram Science (1) : 21 - 28. (C.F. Breed. Abst. 45: 1068).

....., L.; R.S. Paroda and P.K. Behl (1986): Diallel analysis for combining ability over environments in wheat. Wheat Information Service, 61/62 : 74-76.

....., R.P.; R.B Singh; Subedar Singh and G. Misra (1984): Heterosis and inbreeding depression in relation to wheat improvement. American J. Botany, 71: 102.

Singh, R.R.; R.B. Singh; Subedar Singh and G. Misra (1984): Heterosis and inbreeding depression in relation to wheat improvement. American J. Botany, 71:102

Soomro, B.A. and R .Aksel (1976): General and specific combining ability analysis of some quantitative characters in five parent diallel crosses of

- common wheat. Egypt. J. Genet. Cytol. 5: 331 - 345.
- Stuber, C.W.; V.A. Johnson and J.W. Schmidt (1962): Grain protein content and its relationship to other plant and seed characters in the parents and progeny of a cross of (*T. aestivum* L .). Crop Sci., 2: 506 - 508.
- Tamam, A. M. (1989): Inheritance of yield and Aphid resistance in some wheat crosses. M.Sc. Thesis, Assiut. Univ. Egypt.
- Uppal, S.; V.P. Singh and V. Chawla (1987): studies on heterosis, inbreeding d depression and gene effects in bread wheat. Haryana Agric. Univ., Hissar India, 17(2): 114 - 119. (C.F. Breed. Abst. 58: 9253).
- Younis, S.E. A.; M.K. Omara and M.Y. Hussein (1988): A genetic analysis of earliness in wheat and the response to selection for flowering time under favourable clay soil moisture stress conditions in sandy soils. Assiut. J . Agric. Sci., 19 (5): 35 - 48.