

REFERENCES

- Abd El-Aleem, M.M. and S.R.S. Sabry (1994). Wheat population density and distribution under rainfed conditions. Proc. 6th - Conf. Agron., Al-Azhar Univ., Cairo, Egypt, 1: 351-364.
- Abd El-Gawad, A.A.; A.E. El-Tabbakh; A.S. Edris and A.M. Abo-Shetaia (1986). Potential productivity of wheat in Egypt. VIII- Effect of seeding rates on yield and its components. Annual of Agric., Ain Shams Univ. 31: 1173-1182.
- Abd El-Latif, L. and M.K. El-Tuhamay (1986). Effect of nitrogen fertilization levels and seeding rates on growth and yield of wheat. Annuals of Agric. Sci. Fac. Agric., Ain Shams Univ., 3: 265-272.
- Alam, M.T.; M.A. Ghaffer and M.A. Kashem (1994). Critical period of weed competition in wheat (*Triticum aestivum* L.) as influenced by different seed rates. Bangladesh J. of Scientific and Industrial Res. 29: 63-70.
- Al-Marsafy, H.T.; E.E. Hassanein and S.K. Mahmoud (1992). The potential chemical control of wild oats and other weeds in wheat. J. Agric. Sci. Mansoura Univ. 17(4): 705-713.
- Anderson, R.L. (1993). Jointed goatgrass (*Aegilops cylindrica*) ecology and interference in winter wheat. Weed Sci. 41: 388-391.
- Anonymous, (1991). European Weed Res. Society. No. 50.
- Anten, N.P.R. and M.J.A. Werger (1996). Canopy structure and nitrogen distribution in dominant and sub-ordinate plants in a dense stand of *Amaranthus dublius* L. with a size hieracrchy of individuals, Oecoogia, 105: 30-37. [C.F. Weed Abst. 1995, 45: 2131).

- Appleby, A.P. and B.D. Brewster (1992). Seeding arrangement on winter wheat (*Triticum aestivum*) grain yield and interaction with Italian Ryegrass (*Lolium multiflorum*). Weed Technol. 6: 820-823.
- Assey, A.A. and A.A. Ibrahim (1976). Response of local and introduced Mexican wheats to varying seeding rates. J. Agric. Sci. Zagazig Univ., 3: 143-160.
- Assey, A.A.; I.E. Ramadan; E.M. El-Nagar and I.M. Shrief (1983a). Effect of seed rates and weed control methods on growth and grain development of wheat. Zagazig J. Agric. Res. 10: 187-198.
- Assey, A.A.; I.E. Ramadan and E.M. El-Nagar (1983b). Effect of seed rate and weed control methods on weeds and yield of wheat. Zagazig J. Agric. Res. 10: 199-217.
- Balyan, R.S.; R.K. Malik; R.S. Dankar and S.M. Sinkh (1994). Effect of time and method of isoproturon application on weeds in wheat. Haryana Agric. Univ. J. of Res., 24: 158-162. [C.F. Weed Abst., 1996, 45: 261].
- Barhoma, M.A.; M.F.H. Atwa and Sania, Hassan (1990). Effect of Arelon and Brominal on chemical and technological properties of wheat grains. Minufiya J. Agric. Res. 15(2): 2177-2190.
- Barton, D.L.; D.C. Thill and B. Shafii (1992). Integrated wild oat (*Avena fatua*) management affects spring barley (*Hordeum vulgare*) yield and economics. Weed Technol., 6: 129-135.
- Black, C.D.; D.D. Evans; L.E. Ensminger; J.L. White and F.E. Clark (1965). Methods of soil analysis. Part I and Part II. American Soc. of Agron. Inc. Publisher, Madison-Wisconsin, U.S.A.

- Blackshaw, R.E. (1993a).** Downy brome (*Bromus tectorum*) density and relative time of emergence affects interference in winter wheat (*Triticum aestivum*) Weed Sci. 41: 551-556.
- Blackshaw, R.E. (1993b).** Downy brome (*Bromus tectorum*) interference in winter rye (*Secale cereal*). Weed Sci. 41: 557-562.
- Blackshaw, R.E. (1994).** Differential competitiveness ability of winter wheat cultivars against downy brome. Agron. J. 66: 649-654.
- Blue, E.N.; S.C. Mason and D.H. Sander (1990).** Influence of planting date, seeding rate and phosphorus rate on wheat yield. Agron. J. 82: 762-768.
- Buchanan, G.A. and R.D. McLaglin (1975).** Influence of nitrogen on weed competition in cotton. Weed Sci., 20; 324-327.
- Bussan, A.J.; O.C. Burnside; J.H. Orf; E.A. Ristau and K.J. Puettmann (1977).** Field evaluation of soybean (*Glycine max*) genotypes for weed competitiveness. Weed Sci. 45: 31-37.
- Challaiah, R.E.; O.C. Burnside; G.A. Wicks and V.A. Johnson (1986).** Competition between winter wheat (*Triticum aestivum*) cultivars and downy brome (*Bromus tectorum*). Weed Sci. 34: 689-693.
- Christensen, S. (1990).** Varietal differences in spring barley with respect to competitive ability. European Weed Res. Soc. Newsletter No. 46: 15-16.
- Christensen, S. (1995).** Weed suppression ability of spring barley varieties. Weed Res. 35: 241-247.
- Christensen, S.; G. Rasmussen and J.E. Olesen (1994).** Differential weed suppression and weed control in winter wheat. Aspect of Applied Biology, 40: 335-342. [C.F. Weed Abst., 1996, 45: 269].
- Christensen, S.; J.C. Streibig and H. Haas (1990).** Interaction between herbicide activity and weed suppression by spring barley

- varieties. Symposium on Integrated Weed Management in Cereals. Proceeding of an EWRS Symposium, Helsinki, Finland, 4-6 June. Wageningen, Netherlands; European Weed Research Society, 367-374.
- Cosser, N.D.; M.J. Gooding and R.J. Froud-Williams (1995). The effect of wheat dwarfing genes on competition against blackgrass. British Crop Protection Council., 1: 361-362. [C.F. Weed Abst., 1996, 45:1716].
- Cudney, D.W.; L.S. Jordan and A.E. Halla (1991). Effect of wild oat (*Avena fatua*) infestation on light interception and growth rate of wheat (*Triticum aestivum*). Weed Sci. 39: 175-179.
- Cudney, D.W.; L.S. Jordan; J.S. Holt and J.S. Reints (1989). competitive interactions of wheat (*Triticum aestivum*) and wild oats (*Avina fatua*) grown at different densities. Weed Sci., 37: 538-543.
- Dhawan, R.S. (1995). Effects of isoproturon on Hill activity and membrane permeability in leaves of wheat (*Triticum aestivum*), small canary grass (*Phalaris minor*) and wild oat (*Avena ludoviciana*). Indian J. of Agric. Sci., 65: 894-895.
- Doll, H.; U. Holm and B. Sogaard (1995). Effect of crop density on competition by wheat and barley with *Agrostemma gethago* and other weeds. Weed Res. 35: 391-396.
- Donald, W.W. and K. Mohammad (1996). Canada thistle (*Cirsium arvense*) effects on yield components of spring wheat (*Triticum aestivum*). Weed Sci. 4: 114-121.
- Dover, P.a. and J. East (1990). The effect of variety blends and seed rates on disease and weed incidence in wheat grown in organic systems. Monograph-British Crop Protection Council, 45: 26).

- Dunan, C. and R.L. Zimdahl (1991). Competitive ability of wild oats (*Avena fatua*) and barley (*Hordeum vulgare*). Weed Sci. 39: 558-563.
- Easson, D.; A.D. Courteney and J. Picton (1995). The effects of performance of cereal crops. In integrated crop protection towards sustainability. Proceedings of a symposium, Edinburgh, UK, 11-14 September, British Crop Protection Council, 305-312. [C.f. Weed Abst. (1996), 45: 687].
- Eissa, A.M.K. (1979). Response of some wheat varieties to cultural treatments. M.Sc. Thesis, Cairo Univ. Egypt.
- El-Bana, A.Y.A. and H.A. Basha (1994). Response of yield and yield attributes of wheat (*Triticum aestivum* L.) to planting density under newly cultivated sandy soil. Zagazig J. Agric. Res. 21(3A): 671-681.
- El-Deek, M.H. and A.M. Abo El-Kheer (1986a). The influence of substituted urea herbicides on wheat growth and yield as well as its common associated weeds. J. Agric. Sci. Mansoura Univ. 11: 1384-1393.
- El-Deek, M.H. and A.M. Abo El-Kheer (1986b). Effect of some foliar herbicides on weed control and yield of wheat. J. Agric. Sci. Mansoura Univ. 11: 1394-1402.
- El-Deepah, H.R.A. (1989). Evaluation of some herbicides. Annals of Agric. Sci. Moshtohor, 27(3): 1381-1393.
- El-Ghareib, E.A. and M.M. El-Monoufi (1988). Effect of seeding rates and nitrogen fertilization levels on the productivity of Giza 157 Egyptian wheat cultivar. Annuals Agric. Sci. Fac. of Agric. Ain Shams Univ., 33: 1031-1044.

- El-Metwaly, I.M. (1994).** Response of some wheat varieties to some herbicides and nitrogen fertilization. M.Sc. Thesis, Fac. of Agric. Mansoura Univ.
- El-Moursy, S.A. (1989).** Effect of weed control treatments and cycocel concentration on growth, yield and yield components of wheat. J. Agric. Sci., Mansoura Univ. 14: 23-29.
- El-Shaer, M.H.; M.S. El-Sayed; H.M Soliman and M.A. Rady (1988).** Wheat yield and yield components as related to sowing dates and seeding rates as microclimatic modifiers. Annals of Agric. Sci., Moshtohor, 26: 829-842.
- Essa, F.A. (1990).** Effect of plating methods, seeding rate and time of nitrogen application on yield and yield components of some wheat varieties (*Triticum aestivum vulgare* L.) M.Sc. Thesis, Fac. Agric. Mansoura Univ.
- Felicio, J.C. (1984).** Sowing density in three wheat cultivars. Pesquise Agropecuaria Brasileira, 19(4): 455-460. (C.F. Field Crop Abst., 1985, 38(5): 2777).
- Ferrero, A.; A.C. Sparacino; S. Bocchi and N. Riva (1994).** Influence of shading on *Heteranthera reniformis* Ruitzt pavon growth. 46th International Symposium on crop protection: part IV< Gent, Belgium, 3 May. Mededelingen-Faculteit Landbouwkundige en Teogepaste Biologische Wetenschappen, Universiteit Gent, 59(3b): 1333-1337. [C.F. Weed Abst. 1996, 45: 2132].
- Frederick, J.R. and H.G. Marshall (1985).** Grain yield and yield components practices. Agron. J., 77: 495-499.
- Gabr, E.M. (1983).** Effect of some weed control on the growth and yield of what. M.Sc. Thesis, Fac. of Agric. Mansoura Univ.

- Garrity, D.P.; M. Movillon and K. Moody (1992). Differential weed suppression ability in upland rice cultivars. *Agron. J.* 84: 586-591.
- Ghafoor, A. and R.A. Shad (1995). Relative competition of grasses, broad leaved and mixed weed population with wheat. *Emirates J. of Agric. Sci.*, 7: 55-64.
- Ghanem, S.A.I. and A.A.H. El-Khawaga (1991). Growth, yield and its attributes of wheat as influenced by seeding rates and chemical weed control. *Zagazig J. Agric. Res.*, 18: 1403-1416.
- Gill, D.S. and K.S. Sandhu (1994). Response of wheat and sunflower to allelopathic effects of weed residues. *Indian J. of Ecol.*, 21: 75-78.
- Gil, H.S.; N.S. Walia and L.S. Brar (1988). Influence of method of application on the performance of isoproturon in wheat. *Haryana J. of Agron.* 4: 112-115. [C.F. Weed Abst. (1991) 40: 1427].
- Gogol, A.K. and H. Kalita (1995). Effect of seeding methods and herbicide on weeds and growth and yield of wheat (*Triticum aestivum*). *Indian J. of Agron.* 40: 209-211.
- Gomaa, A.A.; S.M. Desouki; M.M. Sadek, E.H. Talaat; H.S. Khalil; S.A. Attia; K.F. Hegazi; E.H. Ghanem; M.M. Hamdi and M.A. Gouda (1979). Giza 157 and Sakha 8 two new bread wheat varieties. First Con. Agric. Res. Center, Giza, Egypt.
- Gomaa, A.A.; O.H.S. Khalil; R.A. Abu-Elenein and S.E. El-Kalla (1977). Effect of plant density and nitrogen level on grain yield and its components for two wheat varieties. *Agric. Res. Rev.*, Cairo, 55: 1-25.
- Gomez, K.A. and A.A. Gomez (1983). Statistical procedures for Agricultural Res. A Wiley Interscience Publication, John Wiley & Son (2nd ed.)

- Gonzalez, P.; R.A. Lamela and M.L. Salas (1990).** Control wild oats by triallate and its effects on growth and grain yield of wheat varieties. *European Weed Res. Soc.* 399-404.
- Gouda, M.A.; M.M. El-Shami and M.S. Sharshar (1994).** Effect of planting methods, seeding rates and use of herbicide on yield and its components of wheat. *J. Agric. Sci. Mansoura Univ.* 19: 39-47.
- Hagras, A.M. (1985).** Influence of seed rates and nitrogen fertilization on yield of durum wheat. *Annual Agric. Sci., Fac. of Agric., Ain Shams Univ.* 30: 929-949.
- Harrison, S.K. and J. E. Beuerlein (1989).** Effect of herbicide mixtures and seeding rate on red winter wheat (*Triticum aestivum*) yield. *Weed Technol.* 3: 505-508.
- Hegazi, K.F.; E.H. Ghanem; A.M.A.S. Ali and E.S. Shokr (1982).** The effect of seeding rate and nitrogen application on the yield and its components of the wheat variety Giza 157. *Annals of Agric. Sci. Moshtohor* 18: 1-13.
- Hooda, I.S. and S.K. Agarwai (1995).** Response of weeds to levels of irrigation, weed control and fertility in wheat. *Brighten Crop Protection Conference: Weeds. Proceedings of an Integration Conference, Brighton, UK* < 20-23 Nov. Farnham, UK; British Crop Protection Council, 2: 679-682. (C.F. Weed Abst., 1996, 45: 1951).
- Hussein, M.A.; A.H. El-Hattab and A.K. Ahmed (1974).** Competition between wheat plants and weeds as affected by nitrogen levels and wheat seeding rates. *Egypt J. Agron.*, 3: 35-48.
- Hussein, M.A.; A. Kandil; El-Sayed Shokr and M.M. Abd El-Aleem (1981).** Effect of seeding methods and nitrogen fertilizers on some agronomic characters of Giza 157 and Sakha 8 wheat

- cultivars. *Triticum aestivum* L. Annals of Agric. Sci., Moshthor, 15: 1-12.
- Johnson, J.W.; W.L. Hargrove and R.B. Moss (1988).** Optimizing row spacing and seeding rate for soft red winter wheat. Agron J. 80: 164-166.
- Joseph, K.D.; M.M. Alley; D.E. Brann and W.D. Grevelle (1985).** Row spacing and seeding rate effects on yield and yield components of soft red winter wheat. Agron. J. 77: 211-214.
- Justice, G.G.; T.F. Peeper; J.B. Solie and F.M. Epplin (1993).** Net returns from cheat (*Bromus secalinus*) control in winter wheat (*Triticum aestivum*). Weed Technol. 7: 459-464.
- Kalita, P. and A.K. Choudhury (1984).** Effects of varieties seed sizes and seed rates of the yield of wheat. Indian J. of Agron. 29: 287-290.
- Khalil, M.K.M. (1989).** Effect of nitrogenous fertilization and chemical weed control on growth, yield and yield components and technological properties of wheat (*Triticum aestivum*, L.) M.Sc. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ.
- Khan, M.; W.W. Donald and T. Prato (1996).** Spring wheat (*Triticum aestivum*) management can substitute for diclofop for foxtail (*Setaria sp.*) control. Weed Sci. 44: 362-372.
- Khera, K.L.; B.S. Sandhu; T.S. Aujla; C.B. Singh and Kumar (1995).** Performance of wheat (*Triticum aestivum*) in relation to small canarygrass (*Phalaris minor*) under different levels of irrigation, nitrogen and weed population. Indian J. of Agricultural Sciences, 65: 717-722.
- Khokhar, M.A.; M.S. Sheikh; M. Siddique and M.S. nazar (1985).** Effect of different seeding densities and nitrogen levels on the

- yield of two wheat genotypes. Pakistan J. of Agric. Res. 6(3): 150-152. (C.F. Wheat and Barley Abst. (1987), 4: 3179).
- Kirkland, K.J. (1993).** Spring wheat (*Triticum aestivum*) growth and yield as influenced by duration of wild oat (*Avena fatua*) competition. Weed Tech o. 4: 890-893.
- Kolp, B.J.; R.G. Sackett; K.E. Bohnenblust and G.P. Roehrkaske (1973).** Effect of rate and data of seeding Shonoshi winter wheat on soil moisture depletion. Agron. J. 65: 929-930. In Tanji *et al.* (1997). Competitive of wheat. Weed Sci. 45: 481-487.
- Koraieem, Y.S.; A.A. Bary and A.A. Khater (1979).** Effects of cultivars and seeding rates on wheat production under mechanized and conventional farming practices. Alex. J. of Agric. Res., 27(2): 323-329.
- Koscelny, J.A.; T.F. Peeper; J.B. Solie and S.G. Solomon (1990).** Effect of wheat (*Triticum aestivum*) row spacing, seeding rate and cultivar on yield loss from cheat (*Bromus secalinus*). Weed Technol., 4: 487-492.
- Kumar, A.; A.K. Sinha; B.P. Sinha; K.M.P. Singh and S.S. Thakur (1991).** Response of wheat genotypes to seed rates and row spacing. Indian J. Agron. 36: 78-82.
- Laila Aboushoba; M. El-Afry and M.A. Gouda (1986).** Effect of two herbicides, Brominal and Tribunil on yield and photosynthetic pigments contents in wheat. J. Agric. Sci. Mansoura Univ., 11(3): 987-992.
- Lee, H.C. (1995).** Non-chemical weed controlled in cereals. Brighton Crop Protection Conference: Weeds, Proceedings of an International Conference, Brighton, UK, 20-23 Nov., Farnham, UK; British Crop protection Council, 3: 1161-1170, [C.F. Weed Abst., 1996, 45: 1940].

- Mahmoud, S.K. (1988).** Study on some agricultural factors affecting growth, yield, yield components and technological characteristics of wheat. Ph.D. Thesis, Fac. of Agric., Minia Univ., Egypt.
- Makwaski, R.M.D. (1995).** Round-leaved mallow (*Malva pusilla*) interference in spring wheat (*Triticum aestivum*) and lintil (*Lense culinaris*) in Saskatchewan. Weed Sci. 43: 381-388.
- Malik, R.K. and S. Singh (1995).** Littleseed canarygrass (*Phalaris minor*) resistance to isoproturon in India. Weed Technol. 9: 419-425.
- Malitius, O.; F. Bergmann; A. Sidler; E. Meister and P. Weisskopf (1996).** Integrated versus intensive crop management. 3. Cereal-dominated rotation. Revue-Suisse-d'agriculture (Switzerland) (Mar.-Apr.), 26: 99-103.
- Marwat, A.Q.; M. Karim; S.K. Khalil; A.L. Wazir and J. Bakhat (1989).** Effect of land preparation methods and seeding rates on weed population and wheat yield. Sarhad J. of Agric. 5: 107-111.
- Mason, M.G. and R.W. Madin (1996).** Effect of weeds and nitrogen fertilizer on yield and grain protein concentration of wheat. Australian J. of Experimental Agric., 36: 442-450.
- Mazurek, J. (1984).** Effect of sowing density on yield components and grain quality in new spring wheat lines and varieties. Biultyn Instytutu Hodowlil Akilimatyzacji Roshlim No. 155: 73-81. (C.F. Field crop Abst., 1985, 38(12): 6859).
- Menshawy, A.M.M. (1996).** Genetic studies on earliness in wheat. M.Sc. Thesis, Fac. of Agric., Zagazig Univ.
- Mosalem, M.E. (1993).** Response of two wheat cultivars to nitrogen level and seeding rate. J. Agric. Res. Tanta Univ., 19: 791-805.

- Mourad, M.A.; E.E. Shalaby and AA. Galelah (1992).** Effect of micronutrients and herbicide application on wheat growth and yield. J. Agric. Res. Tanta Univ., 18: 643-661.
- Muendel, H.H.; R.J. Morrison; R.E. Blackshaw and B. Roth, eds. (1992).** Safflower production on the Canadian prairies. Agric. Canadian Res. Stn. Lethbridge, Alberta. 36 pp. in Blackshaw, 1993. Safflower effects on competition with green foxtail. Weed Sci. 41: 403-408.
- Niemann, P. (1990).** Weed suppression by cereal cultivars. European Weed Res. Soc. Newsl. 46: 15-16.
- Osman, A.M. and Z.M Mahmoud (1981).** Yield and yield components of wheat (*Triticum aestivum* L.) and their inter relationships as influenced by nitrogen and seed rate in the Sudan. J. of Agric. Sci. UK, 97: 611-618.
- Pantone, D.J.; W.A. Williams and A.R. Maggenti (1989).** An alternative approach for evaluating the efficacy of potential biocontrol agents of weeds, 2-path analysis. Weed Sci. Soc. 43: 126 pp.
- Panwar, R.S.; R.K. Malik; R.S. Balyan; S.S. Punia and S.S Rathi (1995a).** Performance of clopyralid, Isoproturon and 2, 4-D on control of Canada thistle (*Cirsium arvense*) in wheat (*Triticum aestivum*). Indian J. of Agron., 40: 102-104.
- Panwar, R.S.; R.K. Malik; R.S. Balyan and D.P. Singh (1995c).** Effect of isoproturon, sowing method and seed rate on weeds and yield of wheat (*Triticum aestivum*) Ind. J. of Agricultural Sciences, 65: 109-111.
- Panwar, R.S.; R.K. Malik; S. Singh and R.S. Balyan (1994).** Influence of tralkoxydim applied alone or as tank mixture on the control of

- grassy weeds in wheat. Haryana Agric. Univ. J. of Res. 24: 25-32.
- Panwar, R.S.; S.S. Rathi and R.K. Malik (1995b).** Effect of isoproturon and 2,4-D combination on weed control in wheat. Haryana Agric. Univ. J. of Res. 25: 101-105 [C.F. Weed, Abst. 1996, 45: 2782].
- Peterson, D.E and J.D. Nalewaga (1992).** Green foxtail (*Setaria viridis*) competition with spraying wheat (*Triticum aestivum*). Weed Techno. 6: 291-296.
- Ponce, R.C. and M.L. Salas (1990).** Competition of wild oat with wheat in comparison to wheat with itself. European Weed Res. Soc. Newsl. No. 46: 16-17.
- Pradhan, L.; D. Rout and G.K. Patro (1989).** Rationalization of plant population, fertilizer and method of weed control in irrigated wheat on farmer's fields in Sambalpur district (Orissa). Orissa J. of Agricultural Res. 2: 143-147. [C.F. Weed Abst. (1991) 40(5): 1426].
- Prasad, K. (1985).** Efficiency of post-emergence weedcides applied at different stages in control of annual weeds of irrigated wheat under mid-Himalayan conditions. Pesticides, 19(6): 20-22.
- Prasad, K. and R.S. Singh (1995).** Influence of weed and nitrogen management on weed growth, nutrient uptake and yield of wheat (*Triticum aestivum*). Indian J. of Agricultural Sciences, 65: 117-122.
- Qasem, J.R. (1995).** Allelopathic effect of some arable land weeds on wheat (*Triticum aestivum*): a survey. Dirasat. Series B, Pure and Applied Sciences, 22(4): 81-87. [C.F. Weed Abst. 1996 45: 4639].

- Radosevich, S.R. and J.S. Holt (1984).** Weed Ecology: Implications for vegetation management. New York: J. Wiley, pp. 97-99.
- Raghuwanshi, R.K.S.; R.S. Thakur; R. Umat and M.L. Nema (1990).** Crop technology for optimum grain production in sorghum-wheat sequence under resource restraints. Indian J. of Agron. 35: 246-250.
- Rajender, K. and J.S. Hooda (1996).** Effect of methods of weed control on yield of wheat and associated weeds. Annual of Biology (Ludhiana), 12: 49-51.
- Rajput, F.K.M.; A.S. Arain; M.J. Rajput; S.M Alam and A.W. Baloch (1989).** The growth and yield of wheat as affected by different seed rates and row spacing. Sarhad J. of Agric. 5: 479-482.
- Rastogi, S.K.; D.K. Gupta and S.K. Katyal (1984).** Economic evaluation of herbicides in control of (*Phalaris minor*, Retz) in sandy loam soil of Haryan. Pesticides, 18(9): 38-39. [C.F. Field Crop Abst. 1985: 3333].
- Romex, B. and S. Dzienia (1994).** The effectiveness of herbicides applied in winter tritical. Efektywnosc herbicidow stosowanych w Bcenzycie Ozimym. Zeszdy Naukowe Akedemii Romlnicej w Szczecinie. Rolnictwo, 58: 207-212. [C.F. Weed Abst. ,1996, 45: 3545].
- Roshdy, A.(1988).** Effect of seeding rates and some weed control methods on yield, yield components and associated weeds f whet. Annals of Agric. Sci., Moshtohor, 26: 1399-1409.
- Sabry, S.R.S.; A.H. Abdel Lateef and M.H. Iskandar (1994).** The effect of row spacing and seeding rate on wheat (*Triticum aestivum* L.) grain yield under different environments of Egypt. Zagazig J Agric. Res. 21: 1015-1021.

- Saif Al-Aghbare, A.O.A. (1988).** Effect of some agricultural treatments on wheat under sprinkler irrigation system. M.Sc. Thesis, Fac. of Agric., Moshtohor, Zagazig Univ. Egypt.
- Salem, A.H.; H.A. Rabie; M.A. Mohamed and M.S. Salim (1990).** The relative importance of yield contributing characters in wheat as influenced by different locations and seeding rates. Proc. 4th Conf. Agron. Cairo, (1): 99-110.
- Salonen, J. (1990).** Interaction between crop density and weed infestation in spring cereals. European Weed Res. Soci. Newsletter, (46): 31-34.
- Samiullah, M.; M.M. Afridi and A. Inam (1990).** Determination of seeding rate for triticale: effect of varying seeding rates on population count, leaf-NPK contents and growth, yield and quality characteristics of four new triticales. Geobios (Jodhpur), 17, (5-6): 255-260. [C.F. Field Crop Abst. (1991) 4: 2804].
- Samuel, A.M. and J. East (1990).** Organically grown wheat-the effect of crop husbandry on grain quality. Aspects of Applied Biology, 25: 199-208. [C.F. Field Crop Abst. (1991) 44: 3514].
- Samuel, A.M. and S.J. Guest (1990).** Effect of seed rates within crop cultivations in organic winter wheat. Monograph British Crop Protection Council No. 45: 49-54.
- Satorre, E.H. and R.W. Snaydon (1992).** A comparison of root and shoot competition between spring cereals and *Avena fatua* L. Weed Res. 32: 45-55.
- Seavers, G.P. and K.J. Wright (1995).** Potential for weed control by suppressive cereal cultivars. British Crop Protection Council 2: 737-742. [C.F. Weed Abst. 1996 45: 1718].

- Shabashon, V.V.; G.G. Duda; A.A. Egorshin and A.V. Baranovski** (1990). Regulation of grain yield and quality of winter wheat. *Agrokimiya*, 1: 42-48. [C.F. Field Crop Abst. (1990) 43: 3685].
- Sharma, M.L.; S.K. Bhardwaji and G.S. Bhardwgi** (1989). Comparative study of chemical weed control in wheat. Haryana J. Agric. 5: 177-178[C.F. Field Crop Abst., 1992, 45: 5919].
- Sharma, P.K. and P.C. Kharwara** (1994). Effect of tillage and weed control on wheat yield in sandy loam soil under irrigated and rainfed condition. *Himachal J. of Agric. Res.* 20: 28-34.
- Sharshar, M.S.** (1989). Effect of some cultural treatments on wheat growth, yield and its components. Ph.D. Thesis, Fac. of Agric., Tanta Univ., Egypt.
- Shehab El-Din, T.M.** (1993). Response of two spring wheat cultivars (*Triticum aestivum* L. em. Thell.) to ten seeding rates in sandy soils. *J. Agric. Sci. Mansoura Univ.* 18: 2235-2240.
- Sidhu, M.S.; J.S. Sawhney and R.S. Navang** (1988). Chemical weed control in wheat-Indian Mustard intercrop System. *Indian J. of Weed Sci.* 20: 92-95.
- Silva, D.B. and Gomes** (1990). Spacing and sowing density in irrigated wheat in the Cerrdo region. *Pesquis Agropecuaria Brasileira*, 25: 305-315. [C.F. Field Crop Abst. (1992) 45: 697].
- Singh, S.; R.K. Malik; R.S. Panwar and R.S. Balyan** (1995). Influence of sowing time on winter wild oat (*Avena ludoviciana*) control in wheat (*Triticum aestivum*) with isoproturon. *Weed Sci.* 43: 370-374.
- Singh, S.; R. Matzen and T.T. Pedersen** (1985). The effect of seed rates and sowing machines on the growth, yield and yield components of spring wheat. *Indian J. Agron.* 30: 55-58.

- Smith, R.J. (1993).** Biological control as component of integrated weed management for rice in the United States. A SPEC Extension Bulletin. pp. 376-399.
- Snedecor, G.W. and W.G. Cochran (1971).** Statistical methods. 6th ed., Iowa State Univ. Press, Ames, Iowa, U.S.A.
- Solie, J.B.; S.G. Solomon,; K.P. Self; T.F. Peeper and J.A. Koscelny (1991).** Reduced row spacing for improved wheat yields on weed-free and weed infested fields. Trans. Am. Soc. Agric. Eng. 34: 1654-1660.
- Soroka, S.V.; L.I. Soroka and A.S. Andreev (1995).** Early spring application of Arelon in winter wheat crops. Zashchita Rastenil (Moskva) 4: 14 [C.F. Weed Abst. 1994, 45: 3176].
- Stahlman, P.W. and S.D. Miller (1990).** Downy brome (*Bromus tectorum*) interference and economic thresholds in winter wheat (*Triticum aestivum*). Weed Sci. 38: 24-228.
- Steinmann, H.H. and B. Gerowitt (1994).** Mechanical control of *Galium aparine* in winter wheat. Maitrise des adventices parvoie non chimique. Communications de la quatrieme conference internationale I.F.O.A.M. Dijon, France, 5-9 July 1993. [Edited by Thomas, J.M.] Quetigny Cedex, France; Association Colloque IFAM Ed. 2, 273-277. [C.F. Weed Abst. (1996), 45: 271].
- Tahoon, R.M. (1994).** Studies on weed control in wheat. M.Sc. Thesis, Fac. of Agric., Al-Azhar Univ., Egypt.
- Tanji, A.; L. Zimdahl and W. Westra (1997).** The competitive ability of wheat (*Triticum aestivum*) compared to rigid ryegrass (*Lolium rigidum*) and cowcockle (*Vaccaria hispanica*). Weed Sci. 45: 48-487.

- Thakur, S.S; I.B. Pandey; S.J. Singh and S.S Mishra (1995).** Response of wheat (*Triticum aestivum*) varieties to weed management under late-sown condition on calcareous soil of north Bihar. Indian J. of Agron., 40: 204-208.
- Thomson , W.T. (1983).** Agricultural chemicals-book II Herbicides. Thomson Publications, Library of Congress Catalog NO. 64-24795, U.S.A.
- Tiwari, J.P.; H.C. Jain and S.K. Dubey (1988).** Efficacy of fluroxypyr control weeds in irrigated wheat. Indian J. of Weed Sci. 20: 35-40.
- Tompkins, D.K.; G.E. Hultgreen; A.T. Weight and D.B. Fowler (1991).** Seed rate and row spacing of no-till wheat. Agron. J. 83: 684-688.
- Valenti, S.A. and G.A. Wicks (1992).** Influence of nitrogen rates and wheat (*Triticum aestivum*) cultivars on weed control. Weed Sci. 40: 115-121.
- Vanova, M.; T. Spitzer and K. Klem (1995).** Control of dicotyledonous weed occurrence in spring barley. Rostinna Vyroba, 41: 83-89. [C.F. Weed Abst. (1996) 45: 116].
- Varge, B. (1980).** Investigation of various wheat varieties reaction upon quantity, time of application and form of N-fertilizers in various agroecological conditions. Polkoprivred na Znanstvena Smotra, 52: 285-297. (C.F. Field Crop Abst., (1985), 38(1): 4).
- Varshney, J.G. and H.G. Singh (1990).** Effect of adjuvants on herbicide efficiency in controlling weeds in wheat. Weeds Sci. 38: 229-236.
- Varshney, J.G. and H.G. Singh (1994).** Nutrient economy through tank mixing of adjuvants with herbicides for wheat. Advances in Agricultural Res. in India, 2: 98-108.

- Verschwele, A. and P. Niemann (1994a).** Managing competition between cultivated plants and weeds by cultivation methods and selective use of herbicides. (3.2. competitive ability of winter wheat cultivars for indirect weed control). Entwicklung eines integrierten Systems der Pflanzproduktion unter Beachtung ökonomischer und ökologischer Aspekte des Pflanzenschutzes im Weizen, 2: 160-174.
- Verschwele, A. and P. Niemann (1994b).** Indirect weed control by selection of wheat cultivars. Maitrise des adventices par voie non chimique. Communications de la quatrième conférence internationale I.F.O.A.M., Digion, France, 5-9 July 1993 [edited by Thomas, J.M.]. Quetigny Cedex, France, Association Colloque IFOAM (1994) Ed. 2, 285-289. [C.F. Weed Abst. 45, 1: 272].
- Vonzounis, N.A. and P.G. Amerikanos (1995).** Post emergence control of weeds in cereals with non-hormone type herbicides. Technical Bulletin-Cyprus Agri. Res. Inst., No. 168, 6 pp.
- Wicks, G.A.; P.T. Nordquist; G.E. Hanson and J.W. Schmidt (1994).** Influence of winter wheat (*Triticum aestivum*) cultivars on weed control in sorghum (*Sorghum bicolor*). Weed Sci. 42(1): 27-34.
- Yadav, P.K.; S.P. Kurchania and J.P. Taiwari (1995).** Herbicide and fertilizer compatibility under normal and stall seedbed sowing of wheat (*Triticum aestivum*) at different levels of nitrogen. Indian J. of Agric. Sci., 65: 265-270.
- Yaduraju, N.T. and K.N. Ahuja (1995).** Response of herbicide resistant *Phalaris minor* to pre and post-emergence herbicides, herbicide mixtures and adjuvants. British Crop Protection Council, 1: 225-230. [C.F. Weed Abst. 1996, 45: 1734].

- Yenish, J.P.; B.R. Durgan, D.W. Miler and D.L. Wyse (1997).** Wheat (*Triticum aestivum*) yield reduction from milkweed (*Asclepias cyriaca*) competition. Weed Sci. 45: 127-131.
- Yongqing, M.; Z. Suying; Z. Guanchang and M. Sumel (1995).** A preliminary investigation of weed growth in stubble of different wheat cultivars. Res. of Agricultural modernization 16(10: 54-56. (C.F. Wed Abst. (1996), 45: 3382).
- Zaher, S.A.M. (1996).** Effect of some agricultural treatments on growth and yield of wheat and associated weeds. Ph.D. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ., Egypt.