

REFERENCES

LITERATURE CITED

- Ahmed, Q.; M.A. Rana, ; and S.U.H. Siddiqui (1991). Sunflower seed yield as influenced by some agronomic and seed characters. *Euphytica*. 56(2) 137- 142.
- Alam, M.S.; M.N. Hossian; A.B. M.A. Khair and M.S. Kham (1987). Genetic parameters and relationship among some agronomic characters in sunflower. *Bangladesh J. of Agric.* 12 (2) 89- 93.
- Alekseev, A.P.; V.P. Voskoboinik, and F.I. Gorbackenko (1980). Combining ability of short forms of sunflower seleksiya i- Semenovodstvo, 1979 No. 4. (1979). U.S.S.R. (C.F. Plant Br. Abstr. So. (1). 523).
- Attia, S.A.M. (1980). Evaluation and correlation studies on sunflower cultivars. M. Sc. Thesis, Fac. of Agric., Kafer El- Sheikh Tanta Univ.
- (1985). Effect of some cultural treatments of sunflower (*Helianthus annuus* L.). Ph. D. Thesis, Fac. of Agric., Al- Azhar Univ.
- Beard, B.H. and S. Geng (1982). Interrelationships of morphological and economic characters of sunflower. *Crop Sci.* 22 (4): 817- 822.
- Burns, R.E. (1970). Head size of sunflower as indicator of plant yield. *Agron. J.*, 62 : 112- 283.
-

- Chaudhary, S.K. and I.J. Anand (1986). Heterosis and inbreeding depression in sunflower. *Genet. Crop Improvement* 84. 11 (1): 15-19. [C.f. Plant Br. Abstr. (1988), Vo. 56. No. 1p. 133).
- Chaudhary, S.K. and I.J. Anand (1993). Relationship of metric traits with seed filling in sunflower (*Helianthus annuus* L.) under open pollination. *International Journal of Tropical Agriculture* 11 (3) 146-148.
- Clement, Y. (1968). Contribution to study of some characteristics of sunflower. *C.R. Acad. Agric. Fr.* 54. 853- 857.
(C.F. Plant Br. Abstr. 39. (3) : 5328).
- Constock, V.E. and J.O. Culbertson (1958). A rapid method of determining the oil content and iodine value. *Agron. J.* 50 : 113-114.
- Cruz, Q.D. (1986). Heterosis and combining ability for yield and yield components in sunflower. *Phillipine J. of Crop Sci.* (1986) 11 (3) 171- 174 [En, 4 Ref.] Res. and Devel. Cent., Central Luzon State Univ., Munoz., Nueva Ecija, Phillipins.
- Dilruba, B.; M.S. Kham and M.A. Khaleque (1988). Genetic and environmental variability in sunflower. *Bangladesh J. of Agric.* 13 (1) : 11- 5 (f ref.).
- Dua, R.P. (1979). Studies on the stability of genetic parameter for the oil and yield components in sunflower (*Helianthus annuus* L.). Thesis Abst. Hissar, India,
(C.F. Plant Br. Abstr. 49 : 318).
-

- Dua, R.P. and T.P. Yadava (1982). Gene effects for some quantitative characters in sunflower. *Indian Journal of Agric. Sci.*, 52 (7) 439-443.
- and ----- (1985). Combining ability in sunflower. *Indian J. of genetics and Plant Breeding* 43 (2) 129- 136.
(C.F. Plant Br. Abstr. 55. 3818).
- D'Yakov, A.B. (1971). Protein content and level of oil accumulation in sunflower seeds. *Vestnik Selekochoze, Nauki* 7 : 57- 63 ves. Inst. Maslick. Kultur. Krasnodor, U.S.S.R.
(C.F. Plant Br. Abstr. 42 : 1070, 1972).
- Earle, F.R.; C.H. Vantten; T.F. Clark and I.A. Wolff (1969). Compositional data on sunflower seed. *J. Amer. Oil Chem. Soci.* 45 : 876- 879.
- El- Ahmmer, B.A. (1989). Oil crops in Egypt. *Oil Crops Proceeding of the three meeting Pantnugar and Hyderabad, India* 4- 17.
- El- Ahmer, B.A.; F.M. El- Rayes; A.O. Omran and A. Ali (1980). Effect of N, P, K and their different combinations and seed, oil content and other economic characters of sunflower (*Helianthus annuus* L.) *Agric. Res. Rev.* 58: 117- 134.
- El- Emam, M.A. (1984). Effect of some cultural treatments on sunflower yield. M. Sc. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ.
- El- Gendy, F.H.; S.A. Attia; W. Kady and S.M. El- Yamany (1992). Response of sunflower to different nitrogen fertilization sources and levels in sandy soils. *Egypt. J. Appl. Sci.*, 7 (12): 717- 727.

- El- Kady, W. (1987). Physiological studies on sunflower. Ph. D. Thesis in Agronomy of Agric. Kafer El- Sheikh, Tanta Univ.
- El- Yamany, S.M.; S.A. Attia and F.H. El- Gendy (1993). Response of sunflower yield to nitrogen application rates under different methods of micronutrients application on sandy soils. *Annals of Agric. Sci., Moshtohor*, Vol. 31 (3), 1381- 1391.
- Enns, H. (1959). A study of the inheritance, and expressivity of some seeding character in the sunflower. *Crop Sci.* (6) : 159, 1966.
- Eric, D.P. and D.G. Dorrel (1975). Breeding for sunflower production. Production oil seed and pulse crops in western Canada. John T. Harapiala pp 185- 201, 1975.
- Fick, G.N. (1975). Heritability of oil content in sunflower. *Crop Sci.* 15 : 77- 78, (1975).
- (1978). Inheritance of oil content in sunflower. In *Agronomy Abstr.* Madison, USA American Soci. of Agron. USDA, Fargo, N. Dak.
(C.F. Plant Br. Abstr. 58 (1) : 548).
- Fiiredi, J. and J. Frank (1981). Investigations into the combining ability of sunflower lines and the genetic analysis of combinations by the Griffing- method. *Novenytermalek.* Tom. 30. No. 4. Hungary.
- Ge, C.F. (1982). Study on the utilization of hybrid vigor in sunflower. *Pl. Br. Cult. Acad. Agric. Sci.* (1981). No. 3 11- 14 China.
(C.F. Plant Br. Abstr. 52 (5). 5957. 1982).
- George, D.L. (1984). Genotype x environmental interactions for seed yield, oil yield and oil percentage in sunflower. Adelaide, Australia,

- Organizing Committee, standing Committee on Agric. 186- 188 [En, 6 ref.] Queensland, Dep. Primary Industries, Hermitage Research Station, Warwick Queensland, Australia.
- Giriraje, K.; S.R. Hiremath; K. Seenappa (1989). Combining ability of converted male- sterile lines of sunflower (*Helianthus annuus* L.). Indian J. of genetics and Plant Br. 47 (3) 315- 317 [En, 4 ref.] Sunflower Scheme, Univ. Agric. Sci., Bangalore, 560065, India.
- Gorbachenko, F.I. (1980). Manifestation of heterosis after crossing short and tall forms of sunflower. J. Vest. Sel' Sho Kohozaistvenno: N. (1979). (8), 35- 40. (C.F. Plant Br. Abstr. 50 (1) : 634).
- Green, V. E. (1980). Correlation and path coefficient analysis of the components of yield in sunflower cultivars (*Helianthus annuus* L.). IX Conference a International del Girosol Tomo II Torremolinas (MALZGAESPANA) 8- 13 de Junio de 1980, p. 12- 21.
- Griffing, J.B. (1956). Concept of general and specific combining ability in relation to diallel crossing system. Australian J. of Biol. Sci. 9 : 463- 493.
- Gumenyuk, A.D. (1987). The basic characteristic oil yield, Maslich. Kul'tury (3), 33- 34 U.S.S.R. (C.F. Plant Br. Abstr. (56), 569).
- Habura, E.C. (1958). Heterosis in yield characters in sunflower. zuchter 28: 285- 287. (C.F. Oil Seed and Pulse Crops in Western canada, John Harapiala 1975).
- Hasan, A.A. (1993). Effect of some cultural practices on sunflower. M. Sc. Thesis, in Agronomy Fac. of Agric., Minofiya Univ.
- Hayman, B. (1954 a). The analysis of variance of diallel tables. Biometertics, 10 : 235- 244.

- (1954 b). The theory and analysis of diallel crosses. *Genetics*, 39: 789- 809.
- Heikal, M.A. (1976). Variation and covarianc in sunflower (*Helianthus annuus* L.). M. Sc. Thesis, Fac. of Agric. Cairo Univ.
- Heiser, C.B. (1951). The sunflower among North American Indians. (C.F. Oilseed and Pulse Crops. In Western Canada 1975. John Harapoak).
- Hunter, M.N. and A.N.Mccosker (1982). Interactions between moisture supply, N and P, in the performance of an open pollinated and a hybrid sunflower variety in a semi- arid environment. Proceedings of the 10 th International sunflower conference. Australia Sunflower Association, 1982: 88- 91.
- Jinks, J.L. (1954). The analysis of continuous variation in a diallel cross of *Nicotians rustica* varieties. *Genetics*, 39: 767- 788.
- (1955). A survey of the genetical basis of heterosis in a variety of diallel crosses. *Heridity*, 9 : 223- 238.
- and D.I. Hayman (1953). The analysis of diallel crosses. *Maize genetics Coop. Newsl.* 27 : 48- 54.
- Jukic, V. (1990). Evaluation of the value of sunflower lines and the way of inheriting of some quantitative characteristics by diallel crossing system. Osiiek, Yugoslavia, Polioprivredni. Institute. P. 50.
- Kadkol, G.P.; I.I. Anand; and R.P. Sharma (1986). Heterosis and combining ability for oil yield in sunflower. *Indian J. of Genetics* 44 (3) 447.

Khanna, K.P. (1972). Factors affecting the production of filled seeds. *Euphytica* 21 : 384- 387.

Kongchuensin, S. and R. Marinkovic (1984). Inheritance and heritability of plant height in diallel crossing in sunflower. *Thai. J. Agric. Sci.* 1984 (170 51- 83.
(C.f. Plant Br. Abstr. 55. 3817.).

Kosteloot, J.A.; J. Heursel and F.M. Poundwels (1985). Estimation of the heritability and genetic variation in sunflower (*Helianthus annuus* L.). *Helia* No. 8, 17- 20 [En, Fr, es, 12 ref.] Fac. Agric. Sci., Mardel Plata Univ., Balcarce Argentina.

Kovacik, A. and V. Skaloud (1972). The proportion of the variability component caused by the environmental and the correlations of economically important properties and characters of the sunflower (*Helianthus annuus* L.). *Scientia Agric. Bohemoslovaca*, 4, 4 : 249- 261.

----- and ----- (1990). Results of inheritance evaluation of agronomic importance traits in sunflower. *Helia*, 13, No. 13, pp. 41- 46.

Kukosk, M.V. (1986). Breeding interline sunflower hybrid under Modavian Conditions.
(C.F. Plant Br. Abstr. 56 (8). 7088).

Kurnik, E. and J. Varsanyi (1963). Variation of the sunflower kernel crops, and of its oil and protein content within a head. *Iregszepese Bull*, 3 : 24- 29.
(C.F. Field Crop Abstr. 18 : 943, 1965).

- Lakshmanalah, V.H. (1981). Genetic variability and association of morphological characters with seed yield and oil content in sunflower. *Mysora J. of Agric. Sci.* 1980. 14 (2) 259. (C.F. Plant Br. Abstr. 51. 9886).
- Manjunath, A. (1978). Genetics of seed yield, oil content and other quantitative characters in sunflower. *Mysore J. of Agric. Sci.* 12, 14 : 667- 668.
- Marinkovic, R. (1993). Components of genetic variability for characters affecting oil yield of sunflower (*Helianthus annuus* L.). Faculty of Agric. Inst. of field and Vegetable Crops. 21000 Novi Sad, Yugoslavia, *J. of genetic and Breeding* Vol. 47 (4) : 289- 293.
- Mather, K. and J.L. Jinks (1971). *Biometrical Genetics* (2 nd ed.), Chapman and Hall Ltd., London.
- Miller, J.F.; G.N. Fick and J.R. Cedon (1977). Improvement of oil content and quality in sunflower (*Helianthus annuus* L.). In *Agronomy Abstr.* Madison, U.S.A. American Society of Agronomy (1977) 64 (En) USDA, Agric. Res. Serv. NDak. State Univ. U.S.A. (C.F. Plant Br. Abstr. 49 : 87370, 1979).
- , J.J. Hammond and W.W. Roath (1981). Comparison of inbred vs. single- cross tester and estimation of genetic effects in sunflower. *Crop Sci.* 20 (6) 703- 706.
- Monotti, M. (1975). Experimental research on cultural techniques for sunflower. 3- Effects of application of increasing rates of nitrogen, phosphorus and potatssium. *Annalidella Facolta di Agrari, Universita degli Studi di Perugia*, 30: 341- 354.

- Moradi, A. and P. Vojadni (1973). Study of the relationship between leaf surface, grain yield and oil percent in different varieties of sunflower. Proceeding of the 6 th International Sunflower Conference, Bucharect, Romania.
- Moursi, M.A.; N. Nour El- Din; O.H. El- Bagoury and T.G. Behairy (1980). Correlation studies among different sunflower characters. Fac. of Agric., Ain Shams Univ.
- Muirhead, W.; A.A. Low and R.J.G. White (1982). The response of irrigated sunflower cultivars to nitrogen fertilizer. Proceedings of the 10 th International Sunflower Conference. Australian Sunflower Association 1982, 82- 85.
- Natali, A.H. and J.M. Shaikh (1970). Correlation of seed yield with other characters in the sunflower. Proceeding of the 21 st - 22 nd Pakistan Sci. Conference. Rajshahi 1970, part III, lahore, W. Pakistan, A55. (C.F. Plant Br. Abstr. 42 : 5958, 1972).
- Niklyae, V.S. (1976). Soviet sunflower varieties in Republic of Guinea. Studies of mangeriol practices proceeding of the 7 th International Sunflower Conference, 27, June- 3 July, 1976. Krasndor, USSR. Vol. 2.
- Nuthan, D. (1981). Study of heterosis and stability index in sunflower (*Helianthus annuus* L.). Mysore J. of Agric. Sci. 14 (1) 125- 126. 1980. (C.F. Plant Br. Abstr. 51. 5379).

- Oka, H.I. and F. Campus (1974). Breeding behavior and genetic variations of sunflower (*Helianthus annuus* L.). Sabras J. S. (1) : 61- 67.
(C.F. Plant Br. Abstr. 44 (11) : 7979).
- Omran, A.O.; A.A. Megahed and F.H. Nofal (1979). Effect of sowing dates on the variability and correlations of mature sunflower characters. Agric. Res. Rev. 57 (9) : 75- 85.
- Pathak, A.R.; B. Sing and M.U. Kukadia (1987). Combining analysis in sunflower. Indian J. of Heredity 1985, 17 (3). 12, 22.
(C.F. Plant Br. Abstr. 56. 566).
- Pirani, V. (1983). Sunflower special effect of nitrogen fertilizer on sunflower cultivars of different genetic characteristics. Informatore Agraric, 39 (20): 25935- 25944.
- Pustovoit, V.S. (1973). Results of sunflower breeding and seed production. Soviet Genetics, 8, 12 : 1501- 1507.
- Putt, E.D. (1943). Association of seed yield and oil content with other characters in the sunflower. Sci. Agric. 23 : 377- 383.
- (1966). Heterosis, combining ability and predicted synthetic from a diallel cross in sunflower (*Helianthus annuus* L.). Can J. Plant Sci. 46 : 59- 67.
- and D.G. Dorell (1977). Breeding for sunflower production Oil seed and pulse crops in Western Canada John T. Harapiala, 1977.

- Rao, P.K. (1981). Diallel analysis of ten quantitative characters in sunflower. Thesis Abst. 1980 6(2) 111- 112. Univ. Agric. Bangalore, India.
(C.F. Plant Br. Abstr. 51. 10821).
- and P. Anand (1981). Diallel analysis of ten characters in sunflower. Mysore J. of Agric. Sci. 1980 14(2) 260.
(C.F. Plant Br. Abstr. 51. 9887).
- Reddy, T.N. (1981). Studies of heritability and genetic advance in certain hybrids of sunflower. Indian J. of Heredity 1979 11 (3). 21- 23.
(C.F. Plant Br. Abstr. 51. 9068).
- Rincon, C.I.; I. Palafox and D.E. Barrada (1986). Recurrent selection for g.c.a. of S1 lines in sunflower. Thesis Univ. Autonoma chap. Mexico 1983, 8 (42) 186.
(C.F. Plant Br. Abstr. 56. 563).
- Ross, A.M. (1939). Some morphological characters of (*Helianthus annuus* L.) and their relationship to the yield of seed and oil. Sci. Agric. 19 : 372- 379.
- Rudranaik, V.; S.R. Hiremath and K. Giriraj (1990). Combining ability for yield and its attributes in sunflower. J. of oil seeds Research, 7 (2) 65- 68 [En, 5 ref.] Univ. of Agric. Sci. GKVK, Bangalore 560065, India.
- Salih, M.A. (1987). Breeding studies on sunflower (*Helianthus annuus* L.). M. Sc. Thesis in Agronomy Fac. of Agric. Zagazig Univ.
-

- Sanford, J.D.; B.L. Arnold and B.R. Eddleman (1979). Variety performance and response to nitrogen of oil type sunflower in Mississippis. *Agric. and Forestry Experiment Station, Mississippi State Univ.*, 4 (16)
- Seetharam, A. (1977). Performance of hybrids of sunflower produced by means of cytoplasmic male sterility. *Sabaras J.* 9 (1) : 51- 55.
(C.F. Genetic Abstr. 10 (1) : 601, 1978).
- Serry, M.H. (1964). Report on sunflower research and production in the A.R. of Egypt. First International Sunflower Conference. Tune 17-18 in Texas A and M University.
- Shabana, R. (1974). Genetic variability of the yield components of oil in different sunflower varieties and inbred lines. Ph. D. Thesis, Novi Sad, Yugoslavia.
- (1978). Effect of nitrogen application on seed yield, oil content and other characteristics of sunflower. Third General Conference and 10 th Anniv. of the Associ. for the Advancement of Agric. Sci. in Africa (AAASA) on Food Crisis and Agric. Prod. in Africa, Problems. Policies and Solutions, Vol. III: 113- 124, 9- 15 April; Ibadan, Nigeria.
- Sheriff, N.M. and R. Appadurai (1987). genetic variability in some quantitative traits of sunflower. *Madras Agric. J.* (1987) 72 (9) 539-540.
(C.F. Plant Br. Abstr. 56. 7287).

- , ----- and M. Rangasany (1987). Estimation the heritability of some character in sunflower through offspring covariance. Madras.Agric.J. (1985). 471 (12) 832- 834.
(C.F. Plant Breed. Abstr. 56. 9859).
- Shrinivasa, K. (1983). Inheritance of fertility restoration and oil content in sunflower (*Helianthus annuus* L.). Thesis Abstr. (1983). 8 (1) 70 71.
(C.F. Plant Br. Abstr. 53. 790).
- Shvestsova, V.P. (1982). Effect of inbreeding and hybridization on the fatty acid composition of sunflower seed oil. Ref. Zh. (60) : 38- 40.
(C.F. Plant Br. Abstr. 52. 5252).
- Singh, K.R. and S.K. Chaudhary (1977). Biometrical Methods in Quantitative Genetic Analysis.
- Singh, P.P.; J.N. Sachan and D. Singh (1977). Variability and correlation in sunflower (*Helianthus annuus* L.). Pantnagar J. of Res. 2, 1 : 27-30.
- Singh, H.; T.P. Yadava and A.K. Yadava (1981). Heterosis in inter-varietal crosses of sunflower. Crop Improvement 1978 5 (1) 84- 85.
(C.F. Plant Br. Abstr. 51. 2412).
- Singh, S.B.; K.S. Labana; and D.S. Virk (1989). Detection of epistasis, additive and dominance variation in sunflower (*Helianthus annuus* L.). Indian J. of Genetics and Plant Br. 47 (3) 243- 247 [En, 14 ref.] Dep. pl. Br., Punjab Agric. Univ., Ludhiana 141004, India.

- Skoric, D. (1980). Production and economic values of new NS sunflower hybrids. Sunflower Newsletter (1977). 1 (2) 5-8.
(C.F. Plant Br. Abstr. 50 8715).
- (1981). Desired model of sunflower hybrid and newly developed NS- sunflower hybrids. International course production and processing of sunflower Novi Sad. Yugoslavia 1981.
- Snedecor, G.W. and W.G Cochran (1967). Statistical Methods. 6 th Ed. The Iowa State Univ. Press, Ames, Iowa.
- Soltani, E. and Y. Arsh (1990). Correlation between oil content and 1000 seed weight and their narrow sense heritability on sunflower variety (ZARYA) in dry farming conditions. Nahal-o- Bazr- Research J. seed and Plant Improvement Institute Karai (Iran Islamic Republic). V. 6 (1, 20 p. 43- 48.
- Sprague, G.F. and L.A. Tatum (1942). general vs. specific combining ability in single crosses of corn. J. Amer. Soc. Agron. 34 : 923- 932.
- Stephens, G.F. and M.D. Finker (1953). Natural crossing in cotton. Econ. Botany, 7, 257- 269.
- Stoyanova, I. (1971). Inheritance of some sunflower characters in the F1. genetic Selek. 4 (1) : 13- 14.
(C. F. Plant Br. Abstr. 42, (1) : 1069).
- Sudhakar, D. (1981). Heterosis and combining ability in sunflower (*Helianthus annuus* L). Mysore J. of Agric. Sci. 1979 (13) 476.
(C.F. Plant Br. Abstr. 51. 3470).

- Taha, R.S. (1993). Heterosis and combining ability in sunflower (*Helianthus annuus* L). Egypt. J. Appl. Sci., 8 (12).
- (1994). Gene action and heritability estimates in sunflower (*Helianthus annuus* L). Proc. 6 th Conf. Agron., Al Azhar Univ. Cairo, Egypt. Sept. (1994) Vol. 11. 817- 832.
- Toit, D.D. (1981). An evaluation of sunflower hybrids and open-pollinated cultivars in respect of certain important agronomic characteristics. Teg. Med. Van 1979, No. 164 pp. 8. (C.F. Plant Br. Abstr. 51. 6443).
- Tomov, T. (1976). Effect of fertilizers on sunflower yield. Pochroznanie i Agrokhimiya II: 101- 108. (C.F. Field Crop Abstr. (27): 9014, 1976).
- Tuberosa, R.; E. Alba; I. Grerio and U. Paradisi (1983). Evaluation of combining ability in pure line of sunflower Sementi. Elett. 1982, 28 (5) 19- 22. (C.F. Plant Br. Abstr. 53. 3308).
- Unrau, J. (1947). Heterosis in relation to sunflower breeding. Sci. Agric. 27 : 414- 427. Oil Seed and Pulse Crops in western Canada. John T Harapiak 1975.
- Varshney, S.K. and R. Singh (1977). Correlation and path coefficient analysis in sunflower (*Helianthus annuus* L). Pantngar J. of Res. 2 (2) : 147- 149. (C.F. Plant Br. Abstr. 3.).

- Velazquez Cagal, M.; A. Carballo and J.I. Rincon Carreon (1991).
Stability of sunflower (*Helianthus annuus* L) in the Valley of Mexico. Revista chapingo, 15 (71- 72) 79- 83 [Es, en, 15 ref.]
Departamento de fitotecnica, Univ. Autonoma de Chapingo, Chapingo, Mexico.
- Velkov, N.N. (1970). Inheritance of stem height in sunflower (*Helianthus annuus* L). Genetika Selekt 3 : 393 - 401.
(C.F. Plant Br. Abstr. 41 (4) : 8607).
- Volf, V.G. and L.P. Dumacheva (1973). Inheritance of the parental on variation in economically valuable characters in sunflower hybrids. Selektsiya Semenovodstva Resp. Mezhved. Temt. Nauch, Sb. 24 : 36- 43.
(C. F. Plant Br. Abstr. 45 : 7598, 1975).
- Vranceanu, A.V. (1983). The performance of sunflower hybrids in production actual and prospective progress. general Siplant tehnice 1984, 48, 67- 74.
(C.F. Plant Br. Abstr. 53. 8993).
- and N. Pirvu (1988). Genetic advance realized in breeding sunflower in Romania. Analele Institutului de Cercetari Pentru Cereale si Plante Techice, Fundulea, 56, 103- 114.
- Vulp, V.V. (1969). A study of some sunflower hybrids produced through male sterility and fertility restoration.

(C.F. Plant Br. Abstr. 39. 5330).

Wright, S. (1921). Correlation and causation. *J. Agric. Res.*, 20: 557-785.

----- (1922). The effect of inbreeding and cross- breeding in quinea pigs. III Crosses between highly inbred fat families. U.S. Dept. Agr. Bul. 1121.

----- (1923). The theory of path coefficient, areply to Niles, Criticism. *Genetics*, 8; 239- 255.

----- (1934). The method of path coefficient. *Annals Math. Statis.*, 5: 161- 215.

Zaffaroni, E.; A.A. Schneiter (1991). Sunflower production as influenced by plant type, plant population and row arrangement. *Agr. J. (USA)*. (Jan-Feb. 1991) V. (1) p. 113- 118.

Zali, A.A.; S.B. Yazadi and M.C. Amirshahi (1977). Multiple regression analysis for different quantitative characters on sunflower. In *Agronomy Abstract*. Madison, U.S.A. American Society of Agronomy (1977) 78 (Em) Dep. Agron., Coll. Agric., Tehran Univ. Iran. (C.F. Plant Br. Abstr. 49 (9) : 8285, 1979).

Zielinska, A. and A. Zielinska (1979). A comparison of sunflower cultivars grown as catch crops with different rates of fertilizer nitrogen. *Zeszyt Naukowe Akademii Rolniczo- Technicznej Wolsztynie Rolnictwo*, 1979, 26 : 119- 126.

Zubriski, J.C.; Deibert and R.P. Schneider (1979). Nitrogen: a limiting factor in sunflower production on non- fallow soils. *North Dakota Farm Research*, 37 (3): 19- 21.