

CONTENTS

| | <i>Page</i> |
|---|--------------------|
| I List of Tables | i |
| II List of Figures | vii |
| I- Introduction | 1 |
| II- Review of literature | 4 |
| III- Material and Methods | 79 |
| IV- results and Discussion | 89 |
| <i>Part I: Stem cuttings studies</i> | 89 |
| IV : 1.1. Mango | |
| IV: 1.1.1. Seasonal changes in some chemical constituent of cuttings as related to excising / collecting date and treatment of mother tree: | 89 |
| IV.1.1.1.A- Total indoles | 89 |
| IV.1.1.1.B- Total phenoles. | 92 |
| IV.1.1.2. Vegetative growth. | 95 |
| IV.1.1.2.1 Rooting Percentage. | 95 |
| IV. 1.1.2.2 Number of roots per cutting. | 104 |
| IV.1.1.2.3 Average root per cutting | 109 |
| IV.1.1.2.4 Number of shoots per cuttings | 115 |
| IV.1.1.2.5 Average shoot length per cutting | 119 |
| IV.1.1.2.6 Number of leaves per rooted cutting. | 125 |
| IV.1.1.2.7 Survival percentage. | 131 |
| IV.1.1.3 Anatomy of initiation and adventitious root formation. | 139 |
| IV.1.2 Pecan | 144 |
| IV. 1.2.1. Seasonal changes in some chemical constituents of cuttings as related to existing "collecting" date and treated mother trees. | 144 |
| IV.1.2.a- Changes in total indoles | 145 |
| IV.1.2.1. b- Changes in total phenols | 147 |
| IV.1.2.2. Vegetative growth. | 149 |
| IV.1.2.2.1. Percentage of rooting | 149 |
| IV.1.2.2. Number of development roots per cutting | 159 |
| IV.1.2.2.3 Average length of initiated roots. | 166 |
| IV.1.2.2.4 Number of developed shoots per cutting | 171 |
| IV.1.2.2.5 Average shoot length (cm) cutting | 176 |
| IV.1.2.2.6 Number of leaves per rooted cutting. | 182 |
| IV.1.2.3 Anatomy of initiation and adventitious root formation | 188 |
| Part II: Invitro propagation. | 194 |
| IV.II.I. Mango | |
| IV.II.1.1. Initiation and establishment of Taymour mango shoot tip and nodal cultures. | 194 |
| IV.II.1.1. A: Surface sterilization | 197 |
| IV.II. 1.1.b. Survival and establishment | 201 |
| IV. II.1.2. Proliferation stage of Taymour mango. | 205 |
| IV.II.1.3 Rooting stage.. | |

| | |
|---|-----|
| IV.II.2. Pecan | 207 |
| IV.II.2.1. Initiation and establishment of curtis pecan shoot tip and nodal cultures. | 207 |
| IV.II.2.1.a: Surface sterilization | 207 |
| IV.II.2.1.b: Survival and establishment | 209 |
| IV. II.2.2. Proliferation stage. | 211 |
| V. IV.II.2.3: Rooting stage. | 214 |
| V. Summary and Conclusions | 216 |
| VI. Literature Cited | 228 |
| VII. Arabic Summary | - |

List of Tables

- Table (1):** Total indoles (mg/ 199 gm.D.wt) of Taymour mango cuttings in response to mother plant treatments and collecting date and their combinations during two successive, 1995-1996 and 1996-1997 seasons. 90
- Table (2):** Total phenols (mg/ 100 gm D.wt.) of Taymour mango cuttings in response to mother plant treatments and collecting date and their combinations during two successive, 1995-1996 and 1996-1997 seasons. 93
- Table (3):** Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on rooting percentage after (14) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons.
- 3-a- Specific effect of mother tree treatments
 - 3-b- Specific effect of collecting date.
 - 3-c- Specific effect of preplanting treatments.
 - 3-d- Specific effect of wounding 99
 - 3-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on rooting percentage in Taymour mango cuttings during 1995-1996 season. 100
 - 3-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on rooting percentage in Taymour mango cuttings during 1996-1997 season. 101
- Table (4):** Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on number of roots after (14) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons.
- 4-a- Specific effect of mother tree treatments
 - 4-b- Specific effect of collecting date.
 - 4-c- Specific effect of preplanting treatments.
 - 4-d- Specific effect of wounding 106
 - 4-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of roots in Taymour mango cuttings during 1995-1996 season. 107
 - 4-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of roots in Taymour mango cuttings during 1996-1997 season. 108

Table (5): Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on average root length after (14) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons.

- 5-a- Specific effect of mother tree treatments
- 5-b- Specific effect of collecting date.
- 5-c- Specific effect of preplanting treatments.
- 5-d- Specific effect of wounding 112
- 5-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average root length in Taymour mango cuttings during 1995-1996 season.
- 5-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average root length in Taymour mango cuttings during 1996-1997 season. 113

Table (6): Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on number of shoots after (14) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons.

- 6-a- Specific effect of mother tree treatments
- 6-b- Specific effect of collecting date.
- 6-c- Specific effect of preplanting treatments.
- 6-d- Specific effect of wounding 116
- 6-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of shoots in Taymour mango cuttings during 1995-1996 season. 117
- 6-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of shoots in Taymour mango cuttings during 1996-1997 season. 118

Table (7): Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on average shoot length after (14) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons.

- 7-a- Specific effect of mother tree treatments
- 7-b- Specific effect of collecting date.
- 7-c- Specific effect of preplanting treatments.
- 7-d- Specific effect of wounding 122

7-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average shoot length in Taymour mango cuttings during 1995-1996 seasons. (123).

7-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average shoot length in Taymour mango cuttings during 1996-1997 season. 124.

Table (8): Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on number of leaves after (14) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons. 127

8-a- Specific effect of mother tree treatments

8-b- Specific effect of collecting date.

8-c- Specific effect of preplanting treatments.

8-d- Specific effect of wounding 128

8-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of leaves in Taymour mango cuttings during 1995-1996 season. 129

8-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of leaves in Taymour mango cuttings during 1996-1997 season.

Table (9): Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments (d) on survival percentage after (12) weeks from planting of Taymour mango cuttings during 1995-1996 and 1996-1997 seasons.

9-a- Specific effect of mother tree treatments

9-b- Specific effect of collecting date.

9-c- Specific effect of preplanting treatments.

9-d- Specific effect of wounding 133

9-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on survival percentage in Taymour mango cuttings during 1995-1996 season. 134

9-f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on survival percentage in Taymour mango cuttings during 1996-1997 season. 135

- Table (10):** Total indoles (mg/ 199 gm.D.wt) of Curtis pecan cuttings in response to mother plant treatments and collecting date and their combinations during two successive, 1995-1996 and 1996-1997 seasons. 164
- Table (11):** Total phenols (mg/ 199 gm.D.wt) of Curtis pecan cuttings in response to mother plant treatments and collecting date and their combinations during two successive, 1995-1996 and 1996-1997 seasons. 148
- Table (12) :** Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments on rooting percentage after (11) weeks from planting of Curtis pecan cuttings during 1995-1996 and 1996-1997 seasons.
- 12-a- Specific effect of mother tree treatments
 - 12-b- Specific effect of collecting date.
 - 12-c- Specific effect of preplanting treatments.
 - 12-d- Specific effect of wounding. 154
 - 12-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on rooting percentage in Curtis pecan cuttings during 1995-1996 season. 155
 - 12 -f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on rooting percentage in Curtis pecan cuttings during 1996-1997 season. 156
- Table (13) :** Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments on number of roots after (11) weeks from planting of Curtis pecan cuttings during 1995-1996 and 1996-1997 seasons.
- 13-a- Specific effect of mother tree treatments
 - 13-b- Specific effect of collecting date.
 - 13-c- Specific effect of preplanting treatments.
 - 13-d- Specific effect of wounding 163
 - 13-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of roots in Curtis pecan cuttings during 1995-1996 season. 164
 - 13 -f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of roots in Curtis pecan cuttings during 1996-1997 season. 165
- Table (14) :** Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments on

average root length after (11) weeks from planting of Curtis pecan cuttings during 1995-1996 and 1996-1997 seasons.

14-a- Specific effect of mother tree treatments

14-b- Specific effect of collecting date.

14-c- Specific effect of preplanting treatments.

14-d- Specific effect of wounding 168

14-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average root length in Curtis pecan cuttings during 1995-1996 season. 169

14 -f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average root length in Curtis pecan cuttings during 1996-1997 season. 170

Table (15) : Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments on number of shoots after (11) weeks from planting of Curtis pecan cuttings during 1995-1996 and 1996-1997 seasons.

15-a- Specific effect of mother tree treatments

15-b- Specific effect of collecting date.

15-c- Specific effect of preplanting treatments.

15-d- Specific effect of wounding 173

15-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of shoots in Curtis pecan cuttings during 1995-1996 season. 174

15 -f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of shoots in Curtis pecan cuttings during 1996-1997 season. 175

Table (16) : Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments on average shoot length after (11) weeks from planting of Curtis pecan cuttings during 1995-1996 and 1996-1997 seasons.

16-a- Specific effect of mother tree treatments

16-b- Specific effect of collecting date.

16-c- Specific effect of preplanting treatments.

16-d- Specific effect of wounding 179

16-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting

treatments x wounding) on average shoot length in Curtis pecan cuttings during 1995-1996 season. 180

14 -f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on average shoot length in Curtis pecan cuttings during 1996-1997 season. 181

Table (17): Specific effect of mother tree treatments (a), collecting date (b), preplanting treatments (c) and wounding treatments on number of leaves after (11) weeks from planting of Curtis pecan cuttings during 1995-1996 and 1996-1997 seasons.

17-a- Specific effect of mother tree treatments

17-b- Specific effect of collecting date.

17-c- Specific effect of preplanting treatments.

17-d- Specific effect of wounding 184

17-e- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of leaves in Curtis pecan cuttings during 1995-1996 season. 185

17 -f- Interaction effect of different combinations between (mother tree treatments x collecting date x preplanting treatments x wounding) on number of leaves in Curtis pecan cuttings during 1996-1997 season. 186

Table (18): Effect of surface sterilization agents on the percentage of contamination and necrosis percentages of Taymour mango shoot tip and nodal cutting explants during 1995 and 1996 seasons. 195

Table (19): Effect of medium type and kind of explant on survival of Taymour mango explants during 1995 and 1996 seasons. 198

Table (20): effect of different concentrations of BA added to the M.S. medium supplemented with IBA at 1.0 mg/L on proliferation rate of Taymour mango explants (number and length of proliferated shoots) during 1995 and 1996 seasons. 202

Table (21): Effect of surface sterilization agents on the percentage of contamination and necrosis percentages of Curtis Pecan shoot tip and nodal cutting explants during 1995 and 1996 seasons. 208

Table (22): Effect of medium type and kind of explant on survival of Curtis Pecan explants during 1995 and 1996 seasons. 211

Table (23): effect of different concentrations of BA added to the W.P.M. medium supplemented with IBA at 1.0 mg/L on proliferation rate of Curtis pecan explants (number and length of proliferated shoots) during 1995 and 1996 seasons. 211

LIST OF FIGURES

| | <i>Page</i> |
|---|-------------|
| Fig. (1): Rooting percentage of Taymour Mango cuttings in response to mother plant treatment, collecting date, some preplanting treatments and wounding after 14 weeks from planting under mist during 1995 - 96 and 1996 - 97 seasons. 1.a. Effect of mother tree treatments. 1.b. Effect of collecting date. 1.c. Effect of preplanting treatment. 1.d. Effect of wounding and unwounding. | 102 |
| Fig. (2): Rooting of May collected cuttings from ethrel sprayed Taymour mango trees as influenced by the different preplanting treatments. 0 = control 1 = 10,000 ppm IBA 2 = 7500 ppm IBA 3 = 5000 ppm IBA 4- 2500 ppm IBA. | |
| Fig. (3): Succeded leafy stem cuttings of Taymour mango cultivar taken in May from trees sprayed with ethrel and treated with vitamin mixture. | 130 |
| Fig. (4): Survival percentage of taymour mango rooted cutting as influenced by mother tree treatments, collecting date, preplanting treatments and wounding after "12" weeks from transplanting during 1995 - 96 and 1996 - 97 seasons. 4-a- Effect of mother tree treatments. 4-b- Effect of collecting date. 4-c- Effect of preplanting treatment. 4-d- Effect of wounding and unwounding. | 136 |
| Fig. (5): Succeded wounding leafy terminal stem cutting of Taymour mango trees sprayed with ethrel, collected in October and treated with IBA at 10.000 ppm. 1- after 5 months from transplanting. 2- after 12 months from transplanting. | 138 |
| Fig. (6): Cross section of unwounding stem cutting of Taymour at planting showing the different layers of the stem : Epiderms (E), Sclernchymatic cells (Sc), Cortex (Co), Phloem (Ph), Cambium (Ca), Xylem (X) and pith (Pi). (X = 3.2 x 10). | 141 |

Fig. (7): Cross section of wounding stem cutting of Taymour mango cv. showing a mass of callus cells was noticed (2 weeks after planting) from wounding zone. (X = 3.2 x 10).

Fig. (8): Cross section of unwounding stem cutting of Taymour mango showing root initiation from the cells of the outer layer of pith parenchyma (10 weeks after planting) xylem and phloem tissue were still undistinguished the development of root. Tip of root primordium is surrounded by the lignified cells at the inner layer of the cortex which prevents emergence of adventitious roots. (X = 3.2 x 10).

Fig. (9): Cross section of wounding stem cutting of Taymour mango cv. showing root initiation from wounding zone (6 weeks after planting). (X = 3.2 x 10).

142

Fig.(10): Cross section of Taymour mango cutting showing the development of adventitious root through the conductive system and reached to the cortex zone as ready to emerge (10 weeks after planting). (X = 3.2 x 10).

Fig. (11): Cross section of Taymour mango cutting showing induced connection between vascular system in the new adventitious roots and vascular bundles of cutting (x = 3.2 x 10).

143

Fig. (12): Rooting percentage of Curtis pecan cuttings in response to mother plant treatment, collecting date, some preplanting treatments and wounding after 11 weeks from planting under mist during 1995 - 96 and 1996 - 97 seasons.

12-a - Effect of mother tree treatments.

12-b- Effect of collecting date.

12-c- Effect of preplanting treatments.

12-d- Effect of wounding and unwounding.

157

Fig. (13): Rooting of May collected cuttings from ethrel sprayed Curtis pecan trees as influenced by the differential preplanting treatments (Wounding application x dipping in IBA and vitamin mixture solution).

187

0 = Control (water). 1= IBA at 1000 ppm.

2 = IBA at 2500 ppm. 3 = IBA at 5000 ppm.

4= Vitamin mixture.

- Fig. (14):** Cross section of unwounding stem cutting of Curtis pecan at planting showing the different layers of the stem :
Priderm (Pr), Sclernchymatic cells (Sc), Cortex (Co), phloem (Ph), Cambium (Ca), Xylem (X) and Pith (Pi). (X = 3.2 X 10) 191
- Fig. (15):** Cross section of unwounding stem cutting of Curtis pecan showing root initiation from the cells of the outer layers of pith paranchyma, was noticed 2-weeks after planting(X = 3.2 x 10).
- Fig. (16):** Cross section of wounded stem cutting of curtis pecan showing the development of adventitious root through the conductive system and reached to the cambium, was noticed 4 - weeks after planting from wounding Zone. (X = 3.2 x 10). 192
- Fig. (17):** Cross section of wounded stem cutting of Curtis pecan showing an adventitious root reached to cortex and priderm tissue from the wounding zone (8 weeks after planting), the developing of adventitious roots through the conductive system and reached to the cortex zone as ready to emeerge. (X = 3.2 x 10).
- Fig. (18):** Cross section of wounded stem cutting of Curtis pecan showing that lignified layers in the priderm and accumulation of resine around the root tips prevent emergence of adventitious roots after transplanting of cutting (X = 3.2 x 10). 193
- Fig. (19):** The normal of Taymour mango explants during establishment stage of (1) shoot tip; (2) single node cutting. 199
19 - a- After 4 weeks from culturing on M.S. medium.
19 - b- After 6 weeks from culturing on M.S. medium.
19 - c- After 8 weeks from culturing on M.S. medium.
- Fig. (20):** Taymour mango proliferated shoots driven from the explants; (1) shoot tip (2) single node cuttring throughout multiplication stage.
20-a- after 2 weeks from subculture.
20-b- after 4 weeks from subculture.
20-c- after 8 weeks from subculture.
20-d- after 10 weeks from subculture. 204
- Fig. (21):** Taymour mango shoots on rooting medium after 20 weeks old (2 week after subculture) but failed to root. 206

Fig. (22): The normal of Curtis pecan explants during establishment stage of (1) shoot tip; (2) single node cutting.

22-a-after 4 weeks from culturing on W.P.M. medium.

22-b-after 6 weeks from culturing on W.P.M. medium.

22-c-after 8 weeks from culturing on W.P.M. medium.

210

Fig. (23): Curtis pecan proliferated shoots derived from the explants; (1) shoot tip; (2) single node cutting in multiplication stage after 8 weeks from subculture.

212

Fig. (24): Curtis pecan shoots on rooting medium after 20 weeks old (4 weeks from subculture) but failed to root.

215