

REFERENCE

REFERENCE

- Abdel- Hamid, M.S. (2001).** Genotype X environments interaction and stability parameters for grain yield and other agronomic traits of white maize hybrids. *J. Agric. Sci. Mansoura Univ.*, 26(6): 3367-3379.
- Agrama, H.A.; C.G. Tarek and S.F. Houssin (1997).** Genetic linkage maps in maize based on RAPD and RFLP markers. *Journal of Genetics and Breeding*, 51: 225-233.
- Ajmone-Marsan, P.; G. Eigdy; G. Monfredini and M. Motto (1993).** Identification and mapping of RAPD markers in maize. *Maize Genet. Newslet.*, 67: 27-28.
- Association of Official Seed Analysis (1983).** Seed Vigour Testing Handbook. Contribution No. 32 to the Handbook on Seed Testing.
- ASTM (1975).** Standard Test Method for Water Using Karl Fischer Reagent. 75 (reapproved 1986). The Society: Philadelphia, PA., USA.
- A.O.A.C. (1990).** Official Methods of Analysis of the Association of Official Agricultural Chemists. 15th Ed. Washington D.C., U.S.A.
- Banchero, C.B.; N. Bartoloni; M.A. Rapela; G. Marrassini and S. Renteria (2000).** Determination of discriminatory traits in maize hybrids. *Agronomie*, 20(8): 875-884.
- Basha, H.A. (1994).** Effect of nitrogen fertilizer application time on growth and yield of some maize varieties. *Zagazig J. Agric. Res.*, 21(2): 329-344.
- Batisella, filho; F. Vitti Moro and N.M. DeCarvalho (2002).** Relationships between physical, morphological, and

- physiological characteristics of seeds developed at different positions of the ear of two maize (*Zea mays* L.) hybrids. Seed Sci., & Technol., 30, 79-406
- Beaumont, V.H.; J. Mantet; T.R. Rocheford and J.M. Widholm (1996).** Comparison of RAPD and RFLP markers for mapping F₂ generations in maize (*Zea mays* L.). Theor. Appl. Genet., 93: 606-612.
- Chaplin, M.F. and J.F. Kennedy (1986)** Carbohydrate Analysis, A Practical Approach; Published Approach Series, Oxford, Washington, D.C.
- Chesneaux, M.T. and A. Kobilinsky (1982).** Possibility of identifying maize varieties at the seedling stage. Agronomie, 2(1): 45-53.
- Ching, T.M., H. Saundra; M.C. Boulger and W.E. Kronstad (1997).** Correlation of field emergence rate and seed vigour criteria in barley cultivars. Crop Sci., 17, 312-314.
- Demeke, T.; B. Sasikumar; P. Hucl and R.N. Chibbar (1997).** Random amplified polymorphic DNA (RAPD) in cereal improvement. Maydica, 42: 133-142.
- Duncan, D.B. (1955).** Multiple range and multiple F test. Biometrics, (11): 1-42.
- El-Batal, M.A.; Boshra, N. Ayad and M.H. Abd El-Gawad (1996).** Flowering of diverse recent maize hybrids in relation to heat units and grain yield. Zagazig J. Agric. Res., 23(4): 447-461.
- El-Batal, M.A.; M.H. Abd El-Gawad and A. Abd El-Aziz (1999).** Phonological stages and kernels dry matter in relation to heat units and grain yield for some maize

genotypes. J. Agric. Sci. Mansoura Univ., 24(10): 5333-5345.

El-Hawary, M.I.; Amal H. Selim and A.M. El-Galfy (2003).

Variability assessment of some maize (*Zea mays L.*) elite inbred lines using morphological and molecular methods. Egypt. J. of plant Breed. 7 (1):109-125.

El-Sabbagh, A.A. (1993). Effect of soil moisture stress and

foliar application of zinc on some maize varieties. Ph.D. Thesis, Fac. of Agric., Moshtohor, Zagazig Univ., Egypt.

Fu, J.H.; L.C. Li; X.Z. Liu; Z.B. Peng and C.L. Huang

(1997). Studies on the protocol optimum and preliminary amplification of RAPD in maize. Acta Agronomica Sinica, 23: 56-61.

Gado, H.E. (2000). Estimates of combining ability of some

yellow maize inbred lines in top crosses. J. Agric. Sci. Mansoura Univ., 25 (3): 1495-1510.

Galarreta, J.I. R. and A. Alvarez (1995). Characterization of

Spanish maize populations according to their agro-ecological zone of origin. Anales de la Estacion Experimental de Aula Dei., 21(3): 183-187 (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).

Galarreta, J.I. R. and A. Alvarez (2001). Morphological

classification of maize landraces from Northern Spain. Genetic Resources and Crop Evolution, 48 (4): 391-400.

Gardner, F.P.; R.Valle and D.E. McCloud (1990). Yield

characteristics of ancient aces of maize compared to modern hybrids. Agron. J., 85: 864-868.

- Gastel, A.J.G. Van and J. Kerely (1988).** Introduction to the practical on varietal identification and verification. Quality Seed Production (Ed.) by Gastel, A.J.G. Van and J. Kerely, pp: 127-136. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Gerdes, J.T. and W.F. Tracy (1994).** Diversity of historically important sweet corn inbred as estimated by RFLPs, morphology, isozymes, and pedigree. *Crop Sci.*, 34: 26-33.
- Gouesnard, B.; J. Dallard; A. Panouille and A. Boyat (1997).** Classification of French maize population based on morphological traits. *Agronomie*, 17(9-10): 491-498.
- Guo, J.L.; J.R. Zhao; Y.F. Kong; D.M. Wei; B.S. Lu and Y.D. Wang (2000).** Studies on the utilization of primer combination in maize inbred line identification using DNA fingerprints. *Acta Agriculturae Boreali Sinica*, 15: 27-31. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Hahn, V.; K. Blankenhorn; M. Schwall; A.E. Melchinger; G. Malecot, and P. Jaccard (1995).** Relationships among early European maize inbreds. III: Genetic diversity revealed with RAPD markers and comparison with RFLP and pedigree data. *Maydica*, 40: 299-310.
- Hamilton, T.S., Hamilton, B.C., Johnson, B.C. and H.H. Mitchell (1951).** The dependence of the physical and chemical composition of the corn kernel on soil fertility and cropping system. *Cereal Chem.* 28: 163-176.
- Heun, M. and T. Helentjaris (1993).** Inheritance of RAPD in F_1 hybrids of corn. *Theor. Appl. Genet.*, 85:961-968.

- Hixon, M.M; M.E. Bauer and O.K.Scholz (1980).** An assessment of landsat data acquisition history on identification and area estimation of corn and soybean. Sixth Annual Symposium, Machine Processing of Remotely Sensed Data and Soil Information System and Remote Sensing and Soil Survey, Purdue Univ. 3-6 June, 72-77.
- IBPGR (1991).** International Board of Plant Genetic Resources. Crop Genetic Resources Center. Maize descriptors. Rome, Italy.
- International Seed Testing Association (ISTA 1993).** International Rules for Seed Testing- Seed Sci. and Technol., 21,187-209.
- Katta, Y.S. and M.S. Abd EL-Aty (2002).** Performance and phenotypic-genetic stability estimates of grain yield and its attributes in different environmental conditions of some maize hybrids. J. Agric. Sci. Mansoura Univ., 27(6): 3647-3661
- Kawata, M. and S. Ueda (1999).** Method of individual discrimination by polymerase chain reaction using MI primer. Utilization of transgenic plant and genome analysis in forage crops. Proceedings of an international workshop held at the National Grassland Research Institute, Nishinasuno, Tochigi, Japan, 17-18 March 1998, publ. 1999, 117-121; NGRI Working Report No. 9. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Lanza, L.L.B.; C.L. De-Souza-Jr; L.M.M. Ottoboni; M.L.C. Vieira and A. P. de-Souza (1997).** Genetic distance of

- inbred lines and prediction of maize single-cross performance using RAPD markers. *Theor. Appl. Genet.*, 94: 1023-1030.
- Liu, X.Z.; Z.B. Peng; J.H. Fu; L.C. Li and C.L. Huang (1997).** Heterotic grouping of 15 maize inbreds with RAPD markers. *Scientia Agricultura Sinica*, 30: 44-51. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Liu, X.J.; Y.L. Zheng and J.L. Liu (1999).** Genetic diversity evaluation of maize recurrent selection population with RAPD marker. *Scientia Agricultura Sinica*, 32: 14-20. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Lubberstedt, T.; A.E. Melchinger; C. Dussle; M. Vuylsteke and M. Kuiper (2000).** Relationships among early European maize inbreds: IV. Genetic diversity revealed with AFLP markers and comparison with RFLP, RAPD and pedigree data. *Crop Sci.*, 40: 783-791.
- Marson, P.A.; G. Egidy; G. Monfredini and M. Motto (1993a).** Identification and mapping of RAPD markers in maize. *Maize Genetics Cooperation Newsletter*, 67: 15-16.
- Marson, P.A.; G. Egidy; G. Monfredini; S. Silvestro; M. Motto and S. Di-Silvestro (1993b).** RAPD markers in maize genetic analysis. *Maydica*, 38: 259-264. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Matinelli-Seneme, A.; C. C. Martins and J. Nakagawa (2000).** Water stress effects on germination and on vigor of

flat and round corn seeds cv. AL.34. Revista Brasileira de Sementes, vol. 22, No. 2, pp. 131-138.

- McDonald, M.B. (1990).** Validation of starch gel electrophoresis for corn seed purity determination. Proceedings of the Annual American Seed Trade Association. Corn and Sorghum Research Conference 45, 43-53. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- McDonald, M.B. (1995).** Standardization of seed vigour tests. pp 88-97 in H.A. van de Venter (Ed) ISTA seed vigour testing seminar . Copenhagen, ISTA. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Medna, P.F. and M.J. Filho (1991).** Evaluation of physiological quality of maize (*Zea mays L.*) seeds. Seed Abstracts, 14, 451.
- Moeller, D.A. and B.A. Schaal (1999).** Genetic relationships among Native American accessions of the great plains assisted by RAPD. Theor. Appl. Genet., 99: 1061-1067.
- Moreno-Martinez, E.; M. E. Vazquez-Badillo; A Rivera; R. Navarete and F. Esquivel- Villagrana (1998).** Effect of seed shape and size on germination of corn (*Zea mays L.*) stored under adverse conditions. Seed Science and Technology, 26 (2): 439- 448.
- Mowafy, S.A.E. (2003).** Response of some maize hybrids to nitrogen fertilizer splitting under drip irrigation system in sandy soils. Zagazig J. Agric. Res., 30(1): 17-34.
- Munoz, A.; A. Orozco; A. Carloallo and A. Santos (1995).** Varietal pattern of maize in the south-east region of the Purepecha Mountainrang. I- Important variables used in its

- definition. *Revista Fitotenia Mexicana*, 18(2): 163-173 (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Nawar, A.A.; S.A. EL-Shamarka and E.A. EL-Absawy (2002).** Diallel analysis of some agronomic traits of maize. *J. Agric. Sci. Mansoura Univ.*, 27(11): 7203-7213.
- Ni, B.; M.R. Paulsen and J.F. Reid (1997).** Corn kernel crown shape identification using image processing. *Transaction of the ASAE*, 40(3): 833-838. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Oraby, F.T. and A.A. Sarhan (2002).** Proper agronomic practices required to maximize productivity of some maize varieties in old and reclaimed soils. IV- Effect of seedling methods on growth and yield of some maize hybrids. *Zagazig J. Agric. Res.*, 29(5): 1457-1470.
- Ortiz, R. and R. Sevilla (1997).** Quantitative descriptors for classification and characterization of highland peruvain maize. *Plant Genetic Resources Newsletter*, 110: 49-52.
- Osipova, E.S.; Y.I. Dolgykh; Z.B. Shamina and S.A. Gostimsky (2000).** RAPD marker variability in maize somaclones produced from inbred A188. *Maize Genetics Cooperation Newsletter*, 74: 52-53. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Osipova, E.S.; Z.G. Kokaeva; A.V. Troitskij; Y.I. Dolgikh; Z.B. Shamina and S.A. Gostimskij (2001).** RAPD analysis of maize somaclones. *Genetika Moskva*, 37: 91-96. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).

- Padma, K.N.; B.M. Reddy; R. Ankaiah and P. Sarada (1996).** Characterization of twenty- three maize inbred lines through seed morphology, leaf stomata studies and response of seedling to added chemicals. *Seed Res.*, 24(1): 15-19.
- Pandy, D.K. (1989).** Short duration accelerated ageing of French bean seeds in hot water. *Seed Sceince and Technology* 17, 107-114.
- Parentoni, S.N.; J.V. Magalhaes; C.A.P. Pacheco; M.X. Santos; T. Abadie; E.E.G. Gama; P.E.O. Guimaraes; W.F. Meirelles; M.A. Lopes; M.J.V. Vasconcelos and E. Paiva (2001).** Heterotic groups based on yield-specific combining ability data and phylogenetic relationship determined by RAPD markers for 28 tropical maize open pollinated varieties. *Euphytica*, 121: 197-208.
- Pejic, I.; P. Ajmone-Marsan; M. Morgante; V. Kozumplik; P. Castiglioni; G. Taramino and M. Motto (1998).** Comparative analysis of genetic similarity among maize inbred lines detected by RFLPs, RAPDs, SSRs, and AFLPs. *Theor. Appl. Genet.*, 97: 1248-1255.
- Rafalski, A.; M. Gidzinska and I. Wisniewska (1998).** PCR-based systems for evaluation of genetic relationships among maize inbreds. *Biuletyn-Instytutu-Hodowli-i-Aklimatyzacji-Roslin*. 1998, No. 208, 131-140. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Ragheb, M.M.; A.A. Bedeer and A.S.A. Gouda (1993).** Effect of row spacing and plant population density on grain yield

- of some maize hybrids. *Zagazig J. Agric. Res.*, 20(2): 581-594.
- Rebourg, C.; B. Gouesnard and A. Charcosset (2001).** Large scale molecular analysis of traditional European maize populations. Relationships with morphological variation. *Heredity*, 86(5): 574-587.
- Revilla, P. and W.F. Tracy (1995).** Morphological characterization and classification of open-pollinated sweet corn cultivars. *J. Amer. Soc. Hort. Sci.*, 120(1): 112-112.
- Rosa, S.D.V.F. da; von E.V.R. Pinho ; M. das G. G. C. Vieira and R. D. Veiga, (2002).** Effect of sample size on some physical parameters of corn ear and physiological quality of seeds. *Ciencia e Agrotecnologia*, 26 (1): 57-65. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Sadek, E.S.; H.E. Gado and M.S. Soliman (2000).** Combining ability and type of gene action for maize grain yield and other attributes. *J. Agric. Sci., Mansoura Univ.*, 25(2): 2491-2502.
- Sadek, E.S.; M.S. Soliman and A.A. Barakat (2003).** Dynamic of yield of fourteen white and yellow maize (*Zea mays*, L.) hybrids grown in Egypt. *J. Agric. Sci., Mansoura Univ.*, 28(2): 759-774.
- Samia, G.A. Mohamed; Sohier, M.S. Amer and S.M. Salama (2002).** Estimating prediction equations of yield and its characters in maize using some macroclimatic and micro environmental factors. *J. Agric. Sci., Mansoura Univ.*, 27(7): 4355-4370.

- Sangakkara, U.R. (1989).** Relationship between seed characters, plant growth and yield parameters of *Phaseolus vulgaris*. J. Agronomy & Crop Sci., 163, 105-105.
- Santiparacha, W; Q. Santiprachho and V. Wongvarodom (1997).** Hybrid corn seed quality and accelerated aging. Seed Sci., and Technology, 25 (2): 203-208.
- Scapim, C.A.; C.G. Carvalho and C.D. Cruz (1995).** A proposal for the classification of coefficients of variation in maize cultivation. Pesquisa Agropecuaria Brasileira, 30(5): 683-686. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Schoorel, F.A. (1956).** The use of soil tests in seed testing. Eleventh International Seed Testing convention, Paris. Report of the activities of the committee on seedling vigour tests. pp.287-302.
- Shieh, G.J. and F.S. Thseng (2002).** Genetic diversity of Tainan-white maize inbred lines and prediction of single cross hybrid performance using RAPD markers. Euphytica, 124: 307-313.
- Sivolap, Y.M.; N.E. Kozhukhova and Y.A. Asyka (1997).** Study of genetic interrelations in maize lines by means of RAPD and zeins. Tsitologiya-i-Genetika, 31: 16-20. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Smith, F.G.(1952).** The mechanism of the tetrazolium reaction in corn embryos. Plant physiol. 27: 445-456.

- Smith, J.S.C. and O.S. Smith (1989a).** The description and assessment of distance between inbred lines of maize: I- The use of morphological traits as descriptors. *Maydica*, 34: 141-150.
- Smith, J.S.C. and O.S. Smith (1989b)** The description and assessment of distance between inbred lines of maize: II- The utility of morphological, biochemical, and genetic descriptors and a scheme for the testing of distinctiveness between inbred lines. . *Maydica*, 34: 151-161.
- Snedecor, G.W. and W.G. Cochran (1969).** Statistical methods. 6th Ed., Iowa State Univ. Press, Ames Iowa, USA: 325-330.
- Solimain, F.H.; A.S. Gouda; M.M. Ragheb and Samia, M. Amer (1995).** Response of maize (*zea mays*, *L.*) hybrids to plant population density under different environmental conditions. *Zagazig J. Agric. Res.*, 22(3): 663-676.
- Steenhoek, L. and C. Precetti (2000).** Vision sizing of seed corn. 2000 ASAE Annual International Meeting, Milwaukee, Wisconsin, USA, 9-12 July 2000 pp: 1-10.
- Sun, G.L.; M. Wiliam; J. Liu; K.J. Kasha and K.P. Pauls (2001).** Microsatellite and RAPD polymorphism Ontario corn hybrids related to the commercial sources and maturity rantings. *Mol. Breeding*, 7: 13-24. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Tekrony, D.M. (1995).** Accelerated ageing. pp 53-72 in H.A. van de Venter (Ed) ISTA seed vigour testing seminar . Copenhagen, ISTA. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).

- Tekrony, D.M. and D.B. Egli (1997).** Relationship between standard germination, accelerated ageing germination and field emergence in soybean. Pp 593-600 in R.H. Ellis; M. Black; A.J. Murdoch, T.D. Hong (Eds) Basic and applied aspects of seed biology. Kluwer Academic. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Torres, S. B. (1998).** Vigor tests to evaluate physiological quality of corn seeds. *Revista Brasileira de Sementes*, 20 (1): 55- 59. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).
- Tottman, D.R. (1987).** The decimal code for the growth stages of cereals. British Crop Protection Council Publication in *Annals of Applied Biology* 100.
- Tyystijarvi, E.; M. Keranen; A. Koski; O. Nevalainen and E.M. Aro (1998).** Chlorophyll fluorescence can be used to identify plant species automatically. Photosynthesis, mechanism and effects. Proc. of the 9th Internl. Cong. on photosynthesis, Budapest, Hungary, 17-22 August, 1998, V: 3857-3860.
- UPOV, (1994).** The International Union for the Protection of New Varieties of Plants Guidelines for the Conduct of Tests for distinctness, uniformity and stability for maize Descriptor no. TG/2/6. Geneva..
- Weber. E.J.(1987).** Lipids of the kernel . pages 311-349 in: corn: Chemistry and Technology. S.A. Watson and P.E. Ramstad, eds. AACC: St. Paul, MN: (C.F. CAB computer

data research, National Agriculture library, Dokki, Giza, Egypt).

Welsh, J. and M. McClelland (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acid Res.* 18: 7213-7218.

Williams, J.G.K.; A.R. Kubelik; K.J. Livak; J.A. Rafalski and S.V. Tingey (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Res.* 18: 6531-6535.

Wook, J.T; S. Kim, C.S. Woo and K. Dalung (2000). Kernel characteristics and germination rate during the grain filling in super sweet corn. *Korean Journal of Crop Science*, 45 (3): 176-180. (C.F. CAB computer data research, National Agriculture library, Dokki, Giza, Egypt).

Wu, M. (2000). Genetic diversity and its relationship to heterosis in maize as revealed by AFLPs and RAPDs. *Maize Genet Newslet.*, 74:43-44.

Zhang, C.L.; Z.L. Sun; D.M. Jin; S.M. Sun; B.T. Guo and B. Wang (1998). Identification of maize inbred lines and validation of genetic relationship among maize inbred lines using RAPD markers. *Maize Genetics Cooperation Newsletter*, 72: 9-10.

Zhao, J.R.; J.L. Guo; Q. Guo; D.M. Wei and Y.F. Kong (1999). Heterotic grouping of China's main maize inbred lines with RAPD marker technique. *Acta Agriculturae Boreali Sinica*, 14: 32-37.