

REFERENCES

IV. REFERENCES

- A.O.A.C. (1995). Official Methods of Analysis of The Association of Official Analytical Chemists. Washington, DC 20044.
- Abd El-Aziz, M.A. (1993). Response of some wheat cultivars to soil salinity. M. Sc. Thesis, Fac. of Agric., Tanta Univ., Egypt.
- Abd El-Ghani, E.K.(2001). Towards *in vitro* selection studies for drought tolerance in wheat. M.Sc. Thesis, Fac. Agric., Alex.Univ., Egypt.
- Abd El-Haleem, A.K.; S.A. Kandil and M.A. Koriem (1995). Growth and yield of six wheat cultivars as affected by different levels of chloride salinization. J. Agric. Sci. Mansoura Univ., 20 (1): 117-126.
- Abu El-Ala, A.E. (1993). Effect of soil salinity and alkalinity on yield and chemical composition of wheat plant. Mansoura, J. Agric., Sci., 24 (5): 2673-2783.
- Ali, A.M.A.S.; A.A. Gomma; O.H.S. Khalil and E.H. Ghanem (1991). Gemmiza 1 and Giza 165, two new bread wheat cultivars in Egypt. Assiut j. Agric Sci., 22 (4)93-102.
- Bahraei ,S. (1996). The use of random amplified polymorphic DNA (RAPD) markers in genetic diversity studies of *Triticum boeoticum* and *T. urartu*. Seed-and-plant. 12: 1, 31-43 (pe), 5(En); 15 ref.
- Banerjee, S.k. and S.Chandre (1974). Modified phenol test for the varieties identification of wheat seed preprint. No. 8-SVI 17th ISTA Congress Warsaw.

VI. REFERENCES

- Barakat, M.N.; M.I. Motawei; S.I. Milad; M.A. Moustafa and Y.H. El-Daoudi. (2000).** Using RAPD markers for evaluating genetic relationships among wheat cultivars. Proc. 9th Conf. Agron., Minu.Univ., Egypt.
- Bassilios, S.I. (1992).** Response of two wheat cultivars (Giza 69 and Sakha 93) to rates and splittin of nitrogen fertilizer .Assiut j. Agric. Sci., 23(2): 165-178.
- Bered, F; N.J.F. Barbosa; B.M. Rocha; F.I.F. Carvalho and B.M. Rocha (2002).** Genetic variability in wheat (*Triticum aestivum* L.) germplasm revealed by RAPD markers. Crop-Breeding and Applied Biotechnology. 2: 4, 499- 505.
- Bered,F.; J.F. Barbosa-Neto; B.M. Rocha; F.I.F. Carvalho and B.M. Rocha; F.I.F. Carvalho (2001).** Genetic variability in common wheat based on morphological traits, coefficients of parentage and RAPDs. Journal-of-New-Seeds. 2001, 3: 2, 73-87; 29 ref.
- Cao,W.G.E. (2002).** Cultivar identification and pedigree assessment of common wheat based on RAPD analysis. Wheat- information-service. No.95, 29-35; 37 ref.
- Chandrashkhar, P.;Joshi and Henry, T. Nauyen. (1993).** Application of the random amplified polymorphic DNA technique for detection of polymorphism among wild and cultivated tetraploid wheats. Plant Molecular Genetics Laboratory, Department of Agronomy, Horticulture, and Enteomology. Mail Stop, 2122 Texas Tech. Univ. Lubbock, Tx 974090,USA.
- Chen, J.; D.R. Sheltonm; B.L. D' Appolonia and K. Khan (1992).** Structural characterization of the carbohydrate

- protein of a glycopeptide from wheat gluten. Cereal chem. 69 (5): 481-484.
- Cooke, R.J. and R.C. Payne (1992).** Handbook of variety testing Electrophoresis for variety identification. The international seed testing association (ISTA), publication, Zurich, Switzerland. PP 3-8.
- Das, J.L.; K. Bose and A.K. Biswas (1992).** Evaluation of seeds and intervarietal relationships revealed by PAGE of seed proteins in wheat. Environmental and Ecology, 10 (2): 18-22.
- De loose, M. and G. Gheysen (1995).** Identification Methods Based on Molecular Techniques. UPOV Document BMT 1312, Geneva pp 22.
- Dhaliwal, H.S.; J.S. Sidhu and J.L. Minocha (1993).** Genetic diversity in diploid and hexaploid wheats as revealed by RAPD markers. Crop-improvement. 20: 1, 17-20; 12 ref.
- Dòvido, R.; O.A. Tanzarella and E. Porceddu (1990).** Rapid and efficient detection of genetic polymorphism in wheat through amplification by polymerase chain reaction. Plant Molecular Biology, 15:169-171.
- Dudley, J.W. (1992).** Theory for identification of marker locus-QTL associations in population by line crosses. Theor. Appl. Genet., 85:101-104.
- Dudley, J.W.; M.A. saghai-Marooof and G.K.Rufener. (1991).** Molecular markers and grouping of parents in corn breeding programs. Crop Sci., 31:718-722.
- Duncan, D.B. (1955).** Multiple range and multiple F test. Biometrics. (11): 1-42.

- Ebrahim, Eman A. (1999).** Chemical studies of some genetic resources of wheat .Ph.D. Thesis, Fac. Agric., Cairo Univ., Egypt.
- El-Bagoury, H. Olfat; A.M. Hegazi; M.T. Mostafa; Kh.T. El-Afandi. (1998).** Influence of organic manure and nitrogen fertilizer on chemical composition and technological characters of wheat under irrigation with saline water. Agron.Dep., Fac., of Agric., Sci. Egyptian Society of Agronomy.
- El-Emam (2004).** Studies on some morphological, physiological and biological characters of some wheat genotypes under saline conditions. Ph.D. Thesis, Fac.Agric., Al-Azhar Univ., Egypt.
- El-Kalla, S.E.; A.A. Kandil; M.S. Suultan; S.A. El-moursy and M. Abd El-Aal (1992).** Response of some wheat varieties to irrigation regime and time of nitrogen fertilization. J. Agric. Sci. Mansoura Univ., 17 (8): 2568-2575.
- Eweis, M.N.A. (2003).** Morphological and biochemical identification of some wheat varieties. Ph.D. Thesis, Fac. of Agric., Al-Azhar Univ. Egypt.
- Fahima, T.; G.L. Sun; A. Beharav; T. Krugman; A. Beiles and E. Nevo. (1999).** RAPD polymorphism of wild emmer wheat populations, *Triticum dicoccoides* in Israel. Theor. Appl. Genet, 98:434-447.
- Ferguson, A. (1980).** Biochemical systematics. Blackie Genetic Resources News Letter, 63: 34-40.
- Figuroa, J.D.C and K. Khan (1993).**Albumin fraction from spring, winter, and soft wheats –protein and associated

- carbohydrate identified by gel filtration chromatography and gel electrophoresis. *Cereal Chem.* 70 (6): 689-695.
- Freitas, LB-de; L. Jerusalinsky; S.L. Bonatto; F.M. Salzano and L.B. Freitas (2000).** Extreme homogeneity among Brazilian wheat genotypes determined by RAPD markers. *Pesquisa- Agropecuaria Brasileira.* 35: 11, 2255-2260; 24 ref.
- Gaber, S.D. (1978).** Identification of seeds of some barley cultivars by chemical staining. *Field Crops Res. Inst., Agric. Res. Center, Ministry of Agric., Egypt*, pp.127-133.
- Gandy, E.K. (1996).** Evaluation of some methods for varietal identification of some crops M..Sc. Thesis, Fac. Agric., Ain Shams Univ., Egypt.
- Ganem,Enayat, H.; O.H.S. Khalil; A.A.A. Gomaa and S.E. Shereef (1991).**The new durum wheat varieties for upper Egypt. *Assiut J. of Agric.Sci.*, 22(3): 208-222.
- Garg, M.; S. Sukhwinder; S. Baldev; S. Kuldeep; H.S. Dhaliwal; S. Singh K. Singh(2001).** Estimates of genetic similarities and DNA fingerprinting of wheats (*Tritium* species) and triticale cultivars using molecular. *Indian-Journal-of-Agricultural-Sciences.* 17:7,438-443;20 ref.
- Ghanem,Enayat H; M.A. Gouda and H.K. Ashour (1987).** Varietal versus environmental effect on certain yield and quality of some promising wheat varieties & lines .*Menia J. Agric. Res.*, 6 Dav. 9 (2): 695-706.
- Guadaluolo, R.; D.S. Bianchi and F. Felber (2001).** Specific genetic markers for wheat, spelt, and four wild relatives:Comparison of isozymes, RAPDs, and wheat microsatellites. *Genome*, Aug., 144 (4):610-621.

- Gubta, R.B.; K.W. Shepherd and F. Macritchie (1991). Genetic control and biochemical properties of some high molecular weight albumins in bread wheats. *J. Cereal Sci.*, 13:221-235.
- Gupta, P. K.; R.K. Varshny; P. C. Sharma and B. Ramesh (1999). Molecular markers and their applications in wheat breeding. *Plant breeding* .118:369-390
- Gurmit Singh (1992). Phenol reaction in some wheat varieties and lines. *Crop Improvement*, 19 (1): 70-71.
- Hassanein, Azziza M. (2005). Molecular fingerprint and genetic purity of some wheat genotypes (*Triticum aestivum* L). Ph.D.Thesis, Fac.Agric., Ain Shams Univ., Egypt.
- Hussein, K.R.F. and H. Stegemann. (1978). Comparison by various electrophoretic methods in polyacrylamide. *A. Acher-und pflanzenbau Agron. J.*, 146: 68-78.
- Hussein, M.; A. Kandil; E. Shok and M. Abd El-Alim (1981). Effect of seeding method and nitrogen fertilizer on some agronomic characters of wheat cultivars (*Triticum aestivum*, L). *Ann. Agric. Sci. Moshtohor*, (15): 3-12.
- Hussein, S.M.A. (1995). Effect of irrigation and microelements on root system, growth and yield of wheat under desert soil conditions. Ph.D.Thesis. Fac. of Agric. Mansoura Univ.
- IBPGR (1985). International Board of Plant Genetic Resources. Crop Genetic Resource Center. Wheat Descriptors AGPG: IBPGR/ 85/ 210. Rome, Italy. PP 1- 19. III.Proceedings of the Third International Triticeae Symposium, Aleppo, Syria, 4-8 May 1997. 1998, 165-170; 20
- ISTA (1996). International Seed Testing Association. *Seed Sci. & Technol.*, 24, supplement, Rules. PP29- 202.

- Joshi, C.P. and H.T. Nguyen. (1993).** RAPD analysis based intervarietal genetic relationship among hexaploid wheats. *Plant Sci., Limerich*, 93(1-2): 95-103.
- Jushi, M.G. and S.K. Banerjee (1970).** Genetic of phenol color reaction in emmer wheats. *Proc. ISTA* 35: 807-811.
- Kandil, A.A. (1985).** Response of some wheat varieties to different rates and forms of nitrogen. *J. Agric Sci. Mansoura Univ .*, 10 (4) : 1041-1053.
- Kandil, A.A. (2000).** Physiological and biochemical study on the responses of some wheat cultivars to salinized water irrigation. *Mansoura J. AGRIC. Sci.*, (1): 13-22.
- Kresoviches, J.Ck. Williams; J.R. McFerson; E.J. Routman and A.B. Schaal. (1992).** Characterization of genetic identifications and relationships of *Brassica oleracea* L. via., a random amplified polymorphic DNA Assay. *Theor. and Appl. Genet.*, 85:190-196.
- Kuz-Mina, K.M. and V.A. Kumakov (1983).** Biological prerequisites for breeding spring wheat for earliness. *Sel-Skokhozyaistvennaya Biologiya*, 10:24-30.
- Laemmli, M.K. (1970).** Cleavage structural protein head bacteriophage T4. *Nature* 227: 680- 685.
- Loarce, Y.; R. Gallego and E. Ferrer (1996).** A comparative analysis of the genetic relationships between rye cultivars using RFLP and RAPD markers. *Euphytica*, 88:107-115.
- Low, C.N. ; J.W. Snape and A.J. Worland (1978).** The genetically relationship between height and yield in wheat. *Heredity*, 40: 133-151.

- Maguire, J.D.; K.M.Steen and K. Grzelok (1975). assification of pacific northwest winter and spring wheat cultivars by phenol reaction.Proc.Assoc.of Seed Anal. 6:143-146.
- Malkawi, H.I.; M.M. Ajouni and A.A. Jaradat(1998).Analysis of genetic diversity of wild emmer and durum wheats by RAPDs. Triticeae
- Mandoulakani, B.A.; B.E.S. Tabatabaei; A.A.S. Bushehri; M.R. Ghannadha and M. Omid (2003). Assessment of genetic diversity among wheat cultivars by RAPD-PCR.Iranian-Journal-of-Agricultural-Sciences. 34(2): 447-454
- Mohamed, S.A; A.A Ali, A.S Fahmy. (2003). Molecular characterization of wheat *Triticum aestivum* cultivars using isoenzyme analysis and RAPD fingerprint. Bulletin-of-the-National-Research-Centre-Cairo. 28(6): 735-747.
- Mohan, M.P.; V. Sathyanarayanan; A. Kumar; M.M. Srivastava and S. Nair (1997). Molecular mapping of resistance. Specific PCR-based marker linked to a gall midge resistance gene (Gmat) in rice. Theor. and appl. Genet, 95(5-6):777-782.
- Munshi, A.; S. Bazaid and A. EL-Tarras (2003). Genetic diversity in diploid KSA wheat as the roles of seed proteins and RAPD-PCR in genotyping variabilities revealed by RAPD analysis. Egypt. J. Genet. Cytol., 32:25-32.
- Nagaoka, T. and Y. Ogihara. (1997). Applicability of inter-simple sequence repeat polymorphisms in wheat for use as DNA marketrs in comparison to RFLP and RAPD markers. Theor. Appl. Genet, 94:597-602.

- Nakamura, T; M. Yamamori; S. Hidaka and T. Hoshino** (1992). Expression of LMW protein in Japanese common wheat (*Triticum aestivum* L.) cultivars. Japanese J. Breeding 42 (3): 681-685.
- Nguyen, H.J.; C.P. Joshi; D. Hoisington and A. McRab.** (1993). RAPD analysis in tetraploid and hexaploid wheats. Progress in Genome Mapping of wheat and Related Species: proceedings of the 3rd public. Workshop of the International Triticeae Initiative, 22-26 september, 1992 CIMMYT, Mexico, 47-50.
- Pommer, E. and K. Oppitz** (1986). Investigations on yield formation in winter wheat cultivars with short and long straw. Agron. J., 156 (3): 153 – 165.
- Qixin-sun, Ni Zhongfu; Liu Zhiyong; G. Jianwei and H. Tiecheng.** (1998). Genetic relationships and diversity among Tibetan wheat, common wheat and European spelt wheat revealed by RAPD markers. Euphytica, 99:205-211.
- Quader, M.; G. Daggard; R. Barrow; S. Walker and M.W. Sutherland.** (2001). Allelopath, DIMBOA production and genetic variability in accessions of *Triticum speltoides*. J. Chen., Ecol. Apr., 27 (4): 747-760.
- Salam, H.A. and S.A.N. Afiah.** (1998). Evaluation and screening some wheat (*Triticum aestivum* L) genotypes in relation to chemical constitutions under saline conditions. Proc. 8th Cnf. Agron. Suez Canal Univ. Ismailia, Egypt, 28-29 Nov. 1-18.
- Selim, Amal H.** (2000). Evaluation of some heat tolerant wheat germplasms for yield and yield quality. Ph.D. Thesis, Fac. Agric Ain Shams Univ., Egypt.

- Selim, Amal H. (2004). Some factors affecting the value of phenol test used in wheat (*triticum aestivum L.*). Egypt J. Agric. Res., 82 (4). 1657- 1664.
- Sendecor, G.W. and W.G. Cochran (1980). Statistical Methods. 6th Ed. The Iowa State Univ. Press. Iowa, USA.
- Shalaby, T.A.; M.A. Tabe; A.M. Omar and A.M. Abo El-Ela (1996). Response of wheat to nitrogen fertilizer under saline conditions. Tanta J. Agric. Res., 22 (3): 376-386.
- Shehata, M.M. (2004). of some wheat (*Triticum vulgare L.*)cultivars. Pakistan-Journal-of-Biological-Sciences. 7(6): 984-994
- Sievert, D.; H.D. Sapirstein and W. Bushuk (1991). Electrophoretic study of some high molecular weight proteins of acetic acid. insoluble residue of wheat flours. Cereal Chem. 68 (5):5 512-515.
- Smith, J.S.C. and O.S. Smith (1989). The description and assessment of distance between inbred lines of maize. II- The utility of morphological, biochemical and genetic descriptors and a scheme for the testing of distinctiveness between inbred lines. Maydica, 34:151-161.
- Smith, O.S. and J.S.C. Smith (1992). Measurement of genetic diversity among maize hybrids; A comparison of isozymic RFLP, Pedigree and Heterosis data Maydica, 37:53-60.
- Tesemma, T.H. ; C. Becker; G. Belay; D. Miliku; E. Bechere and S. Tsegaye (1993). Performance of Ethiopian tetraploid wheat landraces at their collection sites. Euphitica, 71: 221-230.
- Teshale, E.T; B. Sangita; M. Avinash; V.K. Khanna; S. Bansal and A.Mishra (2003). DNA fingerprinting of

- wheat genotypes by RAPD markers. Wheat-Information-Service. No.96,23-27; 16 ref.
- Tomer, R.P.S. and J.D. Maguire (1990)** Seed vigour studies in wheat. Seed Science and Technology, 18: 383- 392.
- Tomer, R.P.S. and M. Steen (1990).** Identification of wheat cultivars by electrophoresis. Seed & Farus. 10 (11-12): 14-15.
- Tottman, D.R. (1987).** The decimal code for the growth stages of cereals. British crop protection council publications in Annals of Applied Biology 100.
- UcVan De Weghe, I. (1991).** Comparative study of electrophoretic methods for cultivar identification of wheat and tritcale. Seed Sci.Technol.19:41-50.
- UPOV. (1990).** Harmonization of stars of exepression and notes of characteristics. The international Union for the protection of new varieties of plants. Document No.TC/26/4 rev. pp 20. Geneva.
- UPOV. (1994).** The International Union for the Protection of new Varieties of plants. Guidelines for the conduct of tests for distinctness, uniformity and stability for wheat No. TG/3/11,pp 36.
- Vierling, A. and H.T. Nguyen (1992).** Use of RAPD markers to determine the genetic relationships of diploid wheat genotypes. Theor-Appl Genet, 84:835-838.
- Wei, J.Z. and R.R.C. Wang (1995).** Genome and species-specific markers and genome relationships of diploid perennial species in Triticeae based on RAPD analysis. Genome, 38:1230-1236.

- Werner, W.E.; J.E. Wiktorowicz and D.D. Kasarda (1994).** Wheat varietal identification by capillary electrophoresis of gliadins and high molecular weight glutenin subunits. *Cereal Chem.* 71 (5): 397-402.
- Williams, J.G.K; A.R. Kubelik; K.J. Livak; J.A. Rafalski and S.V. Tingey (1990).** DNA polymorphism amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Res.* 18: 6531-6535.
- Zhai, W.; Luc, L. Zhu; W. Yang and Q. Zhang (1996).** PCR analysis of half-seeds of cereal crops and its application in markers-assisted and breeding. *Chin. J. Biotechnol.*, 12: 249-255.
- Zhen, Z. and D. Mares (1992).** A simple extraction and one – step SDS – PAGE system for separating AMW and LMW glutemin subunits of wheat and high molecular weight proteins of rye. *J. Cereal Sci.*, 15 (1): 63-78.
- Zheng, Y.L.; Y.Q. Chen; Y.M. Wei; Y.H. Zhou; Z.Q. Zhang; D.C. Liu; X.J. Lan (2001).** Esterase, gliadins and RAPD variations among Sichuan wheat cultivars. *Wheat-Information-Service*. 2001, No.92, 5-8; 21 ref.