

LITERATURE CITED

- A.O.A.C. (1984). Official methods of analysis. 14th. Ed., Association of Official Analytical Chemists. Alington, Virginia, U.S.A.
- Ahmed, H.K. (1999). Heterosis and combining ability in sunflower (*Helianthus annuus* L.). M.Sc. Thesis, Fac. of Agric., Assiut Univ.
- Akhanda, M.A.M. (1979). Evaluation of cultivars and cultural practices of several crops for tropical multiple cropping. Dissertation Abstracts International, B. 39, 3082-3083.
- Alba, E.; P. Barsanti and G. Carrozzo (1979). Analysis of combining ability in new cytoplasmically male-sterile lines and in fertility restorer lines of sunflower. Annali della Facolta di Agraria, Universita di Bari, 31: 363-370.
- Allam, A.Y. and A.H. Galal (1996). Effect of nitrogen fertilization and plant density on yield and quality of sunflower. Assiut J. Agric. Sci., 27(2): 169-176.
- Attia, S.A.M. (1985). Effect of some cultural treatments on sunflower (*Helianthus annuus* L.). Ph.D. Thesis, Fac. of Agric., Al-Azhar Univ.
- Chaudhari, C.S.; Altaf Hussain and J.M. Sheikh (1975). Effect of plant spacings on yield and certain other characters of sunflower var. Ho. L. PAK. J. Bot. 7(1): 41-50.
- Chaudhari, C.S.; A.G. Mahajan; J.T. Paturde (1978). Effect of plant populations and fertility levels on sunflower. Indian J. of Agron. 23(2): 153-1256.
- Chaudhary, S.K. and I.J. Anand. (1986). Heterosis and inbreeding depression in sunflower. Crop Improvement, 11: 15-19.

- Cherzhentseva, M.I. (1985).** Combining ability and inbred lines of sunflower for capitulum diameter. Sel, Skokhozyaistvennaya, Biologia, 10: 15-17. (Computer Research, AN 861655963).
- Cherzhentseva, M.I. (1990).** Evaluation of inbred lines of sunflower for combining ability. Genetika i Seleksionery, Stavropolya Proizvodstvu, 97-104. (Computer Research, AN 911698790).
- Cruz, Q.D. Dela. (1986).** Heterosis and combining ability for yield and yield components in sunflower (*Helianthus annuus* L.). Philippine J. Crop Sci., 11: 171-174.
- Curotti, G.L. and A. Rosania (1971).** Two year results of a spacing trial with sunflower in the Tuscan Maremma. Rivista de Agricoltura Subtropicale e Tropicale. 65 (4-6): 146-164. [C.F. Field Crop. Abst. 25: 2499, 1972].
- Dedio, W. (1993).** Heterosis and prediction of achene oil content in sunflower hybrids from parental lines. Can. J. Plant Sci., 73: 737-742.
- Dozet, B.; J. Atlagic; D. Skoric and R. Marinkovic. (1995).** Development of sunflower inbred lines by interspecific hybridization. Beric 36th Conference About Production and Processing of Oil Crops, Budva, 138-147.
- Dua, R.P. and T.P. Yadava. (1993).** Combining ability in sunflower. Indian J. Genetics & Pl. Breed., 43: 129-136.
- Galeev, G.S. and F.K. Vilichku. (1986).** An effective method for utilizing heterosis in sunflower. Seleksiya, i, Semenovodstvo, USSR, 5: 14-16. (Computer Research, AN 871657393).
- Ge, C.F. (1982).** Study on the utilization of hybrid vigor in sunflower. PI. Br. Cult. Acad. Agric. Sci. (1981). No. 3, 11-14 China. [C.F. Plant Br. Abstr. 52(5): 5957, 1982].

- Geweiff, H.G.M.; Fatma, A.A. Osman and A.Y. Banna (1997).** Response of sunflower to phosphorus and nitrogen fertilization under different plant densities in sandy soil. *Zagazig J. Agric., Res.*, 24(3): 435-448.
- Giriraj, K. and K. Virupakshappa. (1992).** Heterotic effect for seed yield and component characters in sunflower over seasons. *Proceedings of the 13th Intern. Sunflower Conf., Pisia, Italy*, 7: 1043-1047.
- Giriraj, K.; N. Shivaraju and S.R. Hiremath. (1986).** Studies on heterosis and inbreeding depression in selected cross combination of sunflower. *J. Oilseeds Res.*, 3: 37-72.
- Giriraj, K.; S.R. Hiremath and K. Seenappa. (1987).** Combining ability converted male-sterile lines of sunflower (*Helianthus annuus* L.). *Indian J. Genetics & Pl. Breed.*, 47: 315-317.
- Gorbachenko, F.I. (1980).** Manifestation of heterosis after crossing short and tall forms of sunflower. *J. Vest. Sel Sho Kohozaistvenno: N. (1979)*. (8): 35-40. [C.F. Plant Br. Abstr. 50(1): 634].
- Gumenyuk, A.D. (1987).** The basic characteristic oil yield, Maslich. *Kultury* (3): 33-34 U.S.S.R. [C.F. Plant Br. Abstr. (56), 569].
- Habura, E.C. (1958).** Heterosis in yield characters in sunflower. *Zuchter* 28: 285-287. [C.F. Oil Seed and Pulse Crops in Western Canada, John Haraprala 1975].
- Jukic, V. (1990).** Evaluation of the value of sunflower lines and the way of inheriting of some quantitative characteristics by diallel crossing system. *Osijek, Poljoprivredni Institut* 50 p.
- Kadkol, G.P.; I.J. Anand and R.P. Sharma. (1984).** Combining ability and heterosis in sunflower. *Indian J. Genetics and Pl. Breed.*, 44: 444-451.
-

- Kashef, E.A.A. (1996).** Breeding studies on sunflower (*Helianthus annuus* L.). Ph.D. Thesis, Faculty of Agric. Mostohor, Zagazig Univ. Benha Branch).
- Kempthorne, D. (1957).** An Introduction to Genetic Statistics, New York, John Wiley.
- Kesteloot, T.A.; J. Heursel and F.M. Pauwels. (1985).** Estimation of the heritability and genetic variation in sunflower (*Helianthus annuus* L.). *Helia*, 8: 17-20.
- Li, Y.M.; R.L. Chaney; A.A. Schneiter and J.F. Miller. (1995).** Combining ability and heterosis estimates for kernel cadmium level in sunflower. *Crop Sci.*, 35: 1015-1019.
- Madrap, I.A. and V.G. Makne. (1993).** Heterosis in relation to combining ability effect and phenotypic stability in sunflower (*Helianthus annuus* L.). *Indian J. Agric. Sci.*, 63: 484-488.
- Manjunath, A. and J.V. Goud. (1983).** Genetics of quantitative characters in sunflower (*Helianthus annuus* L.). *Genetica Iberica*, 1: 13-23.
- Marinkovic, R. (1993).** Combining ability of some inbred sunflower (*Helianthus annuus* L.) lines. *J. Genetics and Pl. Breed.*, 53: 299-304.
- Marinkovic, R.; D. Skoric and V. Secerov-Fiser. (1987).** A study of the inheritance of some seed characteristics in diallel crosses of inbred lines of sunflower (*Helianthus annuus* L.). *Arhiv za Poljoprivredne Nauke*, 4: 213-221.
- Miller, J.F. and G.N. Fick (1978).** Influence of plant population on performance of sunflower hybrids. *Can. J. Plant Sci.* 58: 597-600.
-

- Miseta, V. (1969).** Oil content per unit area in sunflower as a function of spacing. *Délafoldi Mezogaz dasagi Kisérlati Intézet Kalazemenyei*, Szeged 6: 129-142. [C.F. Field Crop Abst. 25: 878, 1972].
- Mohamed, A.A.E. (1997).** Effect of plant spacing and nitrogen fertilization on yield and yield components of some sunflower cultivars. *J. Agric. Res. Tanta Univ.*; 23(3): 300-308.
- Monotti, M. (1973).** Investigation of the cultural technique for sunflower. 2- Plant density and spacing. *Annali della Facolta di Agraria, Universta degli studi di Periegia* 28: 551-563.
- Moursi, M.A.; Olfat El-Bagaury and T. Behairy (1983).** harvest index in sunflower plant to some cultural treatments. *Proc. First Conf. Agro., 1983, Egypt. Vol. 2.* 535-544.
- Naik, N.M.; B.B. Pawar and A.D. Dumbre. (1987).** Line x tester analysis for combining ability in sunflower. *Current Research Reporter, Mahatma Phule Agric. Univ., 3:* 10-15. (Computer Research, AN 891607376).
- Nasr Allah, A.K.; M.A. Gomma, I.F. Rehabb and I.A. Saleh (1994).** Effect of plant density and weed control on yield and quality of sunflower. *Menofiya J. Agric. Res.*; 6(1): 3125-3140.
- Nuthan, D. (1981).** Study of heterosis and stability index in sunflower (*Helianthus annuus* L.). *Mysor J. of Agric. Sci.* 14(1): 125-126, 198 (C.F. Plant Br. Abst. 51, 5379).
- Ortegon M.A.S. and A.E. Mendoza. (1993).** Combining ability of sunflower hybrids (*Helianthus annuus* L.) for achene yield and oil. *Helia*, 16: 45-54.
- Pathak, A.R.; B. Singh and M.U. Kukadia. (1985).** Combining ability analysis in sunflower (*Helianthus annuus* L.). *Indian J. Heredity*, 17: 12-22.
-

- Pet, Kov, D. (1992).** Effect of the tester in evaluations of general combining ability in inbred lines of sunflower. *Genetika i Seleksiya*, 25: 383-386.
- Putt, E.D. (1966).** Heterosis, combining ability and predicted synthetic from a diallel cross in sunflower (*Helianthus annuus* L.). *Can. J. Plant Sci.* 46: 59-67.
- Putt, E.D. and D.G. Dorell (1977).** Breeding for sunflower production oil seed and pulse crops in Western Canada John T Harapiala, 1977.
- Rao, A.M.; Y.Y. Rao and G.H.S. Reddi (1976).** Investigation on the optimum spacing and time of N application for sunflower. *Indian J. Agric. Res.* 10(2): 79-100.
- Rao, U.R.; B.B. Pawar and A.D. Dumbre. (1992a).** Combining ability studies in sunflower through diallel analysis. *J. Maharashtra Agric. Univ.*, 17: 149-150. (Computer Research, AN 941607127).
- Rao, U.R.; B.B. Pawar and A.D. Dhumre. (1992b).** Heritability and gene action studies in sunflower. *J. Maharashtra Agric. Univ.*, 17: 504-505. (Computer Research, AN 941603221).
- Rashed, R.H. and J.T. Moosa. (1990).** Effect of inbreeding and open pollination on some yield characters of sunflower (*Helianthus annuus* L.). *Mesopotamia J. Agric.*, 22: 253-262.
- Robinson, R.G.; J.H. Ford; W.E. Lueschen; D.L. Robas; L.J. Smith; D.D. Warnes and J.V. Wiersma (1980).** Response of sunflower to plant population. *Agron. J.* 72: 869-871.
- Rozhkova, V.T. and A.V. Anaschenko. (1981).** Specific combining ability of sunflower varieties for achene yield. *Byulleten, vsesoyuznogo Ordena lenina i Ordena Druzhby Narodov Instituta Rasteniievodstva Imeni NI Vavilova* 115; 49-53. (Computer Research, AN 791674720).
-

- Seetharam, A. (1977).** Performance of hybrids of sunflower produced by means of cytoplasmic male sterility. *Sabaras J.* 9(1): 51-55. [C.F. Genetic Abstr. 10(1): 601, 1978].
- Shankara, A.C. (1983).** Evaluation of sunflower inbreeds for their combining ability by line x tester analysis. *Thesis Abstracts*, 9: 284.
- Shanthamallaiah, N.R. Shjivaraj; N.M. Patil and Veeranna (1977).** Effect of plant population and fertilizer levels on growth, yield and yield components in sunflower. *Mysore J. Agric. Sci.* 11(4): 477-480.
- Sheriff, N.M. and R. Appadurai (1987).** Genetic variability in some quantitative traits of sunflower. *Madras Agric. J.* (1987) 72(9): 539-540. (C.F. Plant Br. Abstr. 56. 7287).
- Sheriff, N.M.; R. Appadurai; M. Rangaswamy and N.M. Sheriff. (1985).** Combining ability in sunflower. *Indian J. Agric. Sci.*, 55: 315-318.
- Shrinivasa, K. (1983).** Inheritance of fertility restoration and oil content in sunflower (*Helianthus annuus* L.). *These Abstr.* (1983). 8(1): 7071. [C.F. Plant Br. Abstr. 53. 790].
- Shvestsova, V.P. (1982).** Effect of inbreeding and hybridization on the fatty acid composition of sunflower seed oil. *Ref. Zh.* (60): 38-40. [C.F. Plant Br. Abstr. 52, 5252].
- Sidhu, B.S. and D.S. Bains (1980).** Effect of planting technique on growth and seed yield of sunflower. *J. of Research, Punjab Agric. Univ.* 17(2): 148-151.
- Singh, H.; T.P. Yadava and A.K. Yadava (1981).** Heterosis in intervarietal crosses of sunflower crop improvement 1978 5(1): 84-85 [C.F. Plant Br. Abstr. 51, 2412].

- Skoric, D. (1980).** Production and economic values of new NS sunflower hybrids. Sunflower Newsletter (1977). 1(2): 5-8. [C.F. Plant Br. Abstr. 50, 8715].
- Skoric, D. (1981).** Desired model of sunflower hybrid and newly developed NS-sunflower hybrids. International course production and processing of sunflower. Novi Sad. Yugoslavia 1981.
- Smith, B.C.; A.M. Akhanda; V.E. Green, Jr. and G.M. Prine (1981).** Early planted oilseed sunflower cultivars grown at three rows spacing and plant populations in North Florida. Soil and Crop Sci. Society of Florida, 40: 165-171.
- Srinivas, K. and S.V. Patil (1977).** Effect of spacing and fertility levels on growth and yield of sunflower (*Helianthus annuus* L.). Mysore J. Agric. Sci. 11(1); 41-45. (Computer Research, AN 751921870).
- Stanev, V. (1981).** The photosynthetic activity of sunflower in relation to the conditions of cultivation. International Zeitschrift der Indwirthaft, 21: 152-156.
- Stoyanova, I. (1971).** Inheritance of some sunflower characters in the FI. Genetic Selekt. 4(1): 13-14. [C.F. Plant Br. Abstr. 51, 3470].
- Sturma, M.P. (1970).** Experiments in 1970 on varieties of sunflower. Annali dell Istituto Sperimental per la Colture Industrial 2: 61-74.
- Sudhakar, D. (1981).** heterosis and combining ability in sunflower (*Helianthus annuus* L.). Mysore J. of Agric. Sci. 1979 (13): 476 [C.F. Plant Br. Abstr. 51, 3470].
- Sun, G.Z. (1986).** A study on heterosis in sunflower. Oil Crops of Chile, 1: 30-32.
-

- Swallers, C.M. and G.N. Fick (1973).** Performance of large seeded sunflower at three plant spacings. North Dakota Farm Res. 31(1): 15-16.
- Tersae, M.; D. Vares and P. Vincourt. (1993).** Combining groups in cultivated sunflower populations (*Helianthus annuus* L.) and their relationships with the country of origin. Theoretical and Applied Genetics, 87: 603-608.
- Toit, D.D. (1981).** An evaluation of sunflower hybrids and open-pollinate cultivars in respect of certain important agronomic characteristics Teg. Med. Van 1979, No. 164 pp. 8. [C.F. Plant Br. Abstr. 51, 6443].
- Vagvolgyi, S. (1984).** Estimation of heterotic effect in sunflower. Novenytermeles, 33: 97-104.
- Vanisree, G.; K. Ananthasayana; G.V.S. Nagabhushasnam and C.A. Jagadish. (1988).** Studies on combining ability in sunflower (*Helianthus annuus* L.). Journal of Oilseeds Research, 5: 52-61.
- Vijaylakshmi, K.; N.K. Sanghi; W.L. Pelton and C.H. Anderson (1975).** Effect of plant population and row spacing on sunflower agronomy. Can. J. Plant Sci. 55: 491-499.
- Vranceanu, A.V. (1983).** The performance of sunflower hybrids in production actual and prospective progress. General Siplant Tehnice 1984, 48, 67-74. (C.F. Plant Br. Abstr. 53.8993).
- Vranceanu, A.V. and N. Pirvu (1988).** Genetic advance realized in breeding sunflower in Romania. Analele Institutului de Cercetari Pentru Cereale si Plante Techice. Fundulea. 56, 103-114. (Computer Res., An 891608502).
- Yousaf, M.; B. Akhtar and A. Shakoar (1986).** Effect of spacing and nitrogen on the yield and yield components of sunflower under rainfed conditions. National Agric. Res. Center. Islamabad, Pakistan.
-