

## References

---

## REFERENCES

A.A.C.C. (1962).

American Association of cereal chemists, Approved methods, 8th edition, St. Paul. U.S.A.

Abdel-Khafar, A. (1988).

Bread, Economical view.

Symposium of Arab Bread, Cairo.

Abdel-Salam, A.M. (1982).

Reductones and oxidants of the Maillard reactions: their contribution on the production of aldehyde from alcohol. Ph.D.Thesis.

Universite Catholique de Louvain.

Abu-El Nasr, S. and El-Nahal, A.M. (1968).

Agric. Insects and their applied control methods, Alam El-Kotob. Pub. Cairo.

Anderson, J.W. and Chen, W.J.L. (1979).

Plant fiber. Carbohydrate and Lipid Metabolism.

Am. J. Clin. Nutr. 32:346.

Anderson, Y.; Hedlund, B.; Jonsson, L. and Sik, S. (1981).

Extrusion cooking of high-fiber cereal product with crisbread Character.

Cereal Chem. 58(5): 370-374.

Anon (1969).

Grain processing- Machinery and Equipment for flour mills.

TMS Pardubice CSSR.

Anon (1976).

The practice of flour milling (Vol.1).

Inc. National Association of British and Irish Millers  
limited.

Anon (1978).

Foods of special dietary uses, low caloric bread ,  
Egyptian Standard, 1432.

Egyptian Organization for Standardization.

Anon (1983).

MaCaroni, Egyptian Standard No.49/1983

Egyptian Organization for Standardization.

Anon (1984).

Ministry of supply, Act 515, 1984, for flour extraction  
from wheat.

Anon (1985).

Imported wheat standard quality, General Authority for  
supply commodities, Cairo.

Anon (1987).

Egyptian Standard Project for MaCaroni. Egyptian  
Organization for Standardization.

A.O.A.C. (1980).

Association of official analytical chemists, "Official  
Methods of Analysis".

13<sup>th</sup> ed., Washington.

Arnbjornsson; E.; Asp., N.G., and Westin, S.I. (1982).

Decreasing incidence of acute appendicitis, with  
special reference to the consumption of dietary fiber.  
Acta Chir. Scand, 148:461.

---

- Arunga, R.O. and Morrison, W.R. (1971).  
Structural analysis of wheat flour glycerolipids.  
Lipids 6:768.
- Ashour, H. K. (1978).  
Study of phytic acid hydrolysis in wheat flour,  
chemical and enzymatic during baking.  
M.Sc. Thesis, Al Azhar Univ.
- Ashour, H. K. (1985)  
Biochemical studies on preparation and evaluation  
of some wheat protein concentrations.  
pH.D. Thesis, Ain Shams, Fac. of Agric.
- Bailey, C.H. (1916).  
The relation of certain physical characteristics of  
the wheat kernel to milling quality.  
J. Agric. Sci. 7:432.
- Baldini, M.; Centi; Micco, C. and Stacchini, A. (1983).  
Study on the presence of heavy metals in cereals.  
FSTA 15 No.6 6M 866.
- Berg, S.O. (1947).  
Is the degree of grittiness of wheat flour mainly  
a varietal character?  
Cereal Chem. 24:274.
- Bianchetti, R. and Maria, L.S. (1967).  
The mechanism of repression by inorganic phosphate  
of phytate synthesis in germinating wheat embryo.  
Biochem. Biophys. Acta, 145(2):488-490.
-

Booth, M.R. (1973 ).

Private Communication.

(C.F. Kingswood, K. 1975, The Structure and Biochemistry of the Wheat Grain. Bread Symposium, Applied Science Pub. London).

Bourdet, A. and Feillet P. (1967).

Distribution of phosphorus compounds in the protein fraction of various types of wheat flours.

Cereal Chem. Vol. 44 September,

Brennan, Paul. ( 1982).

Flour Milling Technology. Grain and Oil Seeds.

Handling , Marketing Processing.

Canadian International Grains Institute. Thrid Ed.  
Reveised.

Buckman, H.O.; and Brady, N.C. (1960).

The Nature and Properties of Soils.

6th ed., MacMillan Comp. New York.

Burkwall, M.P. and Glass, R.L. (1965).

The fatty acids of wheat and its milled products.

Cereal Chem. 42, 236-246.

Chen, J.Y.; Bohnsack, K. and Labuza, T.P. (1983).

Kinetics of protein quality loss in enriched pasta stored in a sine wave temperature condition.

J. of Food Science 48, 460-469.

Clements, R.L. (1977).

Quantitative studies of wheat flour lipids extracted with various solvents by an elution method.

Cereal Chem. 54(6):1198-1206.

Crampton, E.W. and Harris, L.E. (1969).

Applied animal nutrition. 2<sup>nd</sup> edition.

W.H. Freeman and Co., San Francisco.

Davidson, S.; Passmore R.; Brock, J.F. and Truswell, A.S. (1975).

Human Nutrition and Dietetics.

6<sup>th</sup> edition. E.L.B.S.

De-Boland, A.R.; Garmer, G.B. and O'Dell, B.L. (1975).

Identification and properties of "phytate" in cereal grains and oilseed products.

J. Agric. Food Chem. 23:1186.

Dexter, J.E.; Dronzek, B.L. and Matsuo, R.R. (1978).

Scanning electron microscopy of cooking spaghetti.

Cereal Chem. 55 (1):23-30

\_\_\_\_\_ and Matsuo, R.R. (1977).

Changes in semolina proteins during spaghetti processing.

Cereal Chem. 54(4):882-894.

\_\_\_\_\_ and \_\_\_\_\_ (1978a).

The effect of gluten protein fractions on pasta dough rheology and spaghetti making quality.

Cereal Chem. 55(1); 44-57.

\_\_\_\_\_ and \_\_\_\_\_ (1978b).

Effect of semolina extraction rate on semolina characteristics and spaghetti quality.

Cereal Chem. 55(6):842-852.

\_\_\_\_\_ and \_\_\_\_\_ (1979a).

Effect of water content on changes in semolina protein during dough mixing.

Cereal Chem. 56(1) : 15-20.

Dexter, J.E. and Matsuo, R.R. (1979b).

Effect of starch on pasta dough rheology and spaghetti.  
Cereal Chem. 56(3):190-195.

\_\_\_\_\_ and \_\_\_\_\_ (1981).

Effect of starchy kernels, immaturity, and shrunken  
kernels on durum wheat quality.  
Cereal Chem. 58 (5) : 395-400.

\_\_\_\_\_; \_\_\_\_\_ and Morgan, B.C. (1983).

Spaghetti stickiness: some factors influencing sticki-  
ness and relationship to other cooking quality chara-  
cteristics.  
J. Food Science, 48, 1545-1551.

Dick, J.W.; Banasik, O.J. and Vasiljevic, S. (1980).

1980 North Dakota durum wheat quality report.

Doekes, G.J. and Wennekes, L.M. (1982).

Effect of nitrogen fertilization on quantity and  
composition of wheat flour protein.  
Cereal Chem. 59(4) 276-278.

Donnelly, B.J. (1982).

Teflon and non Teflon lined dies: Effect on Spaghetti  
quality.  
J. Food Science 47, 1055-1058

Eichner, K. and Karel, M. (1972).

The influence of water content and water activity on  
the sugar-amino browning reaction in model systems  
under various conditions.  
J. Agr. Food. Chem. 20 (2) : 218-223.

El-Gindy, M.M. (1982).

Food science, Vol. 1, cereal technology.

Dar El-Maaref, Pub. (In Arabic).

Farrell, E.P.; Ward, A; Miller, G. and Lovett, L. (1967).

Extensive analysis of flours and millfeeds made from nine different wheat mixes: 1-Amount and Analysis.

Cereal Chem . 44: 39-47.

Felix, G.R.; Boonthong, P., and Ronald, E.W. (1982).

Maillard browning reaction of sugar-glycine model systems; changes in sugar concentration, colour and appearance.

J. Food Science 47, 1376-1377.

Fleming, J.R.; Miller, B.S. and Johnson, J.A. (1961).

A method for the determination of relative amounts of malted-wheat, Fungal (*Aspergillus oryzae*) and Bacterial (*Bacillus subtilis*) Alpha-amylase in mixtures, and its application to malted wheat.

Cereal Chem. 38, 479-486.

Fraser, J.R. and Holmes, D.C. (1959).

Proximate analysis of wheat flour carbohydrates. IV. Analysis of wholemeal flour and some of its fractions.

J. Sci. Food Agric. 10:506.

Frey, A. and Holliger, A. (1972).

Pasta goods under microscope.

Buhler Diagram 53:21.

Frolich, W. and Asp., N. (1981).

Dietary fiber content in cereals in Norway.

Cereal Chem. 58(6):524-527.



Gobky, N. (1972).

Milling Technology.

pub. by the General establishment of Mills and Bakeries.

Goldblith, S.A. and Tannenbaum, S.R. (1966).

The nutritional aspects of the freeze-drying of foods.

In: Knehnau, J. and Cramer, H.D. eds. Proc. Intl. Cong.

Nutr. 7th. Pergman: Oxford.

Golik, V. (1976).

Contents and yield of protein in spring wheat cultivars  
in relation to fertilizer and swing rates.

F.C.A. 31, 2, 979.

Graveland, Aris; Bongers Piet, and Bosveld Pieter (1979).

Extraction and fractionation of wheat flour proteins.

J. Sc. Food Agric. 30, 71-84.

---

\_\_\_\_\_ ; Bosveld, Pieter; Lichtendonk, H.; Mooner, H.  
and Scheepstra, A. (1982).

Extraction and fractionation of wheat flour proteins.

J. Sci. Food Agric. 33, 1117-1128.

Hall, J.L.; Flowers, T.J. and Roberts, R.M. (1984).

Plant cell structure and metabolism. E.L.B.S.

Hall, R.J.; Trimder, N. and Givens, D.I. (1973).

Observations on the use of 2,4,6-trinitrobenzenesul-  
phonic acid for the determination of available lysine  
in animal protein concentrate.

Analyst, 98, 673-686.

Hana, L. Ph.; Abdel-Wahab, A.M. and Yousef, K.E. ( 1970 ).  
The influence of nitrogen fertility level on the  
yield and physico-chemical properties of Egyptian  
wheat Giza 150 and Giza 155 varieties.  
(c.f. El.Bardeny, A., Ms.C Thesis, 1981., El Minia univ.)

Harland, B.F. and Harland, J. (1980).  
Fermentative reduction of phytate in Rye, white,  
and whole wheat breads.  
Cereal Chem. 57(3): 226-229.

Harris, R.H.; Smith, G.S. and Sibbitt, L.D. (1943).  
The effect of sprout damage on the quality of durum  
wheat, semolina, and macaroni.  
Cereal Chem. 20:333-345.

Hinton, J.J. (1959).  
The distribution of ash in the wheat kernel.  
Cereal Chem. 36, 19-31.

Hook, S.C.; Bone, T.G. and Thomas, F. (1982a).  
The conditioning of wheat, the influence of varying  
levels of water addition to U.K. wheats on flour  
extraction rate, moisture and colour.  
J. Sci. Food Agric., 33, 645-654.

\_\_\_\_\_; \_\_\_\_\_; \_\_\_\_\_; (1982b).  
The conditioning of wheat. The effect of increasing  
wheat moisture content on the milling performance of  
U.K. wheats with reference to wheat texture.  
J. Sci. Food Agric., 33, 655-662.

\_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_ ( 1982c)  
The conditioning of wheat. An investigation into the  
conditioning requirements of canadian wester red spring.  
No.1.  
J. Sci. Food Agric., 33, 663-670.

Hook, S.C.; Bone, T.G. and Thomas, F. (1982d).

The conditioning of wheat, the influence of roll temperature in the Buhler laboratory mill on milling parameters .  
J. Sci. Food. Agric., 33, 639-644.

I.F.T. (1979).

Dietary fiber, A scientific status summary by the Institute of Food Technologists' Expert Panel on Food Safety & Nutrition and the Committee on Public Information.  
Food Technol . 33(1):35.

Irvine, G.N. (1971).

Durum wheat and pasta products. Page 777 in: Pomeranz, Y., Wheat chemistry and technology, 2nd ed. Am. Assoc. cereal chem.; St. Paul., MN.

\_\_\_\_\_ and Anderson, J.A. (1951).

Air bubbles in macaroni dough.  
Cereal Chem., 28:240-242.

\_\_\_\_\_ ; Bardely, J.W. and Martin, G.C. (1961).

A Farinograph technique for macaroni dough.  
Cereal Chem., 38, 153-158.

Itoh, T.; Tamura, T. and Matsumoto, T. (1973).

Sterol composition of 19 vegetable oils.  
J. Am. Oil Chem. Soc. 50:122.

Kelsay, J.L. (1981).

Effect of diet fiber level on bowel function and trace mineral balances of human subjects.  
Cereal Chem. 58(1):2-5.

Kent, N.L. (1966 ).

Subaleurone endosperm of high protein content.  
Cereal Chem. 43:585.

---

————— (1983).

Technology of cereals.  
Pergman Press, Third Edition.

Kent-Jones, D.W. and Amos, A.J. (1967).

Modern cereal chemistry, 6<sup>th</sup>, edition,.  
Food Trade Press, London.

Kent, N.L. and Evers, A.D. (1969).

Variation in protein composition within the endosperm  
of hard wheat.  
Cereal Chem., 46, 293.

Kerr, R.W.; Cleveland, F.C. and Katzbeck, W.J. (1951).

The action of amyloglucosidase on amylose and amylo-  
pectin.  
J. Am. Chem. Soc., 73, 3916-21.

Kingswood, K. (1975).

The structure and biochemistry of the wheat grain.  
Bread Symposium, Applied Science Pub. London. 47-66.

Kobrehl, K.; Laignelet, B. and Feillet, P. (1974).

Study of some factors of macaroni brownness.  
Cereal Chem. 15(5): 675-684.

Koch, R.B.; Geddes, W.F. and Smith, F. (1951).

The Carbohydrates of the Gramineae.  
1. The sugars of the flour of wheat (*Triticum vulgare*)  
Cereal Chem. 28:424.

---

Krol, T.; Brozowska, E. and Kalisiewicz, B. (1983).

Use of natural cellulose sources for production of doughs with increased cellulose content.

F.S.T.A. 15. No.6, 6M, 830.

Kulp, K.; Ranum, P.M.; Williams, P.C., and Yamazaki, W.T. (1980)

Natural levels of nutrients in commercially milled wheat flours:

I- Description of samples and proximate analysis, protein, ash, flour colour

Cereal Chem. 57, 1.

Labuza, T.P. (1970).

Properties of water as related to the keeping quality of foods. Proc. Third Int'l. Congress Inst. Food Technol., 618.

(C.F. Stamp and Labuza, 1983, J. Food Science, Vol. 48, 543-544).

Labuza, T.P.; Bohansack, K. and Kim, M.N. (1982).

Kinetics of protein quality change in egg noodles stored under constant and fluctuating temperatures.

Cereal Chem. 59(2):142-148.

Labuza, T.P. and Saltmarch, M. (1982).

Kinetics of browning and protein quality loss in sweet whey products under steady state and non-steady state storage conditions.

J. Food Sci., 47:92.

———; Tannenbaum, S.R. and Karel M., (1970).

Food Technol. 24(5), 35.

(C.F. Eichner and Karel, 1972, J. Agr. Food Chem., Vol. 20.2).

- Lier, J.B. and Lacroix, L.J. (1974).  
Carotenoids of durum wheat; induced high pigment levels obtained by treatment of the growing plant with chlorophenylthylamine (CPTA) hydrochloride. Cereal Chem. 51 (2) 189-194.
- Lolas, G.M. and Markakis, P. (1977).  
The phytase of Navy beans (*Phaseolus vulgaris*). J. Food Sci., 42:1094.
- , Palamidis, N. and Markakis, P. (1976).  
The phytic acid total phosphorus relationship in Barley, oats, soybeans, and wheat. Cereal Chem. 53:687.
- Lopez, J.; Gordon, D.T. and Fields, M.L. (1983).  
Release of phosphorus from phytate by natural lactic acid fermentation. J. Food Sci.,
- Lorenz, K., and Lowewe, R. (1977).  
Mineral composition of U.S. and Canadian wheat and wheat blends. J. Agric. Fd. Chem. 25:806.
- , ———, Weadon, D. and Wolf, (1980).  
Natural levels of nutrients in commercially milled wheat flours: III mineral analysis. Cereal Chem. 57, 1.
- Maga, J.A. (1982).  
Phytate: its chemistry, occurrence, food interactions nutritional significance, and methods of analysis. J. Agric. Fod Chem. 30(1) reviews 1-9.
-

Mecham, D.K. (1971).

Lipids in pommeranz, Y. (ed), Wheat chemistry and Technology, p.393.

Am. Assoc. Cereal Chem. St. Paul

MedCalf, D.G. and Gilles, K.A. (1965).

Wheat starches. I- Comparison of physicochemical properties.

Cereal Chem. Vol. 42, 558-568.

McDermott, E.E. and Pace, J. (1960).

Comparison of the amino acid composition of the proteins in flour and endosperm from different types of wheat.

J. Sci. Fd. Agric. 11:109.

Menger, A. (1961).

Possible relations between the soluble carbohydrates of durum wheat and the non-enzymic browning (of wheat pastes).

Ber. Getreidechemiker-Tagung, Detmold, 101-114.

(C.F. Matsuo and Irvin, 1967, Cereal Chem. Jan Macaroni Brownness).

Menger, A.; Cleve, H. and Ander, E. (1969).

Versuche zur verbesserung der Teigwaren qualitat durch wa rmebehandlung der Getreiderohstoffe. Brot Gebaeck 9.179,

(C.F. Kobrehel et al., 1974, Cereal Chem. Vol.51,675).

Mineev, .G.; Tishchenkv, A.T.; and Semikhova, D. (1977).

Protein content and technological qualities of grains of wheat grown in rotation given fertilizers regularly.

F.C.A. Vol. 30, No. 5, 2529.

- Morad, M.M.; Magoli, S.B. and Afifi, A.S. (1980).  
Macaroni supplemented with lupins and deffated soy-bean flours.  
J. Food Sci., 45(2), 404-405.
- Morrison, W.R. (1978).  
Wheat lipid composition.  
Cereal Chem. 55(5) 548-558.
- 
- ; Mann, D.L.; Wong, S. and Coventry, A.M. (1975).  
Selective extraction and quantitative analysis of non-starch and starch lipids from wheat flour.  
J. Sci. Food Agric. 26:507.
- Nayini, N.R. and Markakis, P. (1983).  
A research note: Effect of fermentation time on the inositol phosphates of bread.  
J. Food Sci. 48, 262-263.
- Nielsen, H.C.; Babcock; G.E. and Senti F.R. (1962).  
Molecular weight studies on glutenin before and after disulphide bond splitting.  
Arch. Biochem. Biophys. 96:252.
- O'Dell, B.L.; De Boland, A.R., and Koirtyohann, S.R. (1972).  
Distribution of phytate and nutritional important among the morphological components of cereal grains.  
J. Agr. Food Chem., 20 (3), 772.
- Osborne, T.B. (1907).  
The proteins of the wheat kernel. Camegie Inst. Wash.,  
Pub., No. 84.
-



Percival, J. (1921).

The wheat plant.

Duckworth , London, Reprinted (1975).

Phelepophetch, U ., (1980).

Essential of Biochemistry, Vol. 1, Mere,USSR, Pub .  
(IN Arabic).

Polizzoto, L.M.; Tinsley, A.M.; Weber,C.W., and Berry,J.W.  
(1983).

Dietary fibers in Muffins.

J. Food Sci. 48, 111-118.

Pomeranz, Y., (1978).

Wheat chemistry and technology, pp.(791-795),Vol.III,  
Pub., American Association of Cereal Chemists, Inc.  
St. Paul, Minnesota, U.S.A.

Pomeranz,Y., and Shellenberger, J.A. (1961).

Histochemical characterization of wheat and wheat  
products.

IV- Mapping the free fatty acids in Germinating wheat.  
Cereal Chem. 38 (2) 122-131.

Quisenberg, K.S. and Reitz,L.P. Editors (1967).

Wheat and wheat improvement monograph. No. 13,Agronomy  
Series, Am. Zoc. Agronomy. Inc. Madison, Wis.

Ranhotra, G .S. (1972).

Hydrolysis during breadmaking of phytic acid in wheat  
protein concentrate.

J. Food Sci. 37, 12-13.

---

; Loewe, R.J. and Puyat, L.V. (1973).

Effect of dietary phytic acid on the availability of  
iron and phosphorus.

Cereal Chem. 51(3) 323-329.

---

Rasmusson, D.C.; Hester, A.J. Fick, G.N. and Byrne, I. (1971).  
Breeding of mineral content in wheat and barley.  
Crop Sci. 11:623.

Refai, F.Y. (1965).

Essential of milling industry. Pub. by the General  
Establishment of Mills and Bakeries. (In Arabic).

\_\_\_\_\_(1982).

Pasta Production and Technology.  
Pub. by American Wheat Board.

Sandstead, H.H.; Munoz, J.M.; Jacob, R.A.; Klevay, L.M.; Reck,  
S.J.; Logan, G.M.; Dintzis, F.R.; Inglett, G.E.; and  
Shuey, W.C. (1978).

Influence of dietary fiber on trace element balance.  
Am. J. Clin. Nutr. 31:S 180.

Schrenk, W.G. (1955).

Minerals in wheat grain. Kansas Agric. Exp. Sta., Tech.  
Bull. 136.

Schweizer, T.F.; Frolich, W.; Vedovo, S.; and Besson, L. (1984).  
Minerals and phytate in the analysis of dietary fiber  
from cereal.

Cereal Chem. 61(2):116-119.

Seckinger, H.L. and Wolf, M.J. (1967).

Lipid distribution in the protein matrix of wheat  
endosperm as observed by electron -microscopy.  
Cereal Chem. 44, November.

Shellenberger, J.A. (1961).

World wide review of milling evaluation of wheats.  
Ass. Oper. Millers Bull. 2620.  
(C.F. Kent, 1983, Technology of Cereals. Pergman  
Press, Third Edition).

Simmonds, D.H. (1978).

Structure, composition and biochemistry of cereal  
grains. In cereals' 78: Better Nutrition for the  
World's Millions, p.105 Pomeranz.4. (Sci.Ed.) AACC,  
Paul, Minn., U.S.A. on Behalf of the sixth inter-  
national cereal and bread congress..

---

\_\_\_\_\_ ; Barlow, K.K. and Wrigley C.W. (1973).

The biochemical basis of grain hardness in wheat.  
Cereal Chem., 50(5), 553-562.

Sosulski, F.W. and Cadden, A.M. (1982).

Composition and physiological properties of several  
sources of dietary fiber.  
J. Food Sci. Vol. 47:1472-1477.

Southgate, D.A.; Branch, W.J.; Drasar, B.S., Walters, R.L.;  
Davies, P.S. and Mclean, B.I. (1976).

Metabolic responses to dietary supplements of bran.  
Metabolism 25:1129.

Spiller, G .A.; Shipley, E.A. and Blake, J.A. (1978).

Recent progress in dietary fiber (plantix) in human  
nutrition.  
Crit. Rev. Food Sci. Nutr. 10:31.

Staudt, E. and Ziegler, E. (1973).

Flour chemistry.  
Buhler Brothers L. Switzerland.

Stenvert, N.L. and Kingswood, K. (1977).

The influence of physical structure of the protein matrix on wheat hardness.

J. Sci. Fd. Agric. 14:284.

Stevens, D.J.; McDermott, E.E. and Pace, J. (1963).

Isolation of endosperm protein and aleurone cell contents from wheat and determination of their amino acid composition.

J. Sci. Fd. Agric. 28:11.

Tabekhia, M.M., and El Esly, N. (1982).

Effect of various methods of common Egyptian Bread-making ( native bread) on hydrolysis of phytic acid.

F.S.T.A. Vol. 14, No.2, 2M. 302.

Tangkongchiter, U.; Seib, P.A., and Hoseney, R. (1981).

Phytic acid: I. Determination of three forms of phosphorus in flour, dough and bread.

Cereal Chem. 58(3) 226-228.

Tawfik, M. and Mansour S. (1983).

Cooking quality of Macaroni produced from different flours in comparison with durum flour and semolina.

Annals Agric., Sci., Fac., Agric. Ain. Chams, 28(3), 1525-1537.

Thompson, J.B. ( 1950).

A browning reaction involving copper proteins, copper metabolism, a symposium on animal, plant and soil relationship.

Johns Hopkins Press: Baltimore, Md.

Vakar, A.B. (1961).

Wheat Gluten.

Izdatelstvo Akademii Nauk SSSR, Moscow.C.

(C.F. Cereal Technology, Kent, 1983).

Walsh, D.E.; and Gilles, K.A. (1974).

Wheat production and utilization.

Macaroni products (ch.12). TVI Pub.

\_\_\_\_\_; Younce, V.L., and Gillas, K.A. (1970).

Inhibition of durum wheat lipoxidase with L-Ascorbic acid.

Cereal Chem. Vol. 47. No. 2. 119-125.

Waggle, D.H.; Lambert, M.A.; Miller, G.D.; Farrell, E.P., and Deyoe, C.W. (1967).

Extensive analysis of flours and millfeeds made from nine different wheat mixes.

I- Amounts and Analysis.

II- Amino acids, Minerals, Vitamins and Gross Energy.

Cereal Chem. Vol. 44, Jan., 39-60.

Wang, W.L.; Swain, E.W., and Hesseltine, C.W. (1980).

Phytate of molds used in oriental food fermentation.

J. Food Sci. 45: 1262-1266.

Watson, C.A. and Johnson, J.A. (1965).

Studies on the gelatinization of starch.

I- Competition for water by protein and cellulose derivatives.

J. Food Sci. 30, 450.

Wenlock, R.W.; Sivell, L.M.; King, R.T.; Scuffam, D.; and

Wig gins, R. (1983).

The nutritional composition of British bread a nationwide study.

J. Sci. Food Agric. Vol. 34, 1302-1218.

- Woychik, J.H.; Boundy, J.A. and Dimber, R.J. (1961).  
Starch gelelectrophoresis of wheat gluten protein  
with concentrated urea.  
Arch Biochem. Biophys. 94:477.
- Wozenski, J. and Woodurn, M. (1975).  
Cereal Chem. 52, 665.  
(C.F. Maga, J.A., 1982, J. Agric. Food Chem. Vol. 30,  
No. 1, Reviews).
- Wyland, A.R., and D'Appolonia, B.L. (1982).  
Influence of drying temperature and farina blending  
on spaghetti quality.  
Cereal Chem. Vol. 59, No. 3, 199-201.
- Yadkin, M. and Offord, R. (1980).  
Comprehensible Biochemistry.  
E.L.B.S.
- Yasunaga, T.; Bushuk, W. and Irvine, G.N. (1968).  
Gelatinization of starch during baking.  
Cereal Chem. 45, 269.
-