

LITERATURE CITED

- Anand, I.J. and R.R. Downey. 1981.
A study of Erucic acid alleles indigenomic rapeseed
(*B. napus* L.).
Can. J. Plant Sci. 61 (2): 199-203.
- A.O.C.S. 1975.
Official and tentative methods of analysis 2nd ed.
Chicago, Illinois, U.S.A.
- Badwal, S.S.; Gupta and K.S. Labana. 1986.
Combining ability as influenced by environmental
variation in Indian mustard.
Crop Improvement, 13 (1): 86-91.
(c.f. Plant Breeding Abstracts, 57; 3, Abst., 2213
1987).
- Beversdorf, W.D.; L.R. Erickson and I. Grant. 1985.
Hybridization process utilizing a combination of
cytoplasmic male sterility and herbicide tolerance.
United States Patent and Trademark Office Serial No.
4517763.
(c.f. Can. J. Genet. Cytol. 27: 472-478, 1985).
- Brandle, J.E. and P.B.E. McVetty. 1989.
Heterosis and combining ability in hybrids derived from
oil seed rape cultivars and inbred lines.
Crop Sci. 29: 1191-1195.
- Buson, M. 1980.
Heterosis and genetic parameters in winter rape (*B.
napus* L.).
Cruciferae Newsletter, 5: 13-14.
(c.f. Plant Breeding Abstracts, 52 (1): Abst., 651, 1982).

Calhoun, W.; J.M. Crane and D.L. Stamp. 1975.

Development of low glucosinolate, high Erucic acid rapeseed breeding program.

J. of the American Oil Chemists' Society, 52 (9): 363-365.

Chauhan, Y.S. and D. Singh. 1980.

Inheritance of seed weight in Indian mustard.

Indian Journal of Genetics and Plant Breeding, 40 (3): 597-599.

(c.f. Plant Breeding Abstracts, 52 (4): 3235, 1982).

Degenhardt, D.F. and Z.P. Kondra. 1981.

The influence of seeding date and seeding rate on seed yield and yield components of five genotypes of (*B. napus* L.).

Can. J. Plant Sci. 61 (2): 175-183.

Diepenbrock, W. and R.F. Wilson. 1987.

Genetic regulation of Linolenic acid concentration in rapeseed.

Crop Sci. 27: 75-77.

Dorrell, D.G. and R.K. Downey. 1964.

The inheritance of Erucic acid content in rape seed (*B. campestris*).

Can. J. Plant Sci. 44 (6): 499-504.

Downey, R.K. and B.L. Harvey. 1963.

Methods of breeding for oil quality in rape.

Can. J. Plant Sci. 43 (4): 271-275.

Downey, R.K.; B.M. Craig and C.G. Youngs. 1967.

Breeding rapeseed for oil and meal quality.

Symposium: Modification of oil seeds through plant breeding.

Conducted by The American Oil Chemists' Society at its 41st Fall Meeting, Chicago, Illinois, October 15-18.

Gilbert, N.E.G. 1958.

Diallel cross in plant breeding.

Heredity, 12: 477-492.

Govil, S.K.; C.N. Chaubey and Y.S. Chauhan. 1981.

Combining ability studies in Indian mustard.

Indian J. Agric. Sci., 51 (9): 623-626.

(c.f. Plant Breeding Abstracts, 52 (7): Abst., 5951, 1982).

Grabiec, B. 1981.

Evaluation of winter swede rape hybrids with low Erucic acid content in the oil and various contents of thioglycosides in seed meal.

Biuletyn Instytutu Hodowli: Aklimatyzacji Roslin, 146: 71-86.

(c.f. Plant Breeding Abstracts, 53 (8): 6646, 1983).

Grami, B. and B.R. Stefansson. 1977.

Gene action for protein and oil content in summer rape.

Can. J. Plant Sci. 57 (4): 625-631.

----- and ----- . 1977 a.

Paternal and maternal effects on protein and oil content in summer rape.

Can. J. Plant Sci. 57 (4): 945-949.

----- ; R.J. Baker and B.R. Stefansson. 1977.

Genetics of protein and oil content in summer rape:

Heritability, number of effective factors, and correlations.

Can. J. Plant Sci. 57 (4): 937-943.

Grant, I. and W.D. Beversdorf. 1985.

Heterosis and combining ability estimates in spring
planted oilseed rape (*B. napus* L.).

Canadian J. of Genetics and Cytology, 27 (4): 472-478.

Griffing, B. 1956.

Concept of general and specific combining ability in
relation to diallel crossing systems.

Australian J. Biol. Sci. 9: 463-493.

Gupta, S.K.; S.K. Thakral; T.P. Yadava and P. Kumar. 1985.

Combining ability and genetic architecture of oil
content in Indian mustard.

Haryana Agricultural University Journal of Research, 15
(4): 467-470.

(c.f. Plant Breeding Abstracts, 56 (10): Abst. 8970,
1986).

Harvey, B.L. and R.K. Downey. 1964.

The inheritance of Erucic acid content in rapeseed
(*B. napus*).

Can. J. Plant Sci. 44 (1): 104-111.

Hutcheson, D.S.; R.K. Downey and S.J. Campbell. 1981.

Performance of naturally occurring subspecies hybrid
in *B. campestris* L. var *oliefera* Metzg.

Can. J. Plant Sci. 61: 895-900.

Ibrahim, H.K.R. 1989.

Studies on some diallel crosses in rapeseed.

M. Sc. Thesis, Fac. Agric. Alex. Univ.

Iwona Bartkowiak-Broda. 1978.

Inheritance of fat content and fatty acid composition
in seeds of zero-Erucic winter rape (*B. napus*).

5th International Rapeseed Conference, Valmo, Sweden,
June; 12-16.

Jindal, S.K. and K.S. Labana. 1986.

Combining ability in a complete diallel cross of
Indian mustared.

Indian J. Agric. Sci., 56 (2): 75-79.

(c.f. Plant Breeding Abstracts, 56 (7): 6015, 1986).

Jonsson, R. 1977.

Erucic-acid heridity in rapeseed (*B. napus* L. and
B. campestris L.).

Hereditas, 86: 159-170.

Kondra, Z.P. and B.R. Stefansson. 1970.

Inheritance of the major glucosinol of rapeseed (*B.*
napus) meal.

Can. J. Plant Sci., 50: 643-647.

Krzymanski, J. and R.K. Downey. 1969.

Inheritance of fatty acid composition in winter forms
of rapeseed (*B. napus*).

Can. J. Plant Sci., 49 (3): 313-319.

Kundu, S. and S.R. Khurana. 1988.

Stability for seed yield and its components in toria
(*B. campestris*).

Indian Journal of Genetics and Plant Breeding, 48 (3):
389-391.

Lee, J.II.; K. Takayanagi and T. Shiga. 1974.

Breeding for improvement of fatty acid composition in rapeseed, *B. napus* L. I. Fatty acid composition in rapeseed oil of Asian and European varieties.
Bull. Nat. Inst. Agr. Sci., 25: 1-16.

Lee, J.II.; M. Saito; T. Shiga; K. Takayanagi and S.

Sugiyama. 1974 a.

Breeding for improvement of fatty acid composition in rapeseed, *B. napus* L. II. Introduction of zero-Erucic acid genes to Japanese varieties.

Bull. Nat. Inst. Agr. Sci., 25: 17-30.

Lefort-Buson, M.; B. Guillot-Lemoine and Y. Dattee. 1987.

Heterosis and genetic distance in rapeseed (*B. napus* L.): crosses between European and Asiatic selfed lines.
Genome, 29: 413-418.

Morice, J. 1978.

The selection of rape for improvement of yield.

Proc. 5th Int. Rapeseed Conf., Malmo. Sweden.

pp. 36-47.

(c.f. Can. J. Genet. Cytol., 27: 47; 478, 1985).

Olivieri, A.M. and P. De Caneva. 1984.

Relationship between filling phas yield per plant in rapeseed.

Cruciferae Newsletter No. 9, 46-47.

(c.f. Plant Breeding Abstracts, 55 (5): 3796, 1985).

Pal, R.; H. Singh and D.S. Jatasra. 1981.

Genetics of yield and yield components in Indian rapeseed.

Indian J. Agric. Sci., 51 (8): 550-553.

(c.f. Plant Breeding Abstracts, 52 (5): 4185, 1982).

Renard, M. and J. Mesquida. 1983.

Pollinisation entomophile due colza male sterile en production de semences hybrides F₁ dans differentes regions de France.

Proc. 6th Int. Rapeseed Conf. Paris, France; pp. 552 - 557.

(c.f. Can. J. Genet Cytol., 27; 472-478, 1985).

Selim, A.K.A.; M.H. Serry; A.A. Tayel; M.I. Nasr and S.A.N. Afiah. 1981.

Genetic behaviour of seed oil content in a cross between two rape varieties under Egyptian conditions. Annals Agric. Sci., 26 (1-2): 143-157.

Sernyk, J.L. and B.R. Stefansson. 1983.

Heterosis in summer rape (*B. napus* L.)

Can. J. Plant Sci., 63: 407-413.

Shelkoudenko, V.G. 1974.

Combining ability in varieties of winter rape.

Byul-nauch-tekhn-inform. pomasic Kul'turam, 1; 8-11.

(c.f. Plant Breeding Abstracts 45 (12): 10256, 1975).

Shiga, T. 1976.

Studies on heterosis breeding using cytoplasmic male sterility in rapeseed *B. napus* L. (Japanese with English summary).

Bull. Nat. Inst. Agric. Sci. Series D. No. 27: 101 pp.

(c.f. Can. J. Plant Sci., 63; 407, 413, 1983).

Singh, S.P. 1973.

Heterosis and combining ability estimates in Indian mustard, *B. juncea* (L.) Czern, and Coss.

Grop Sci., 13 (5): 497-499.

Singh, H. and C.K. Yadav. 1980.

Gene action and combining ability for seed yield,
flowering, and maturity in rapeseed.

Indian J. Agric. Sci., 50 (9): 655-658.

(c.f. Plant Breeding Abstracts, 52 (7): 5952, 1982).

Singh, A.B.; Y.S. Chauhan and P. Singh. 1981.

Genetics of yield in Indian mustard.

Indian Journal of Genetics and Plant Breeding, 41 (1):
130-136.

(c.f. Plant Breeding Abstracts, 52 (5): 4159, 1982).

Stefansson, B.R. and F.W. Hougen. 1964.

Selection of rape plants (*B. napus*) with seed oil
practically free from Erucic acid.

Can. J. Plant Sci., 44 (4): 359-364.

Thompson, K.F. 1972.

Cytoplasmic male sterility in oil seed rape.

Heredity, 29: 253-257.

Vogel, A.R. 1975.

A textbook of practical organic chemistry 3rd ed.

English Language. Book Society and Longman Group Ltd.

Wang, Z.M. and Y.F. Wang. 1986.

A preliminary study of combining ability and
heritability for the main agronomic characters of *B.*
juncea.

Oil Crops of China, No. 1: 11-15.

(c.f. Plant Breeding Abstracts, 57 (2): 1372, 1987).