

REFERENCES

A.A.C.C. (1982).

American Association of Cereal Chemists. Approved methods of A.A.C.C. The Association, 10 th ed. st., paul, Minn., U.S.A.

A.A.C.C. (1983).

American Association of Cereal Chemists. Approved methods 8 th ed. st., paul, Minn., U.S.A. 24-28.

Abdel-Latief, B.M. (1986).

Studies on the fermentation in some types of Egyptian bread and its effect on the bread properties. M.Sc. Thesis, Submitted to Food Science Dept., Faculty of Agric., Moshtohor, Zagazig Univ., Banha Branch.

Alxford, D.W.E.; Colwell, K.H.; Confor, S.J. and Elton, G.A.H. (1968).

Effect of loaf volume on the rate and extent of staling in bread. J. Sci. Food Agric. 19: 95.

A.O.A.C. (1984).

Official Methods of Analysis of the Association of Official Analytical chemists, 14th ed., pub. by A.O.A.C., Vaiginia, DG.

Bacigalupo, A.; Aguilar, T.S.; Lunade La Fuent, R. and Valle, R.J. (1967).

Bread enrichment with cottonseed flour. Cereal Sci. Today. 12: 431.

Bau, H.M.; Mohtadi-Nia, D.; Mejean, L.; and Detry, G. (1983).

Preparation of colorless sunflower protein products:
Effect of processing on physiochemical and nutritional
properties. J.A.O.C.S., 60, 1141-1148.

Bennion, E.B. and Bamford, G.S. (1973).

Cake making processes P.P. 245 In: The Technology of
cake making 5th Leonard Hill Book Print in Great Britain
By Billing and Sons Ltd Guildford and London.

Brule, D. and Savoie, L. (1988).

In vitro digestibility of protein and amino acids in protein
mixtures. J. Sci. Food Agric. 43: 301.

Burns, E.E.; Talley, L.J. and Brummett, B.J. (1972).

Sunflower utilization in human foods. Cereal Sci. Today
17(9): 287-291.

Cancalon, P. (1971).

Chemical composition of sunflower seed hulls.
J.A.O.C.S., 48, 629-632.

**Canella, M.; Bernardi, A.; Castriotta, G.; and Russomanno, G.
(1984).**

Functional properties of fermented sunflower meal. Le
bensmittel-Wissenschaft und-technologie (1984) 17(3):
146-150.

Clandinin, D.R. (1958).

In sunflower seed oil meal in processed plant protein.
Food stuffs, Altschul, A.M. (ed). Academic press, New
York. 557p.

Claughton, S.M.; and Pearce, R.J. (1980).

Protein enrichment of sugar-snap cookies with sunflower
protein isolate. J. of food Sci., 54(3): 354-356.

Dendy, D.A.V.; Clark, P.A.; and James, A.W. (1970).

The use of blends of wheat and non-wheat flour in bread making. *Tropical Sci.*, 12: 131.

Dreese, P.C.; Faibion, J.M. and Hoseney, R.C. (1988).

Dynamic rheological properties of flour, gluten, and gluten-starch doughs. 1. Temperature dependent changes during heating. *Cereal chem.* 65: 348.

Dubois, M.; Gilles, K.A.; Hamilton, J.K.; Rebers, P.A. and Smith, F. (1956).

Calorimetric method for determination of sugars and related substances. *Anal. Chem.* 28: 350.

Earle, F.R.; Van Etten, C.H.; Clark, T.F. and Wolff, I.A. (1968).

Compositional data on sunflower seed. *J.A.O.C.S.*: 45, 876-879.

El-Talawy, F.; Hassan, Sania, A.; Basyony, A.E. and Sadik, Mona (1988).

Functional breadmaking properties of some oil-seed flours. *Annals of Agric. Sci. Moshtohor*, Vol. 26(1): 53-60.

Farag, R.S. (1966).

Biochemical studies on some of the oily seeds grown in U.A.R. M.Sc. Thesis, Dep. of Biochemistry, Faculty of Agric. Cairo, Univ.

Fraizer, and Holmes (1959).

Proximate analysis of wheat flour carbohydrate, analysis of whole meal flour and some of its fractions. *J.Sci. Food Agric.* 10: 506.

Gharib, H.E.E. (1995).

Biochemical studies on some cereals. Ph.D. In Agric., Biochemistry, Faculty of Agric., Cairo University.

Gheyasuddin, S.; Cater, C.M. and Mattile, K.F. (1970).

Effect of several variables on the extractability of sunflower seed protein. *J. Food Sci.* 35: 453.

Ghiasi, K.; Hoseney, R.C. and Zeleznak, K. and Rogers, D.E. (1984).

Effect of Waxy a barely starch and reheating of firmness of bread crumb. *Cereal Chem.* 61: 281.

Hafez, S.A. (1996).

Effect of adding defatted soyflour on the chemical composition and physical properties of bread. *Egypt, J. Agric. Res.* 74(2), 409-418.

Hamed, A.F. and Gordon, L.R. (1982).

Experimental baking method for Egyptian balady bread production and effect of oven baking on bread starch gelatinization and shelf life. *First Egyptian Conf. of Bread Res.* 9-11 Nov., Giza, A.R.E. 106-128.

Huffman, L.M.; Brummett, B.J and Burns, E.E. (1973).

Sunflower as food (in: legume for international food Education News letter, March 1973, ed. S.M. Weisberg, Washington.

Ibrahim, H.M.; Shams El-Din, M.H.A and Atia, A.M. (1992).

Comparative studies on safflower and sunflower meals. *Egypt, J. Food Sci.* 20(2): 273-284.

Ibrahim, N.A.; Fahmy, A.H. and Afaf, A. Attia (1990).

Production of new reduced caloric bread. 1-Effect of Alpha cellulose on dough Rheology and bread properties. *International Conference on Foods for Special Dietary and Medicinal Uses.* March (19-22) Cairo.

Ibrahim, O.S. (1988).

Effect of some additives on wheat flour dough and bread characteristics. M.Sc. Thesis, Fac. of Agric. Cairo Univ.

Jimge, R.C. and Hoseney, R.C. (1981).

A mechanism by which shortening and certain surfactants improve loaf volume in bread. Cereal chem. 58(5): 408-412.

Kazakov, E.D. (1973).

Cultivation, Chemistry and technology of grains and seeds. Pub. House Kolas Moscow.

Kerolles, S.Y. (1986).

Studies on Certain dietary foods. Ph.D. F. Sci. Agric. Ain Shams Univ.

Khairy, M.E.A. and Tayel, S.A. (1988).

Physical and mechanical properties of sunflower seeds. Misr J. Agric. Eng., 5, 251-259.

Khan, M.N. and Lawhon, J.J. (1980).

Baking properties of oilseed protein isolated and produced with industrial membrane system. Cereal Chem. 57 (6): 433-436.

Khan, M.N.; Wan, P.J.; Rooney, L.W. and Lusas, W. (1980).

Sunflower flour: A potential bread ingredient. Cereal Food World. 25(7): 402-404.

Khan, N.A.; Zamon, R. and Elahi, M. (1986).

Effect of processing on the phytic acid content of wheat products. J. Agric. Food Chem. 34: 1010/

Kilara, A. and Humbert, E.S. (1972).

Nitrogen extractability and moisture absorption characteristics of sunflower seed protein. *J. Food Sci.*, 37-771.

Kim, S.K. and D'Appolonia, B.L. (1977a).

Bread staling studies. I. Effect of protein content on staling rate and bread crumb poring properties. *Cereal Chem.* 54: 207.

Kim, S.K. and D'Appolonia, B.L. (1977b).

Bread staling studies. II. Effect of protein content and storage temperature on the role of starch. *Cereal Chem.* 54: 216.

Kim, S.K. and D'Appolonia, B.L. (1977c).

The role of wheat flour constituents in bread staling. *Bakers Dig.* 51(1): 38.

Kitterman, J.S. and Rubenthaler, G.L. (1971).

Assessing the quality of early generation wheat. Selection with the micro A.W.R.C. test. *Cereal Sci. Today* (16): 313:322.

Knowles, P.F. (1978).

In "Sunflower Science and Technology" (J.F. Catter, ed.), PP. 55-87. Am. Soc. Agron., Madison, Wisconsin.

Lathia, D.N. and Koch, M. (1989).

Comparative study of phytic acid content, in-vitro protein digestibility and amino acid composition of different types of flat breads. *J. of the Sci. of Food and Agric.*, 47(3): 353-364.

Lorenz, K. (1978).

"Non-oilseed" sunflower products in bakery foods. Baker.
Dig. 52(4): 28.

Lusas, E.W. (1985).

Sunflower seed protein in : New protein foods.
"Academic press publ, INC., New York., Vol. 5, 393.

Maleki, M. and Seibel, W. (1972a).

Über das altbackenwerden von Bort. III- Der Einfluss
Fett, Emulgatoren und schuttwassermenge Getreide Mehl
Bort 26: 258.

Maleki, M. and Seibel, W. (1972b).

Über das altbackenwerden Von Bort. IV. Einfluss der
weizensorte aufdas altbackenwerden Getreide Mehl Bort
26(7): 190.

Maleki, M.; Hosney, R.C. and Mattern, P.J. (1980).

Effect of loaf volume moisture content and protein quality
on the softness and staling rate of bread. Cereal chemists.
57(2): 138-140.

**Mansour, S.M.; Mervet, A. Osman; Hassan, H.F. and Khorshid,
A.M. (1989).**

Sunflower flour; Apotential bread ingredient. Agric.
Research Review, 67(4); 641-649.

Martin, M.L.; Zeleznak, K.J. and Hosney, R.C. (1991).

Amechanism of bread firming. 1- Role of starch swelling.
Cereal Chem. 68(5): 498.

Matthews, R.H.; Sharpe, E.J. and Clark, W.L. (1970).

The use of some oil seed flours in bread. Cereal Chem.,
47(2): 181-185.

Manistry of Agricultural (1984).

"Sunflower seed" Agric. Comunication and Education,
No. 63.

Mohye, E.M. (1985).

Technological studies on tritcale. M.Sc. Thesis, Fac.
Agric., Al-Azhar Univ.

Morad, M.M. and D'Appolonia, B.L. (1980).

Effect of surfactants and baking procedure on total water-
soluble and soluble starch in bread crumb. Cereal Chem.
57(2): 141-144.

Morgenstern, G. (1994).

Freshness retention in rye bread varieties-methods and
limitations. Getreide, Mehl und Bort 1994, 48(1): 30, 35-
40.

Morrison, W.H.; Akin, D.E. and Robertson, J.A. (1981).

Open pollinated and hybrid sunflower seed structures that
may affect processing for oil. J.A.O.C.S., 58-969.

Mostafa, M.K.; Said, A.S. and Foda, X.H. (1980).

Enrichment of wheat flour with dry skim-milk and its
effect on the baking quality. Egypt J. Food. Sci., 8: 33-39.

Othman, M.A. and Mostafa, M.K. (1986).

Changes in fermentation times and their relation to bread
characteristics. First conf. Food Sci. and Technol. for
Mediterranean Sea Countries, March, 1986, Cairo, Egypt.

Platt, W. and Powers, S.R. (1940).

Compressibility of bread crumb. Cereal Chem., 17: 601.

Pomeranz, Y. (1971).

Glycoliped protein interaction in bread Bakers Dig., 45:
26.

Ponte, J.C.; Titcomb, S.T. and Cotton, R.H. (1962).

Flour as a factor in bread fermenting. *Cereal Chem.*, 38: 449.

Prentic, N.; Guendet, L.S. and Geddes, W.E. (1954).

Studies on bread staling V. Effect of flour fraction and various starches on the fermenting of bread crumb. *Cereal Chem.*, 31: 988.

Rakowska, M.; Racznska-Bdjanowska, K.; Macewicz, I.; Kubiec, R. and Duma, Z. (1989).

Non digestible soluble compounds and grain structure of rye and wheat. *Science* 9: 159.

Ratkowski, A. (1971).

The feed value of rapeseed meal. *J.A.O.C.S.*, 48, 865-868.

Robertson, J.A.; Thomas, J.K. and Donald, B. (1971).

Chemical composition of the seed of sunflower hybrids and open pollinated varieties. *J. Food Sci.*, 36, 873.

Robinson, R.G. (1975).

Amino acid and elemental composition of sunflower and pumpkinseeds. *Agron. J.*, 67, 541-544.

Rogers, D.E.; Zeleznak, K.J.; Lai, C.S. and Hoseney, R.C. (1988).

Effect of native lipids shorting and bread moisture on bread firming. *Cereal Chem.* 65: 398.

Rooney, L.W.; Gustafson, C.B.; Clark, S.P. and Cater, C.M. (1972).

Composition of Baking properties of several oilseed flours. *J. Food Sci.*, 37, 14-18.

Roozen, J.P.; Franke, E.N. and Kinsella, J.E. (1994).

Enzymic and autoxidation of lipids in low fat foods: Model of linoleic acid in emulsified triolein and vegetable oils. *Food Chemistry* 1994, 50 (1): 39-43.

Rueh, H.; Matthews, E. and Willam, C. (1970).

The Use of some oil-seed in bread. Cereal Chem. 47(2): 181-185.

Sabir, M.A.; Sosulski, F.W. and Hamon, N.W. (1975).

Sunflower carbohydrates. J. Agric. Food Chem., 23, 16-19.

Saeed, M. and Cheryan, M. (1988).

Sunflower protein concentrate and isolates low in polyphenols and phytate. J. Food Sci., 53, 1127-1131.

Sania, A.E. (1980).

Separation and evaluation of cereal and legume proteins. Ph.D. in Food Science and Technology. Cairo University.

Santosh, K. and Chauhan, B.M. (1986).

Effect of domestic processing and cooking and in vitro protein digestability of moth bean. J. Food Sci., 51(4): 1083-1084.

Sarhan, M.A.; Mostafa, M.K. and Mortada, N.S. (1986).

Effect of soybean flour and improvers on bread quality. Egypt, J. Food Sci., 14, No. I.PP. 183-190.

Sastry, M.C. and Subramanian, N. (1984).

Preliminary studies on processing of sunflower to obtain edible protein Concentrates. J.A.O.C.S. 61, 1039-1042.

Sathe, S.K.; Ponte, J.G.; Rangekar, P.D. and Salunkha, D.K. (1981).

Effect of addition of great Northern bread flour and protein concentrate on rheological properties of dough and baking quality of bread. Cereal Chem. 58(2): 97-100.

Saunders, R.M. and Kohler, G.O. (1971).

In vitro determination of protein digestibility in wheat mill field for monogastric animals. Cereal Chem. 49: 98.

Savrasova, L.L.; Knyazhanskaya, B.A.; Vedernikova, E.L. and Bobova, L.S. (1983).

Use of sunflower seed protein isolate in breadmaking. Khlebopekarnaya Konditerskaya promyshlennost (1983) No. 2: 33-34.

Shahat, M.S. (1990).

Chemical and Technological studies on some oil-seeds proteins. Ph.D. Thesis, Dep. of Food Sci. and Technol., Faculty of Agric., Al-Azhar Univ.

Shaheen, A.; Osman, M.A.; Hashem, H. and Khalifa, A.A. (1991).

Production of Halva Tahinia from sunflower (*Helianthus annus*) seeds. 1: Chemical evaluation of sunflower and sesame seeds Egypt J. Food Sci. 19: 179187.

Shellenberger, F.A. (1982).

Processing and utilization in Agriculture. Vol. II, Part I, 91, J.A. Wolfed. CRC press, FNC. Boca. Raton. Pl.

Silaula, S.M.; Lorimer, N.L.; Zabik, M.E. and Ueversak, M.A. (1989).

Rheological and sensory characteristics of bread flour and whole wheat flour doughs and breads containing dry Roasted Air-classified pinto and Navy bean High-protein Fractions. Cereal Chem. 66: 486-490.

Smith, K.J. (1968).

A review of nutrition value of sunflower meal. Feed stuffs (40), 20, 22, 28, 60.

Smith, K.J. (1971).

Nutritional Framework of oil seed proteins. J.A.O.C.S., 48, 625-628.

Snedecor, G.W. and Cochran, W.G. (1967).

Statistical Methods. 6th ed. The Iowa State Univ. Press, Iowa, U.S.A.

Sosulski, F.W. (1979a).

Food uses of sunflower proteins. J.A.O.C.S., 56, 438.

Sosulski, F.W.; Kilara, A. and Humbert, E.S. (1974).

Nitrogen extractability and moisture absorption characteristics of sunflower seed products. J. Food Sci., 37, 771-773.

Sosulski, F.W.; Mc Cleary, C.W.; and Soliman, F.S. (1972).

Diffusion extraction of chlorogenic acid from sunflower kernels. J. Food Sci., 37, 253-256.

Sosulski, F.W.; Sabir, M.A. and Fleming, S.F. (1973).

Fontinuous diffusion of chlorogenic acid from sunflower kernels. J. Food Sci., 38, 468.

Szerdelyl, F. (1969).

Processing sunflower seed harvested in 1967/68. Olaj szappan kozmetika, 18, 105. C.F. FSTA, 12, d 710.

Szerdelyl, F. (1970).

Processing sunflower seed harvested in 1968/69. Libid, 19,2-6. C.F. FSTA, 2, 10N 404.

Szerdelyl, F. (1973).

Quality of Hangarian sunflower seed processed in season 1972/73. Libid, 22, 97-102. C.F. FSTA, 7, 164.

Timms, M.F.; Bottomley, R.C.; Ellis, U.R.C. and Schofield, J.D. (1981).

The baking quality and protein characteristics of a winter wheat grown at different levels of nitrogen fertilization. *Journal of science of food and Agriculture* 1981, 32(7): 684-698.

Varriano, M.E.; Huang, G. and Ponte, J. (1980).

Comparison of methods to determine starch gelatinization in bakery foods. *Cereal Chem.* 57: 242-246.

Wan, P.J.; Balcer, G.W.; Clark, S.P. and Matlock, S.W. (1979).

Characteristics of sunflower seed and meal. *Cereal Chem.* 56(4): 352-355.

Yamzaki, W.T. (1953).

An alkaline water retention capacity test for the evaluation of cooki baking potentialities of soft winter wheat flour. *Cereal Chem.*, 30: 242-249.

Youssef, S.A.M. and Abdel-Rahman, A.Y. (1976).

Quality and chemical constituent of sunflower seeds from different proteins of the bead. *Geasas Y Aceites*, 27.

Yue, P.; Hettiarachchy, N. and D'Appolonia, B.L. (1991).

Native and succinylated sunflower proteins use in bread baking. *Journal of food science* 1991, 56(4): 992-995, 998.

Zhuge, Q.; Persaud, J.N.; Posner, E.S.; Deyoe, C.W.; Seib, P.A.; Chung, D.S. (1991).

Isolation of gluten and starch from ground pearled wheat compared to isolation from flour. *Cereal Chem.* 68(4): 336-339.

Zimmermann, R. and Meith, G. (1984).

Functional properties of sunflower protein in wheat dough.

IV. Comparative investigations of the effectiveness of preliminary swelling of sunflower and soybean proteins isolats at different pH-values. *Nahrung* 1984, 28(9): 975-981.