

R E F E R E N C E S

Abbey, B.W. and Mark-Balm, T. (1988).

Nutritional quality of weaning foods prepared from composite flour of maize, ungerminated and germinated cowpea.

Nutr. Rep. Inter. 38(3):519-526.

Abdel-Kader, M.M. (1984).

Preparations of high quality-low cost meals. Project of Nutrition.

Academy of Scientific Research and Technology. (c.f. Seleet, 1990).

Abrahamsson, L.; Hambraeus, L. and Bengtsson, O. (1977).

Protein quality of home-prepared and industrially produced baby food. Nutr. and Metabolism, 21(1): 160-162.

Abul-Seoud, E.I.Y. (1984).

Natural toxic constituents in some Egyptian legumes M.Sc. Thesis, Fac. of Agric., Ain Shams. Univ.

Agren, G.; Hofvander, Y.; Selinus, R. and Vahlquest, B. (1969).

In: Protein Enriched Cereal Foods for World Needs. Ed. Max., Milner, Pub. Am. Assoc., Cereal Chemi Minnesota p. 278.

Ahmed, R.U.; Lowgren, M.; Velarde, N.; Abrahasson, L. and Hambraeus, L. (1981).

Study of home-prepared weaning foods for consumption in Bangladesh, with special reference to protein quality.

Ecology of Food Nutr. 11(2):93-102.

- Ahn, K.M.; Moon, S.J.; Choi, H.S. and Kwak, T.K. (1985).**
Development and evaluation of the supplementary
foods for Korean infants and Children.
Korean J. of Nutr. 18(4):259-271. (c.f. FSTA
19(6), 1987).
- Albrecht, W.J.; Mustakas, G.C. and Mcchee, J. E. (1966).**
Rate studies on atmospheric steaming and immersion
cooking of soybeans.
Cereal Chem. 43:400.
- Alekaer, N.S. (1976).**
Amino acid composition and biological value of
rice protein.
Voprosy Pitaniya 5:76-79. (c.f. FSTA 9(3), 317,
1977).
- Alvin, S. and Brain, F. (1976).**
Food legume processing and utilization. Ottawa,
Canada p. 15.
(c.f. Abul-Seoud, 1984).
- Alvina, M.; Vera, G. and Araya, H. (1989).**
Legume consumption in preschool children, effect
of energy density and type of preparation. Archi-
Archivos Latinoamericanos de Nutrition 39(2):
129-140. (c.f. FSTA, 23(6), ,1991).
- Ambrose, V.D.; Venkiat, S.; Mahadera, S. and Hussin, P.
(1969).**
Nutritive value and amino acid supplementation
of protein.
J. Nutr. Diet, 9,6(1):15-21.

Anon (1971).

Infant formula mixtures.

Agriculture Research (Washington) 19(3):3-4.

(c.f. FSTA 3(9), 329, 1971.

Antunes, P.L. and Sagrhieri, V.C. (1980).

Effect of heat treatment on the toxicity and nutritive value of dry bean (Phaseolus vulgaris var. Rosinka G₂) proteins.

J. Agric. Food Chem., 28:938.

A.O.A.C. (1980).

Official Methods of Analysis (13th Ed.). Assoc. of Official Analytical Chem. Washington. D.C. U.S.A.

Asfour, R.Y.; Tannous, R.I.; Sabry, Z.I. and Cowan, J.W. (1965).

Preliminary clinical evaluation with loubina mixtures.

Am. J. clin. Nutr. 17(3):148.

Ashworth, A. and March, Y. (1973).

Iron fortification of dried skim milk powder maize-soybean milk mixture (CSM). Availability of iron in jamaican infants.

Br. J. Nutr. 30:577.

Aykroyed, W.R. and Doughty, J. (1969).

Legumes in human nutrition.

FAO Nutrition studies No. 19.

Badi, S.; Pedersen, B.; Monowar, L. and Eggum, B.O. (1991).

The nutritive value of new and traditional sorghum and millet foods from Sudan.

Plant Foods for Human Nutrition 40(1):5-19.

Barja, I.; Munoz, P.; Soliman, G.; Vallejos, E.; Radrigan, M.E. and Tagle, M.A. (1974).

Infant feeding formula based on chickpeas (Cicer arictinum). Its use as the sole food in healthy infants.

Indian J. Nutr. and Diet. 11:335.

Bender, A.E. (1978).

Food processing and nutrition.

London, Academic Press pp. 95.

Bender, A.E. and Doell, B.H. (1957).

Biological evaluation of proteins. A new Concept.

Brit. J. Nutr. 11:140. (c.f. Sidky, 1986).

Billerbeck, F.W.; Hing, F.S. and Purvis, G.A. (1969).

Infant cereals. (Gerber products Co.) U.S.

Patent, 3:429, 711. (c.f. Chem. Abst. 70, 105309f)

Bognon, P. (1985).

[Corn broth].

Franch Patent application FR 2559035 A. (c.f.

FSTA 18(7), V176, 1986).

Bohm, C. (1963).

Rearing premature infants on new early food adepted to requirement.

Med. Ernährung, 4:40. (c.f. Nutr. Abst. Rev. 33:1087).

Bourne, G.H. (1968).

The nutrition and feeding of infants and preschool children in the developing countries.

In: Worl Rev. of Nutr. and Diet. by S. Karger Basel (Switzerland), 9, 85.

Brandizoeg, B.; Malleshi, N.G.; Svanberg, U.; Desikachar, H.S.B. and Mellander, O. (1981).

Dietary bulk as a limiting factor for nutrients with special reference to feeding of preschool children. 3-studies of malted flour from ragi sorghum and green gram.

J. Tropical Pediatrics, 27(4):184-189.

Bressani, R. and Scrimshaw, N.S. (1961).

In progress in meeting protein needs of infants and preschool children.

Proceedings of an International Conference. Washington, D.C. August 21.

Bressani, R.; Wilson, D.L.; Chung, M.; Behar, M. and Scrimshaw, N.S. (1963).

Supplementation of cereal proteins with amino acid: VI. Effect of amino acid supplementation of rolled oats improvement of food and proteins. M. Friedman Ed. Plenum Press, New York. (c.f. Valencia et al., 1988).

Brian, A.F. and Allan, G.C. (1977).

A chemical approach (3th Ed) Hodder and Stoughton London, Sydney, Auckland and Toronto. (c.f. Sidky, 1986).

British Nutrition Foundation (1981).

Salt in the diet.

Briefings paper, British Nutrition Foundation, No. 2, 24 pp.

Buffa, A. (1971).

Food Conservation on service, UNICEF, Paris, France. In: Food Technology and Development Part I. Processing Low Cost Nutrition Food for World's Hungry Children. Factors, Formula Processes. Unicef, Paris, France. (c.f. Food Engin. 43:79-86, 89-90.

Carballido, A. and Barragan, R.M.R. (1983).

Studies on prepared foods for infants. V. content of free sugars and starch.

Anales de Bromatologia 35(1):111-122.

(c.f. FSTA 16(12), , 1984).

Care India (1975a).

Project poshak. An integrated health-nutrition macro pilot study for preschool children in rural and tribal madhya pradesh. Vol. 1, New Delhi: Vienrendra printers.

----- (1975b).

Project poshak. An integrated health-nutrition macro pilot study for preschool children in rural and tribal Madhya pradesh. Vol. 2, New Delhi: Vienrendra printers.

Chan, J. and Lumen, B.O. (1982).

Properties of trypsin-inhibitor from winged bean seed isolated affinity chromatography.

J. Agric. Food Chem., 30:42.

Chandrasekhara, M.R.; Aswathanayan, S.; Shurpalekar, S.R. and Rao, G.H.S. (1969).

Pilot plant production of infant food based on ground nut and milk solids.

J. Food Sci. Tech. (India) 6:267.

Chatterjee, S.R.; Gassi, S. and Abrol, Y.P. (1974).

Amino acid composition of barley varieties.

J. Food Sci. Tech. (India) 11:85.

Church, C.F. and Church, H.N. (1969).

Food values of proteins commonly used. 11th

Ed. J.B. Lippincott Co., Phila., Toronto.

- Churella, M.R.; Yao, B.C. and Thomson, W.A.B. (1976).
Soybean trypsin-inhibitor activity of sors infant
formula and its nutritional significants for
the rat.
J. Agric. Food Chem. 24(2):393-397.
- Danial, A.V.; Kurien, D.N.; Swaminthan, M. and Parpia,
A. (1972).
Protein efficiency ratio of protein enriched
Cereal.
Nutr. Prep. Int., India, 5(5):346-349.
- Danial, A.V.; Leela, R.; Haribaran, K.; Swaminthan, M.
and Parpia, A. (1965).
Nutritive value and amino acid supplementation
of protein.
J. Nutr. Dietet. Coimbttra, India, 17:20.
- Davidson, S.; Passmore, R.; Broce, J.E. and Truswell, A.S.
(1979).
Human Nutrition and Dietetics. (7th Ed.).
The English Language Book Society and Churchill
Livingstone.
- Del Angel, A.R. and Sotelso, A. (1982).
Nutritive value of mixtures using ckickpeas with
wheat, triticales, normal and opaque -2 corns.
J. Nutr. 112:474.
- Desai, B.L.M.; Daniel, V.A.; Rao, S.V.; Swaminatham, M.
and Parpia, H.A.B. (1970).
Studies on low cost balanced foods suitable for
feeding weaned infants in developing countries.
Indian J. Nutr. Dietetics 7(1):21-26.

Devadas, R.P.; Jamala, S.; Chandraschara, U. and Murthy, N.K. (1974).

Nutrition evaluation of maize-based indigenous infant food.

Indian J. of Nutr. of Diet., 11(5):257-268.

Difco (1977).

Difco manual of dehydrated culture media reagents for microbiological and clinical laboratory procedures.

Difco, Lab. Inc. Detroit I, Michigan, USA, pp. 350.

Dodd, N.S. and Dutta, S. (1988).

Fat content and fatty acid composition of infant and weaning foods.

J. Food Sci. Tech., Indian 25(5):319-320.

Dodd, N.S. and Juoti, R. (1988).

Protein content and amino acid profile of selected commercial infant and weaning foods.

J. Food Sci. Tech., Indian 25(5):317-318.

Easwaron, P.P.; Gopinath, S.; Jamala, S. and Devade, R.P. (1972).

Evaluation of the protein quality of two selected vegetable protein mixtures using albino rats.

Indian, J. Nutr. Diet. 9(6):327-330.

Eggers, H. (1965).

The practice of artificial feeding of infants and the feeding of toddlers in greches and homes. Ernahrungs Forschung, 10:598. (c.f. Nutr. Abst. & Reviews 36,809).

Egyptian Organization for Standardization (1977).

Vegetarian baby food fortified with milk.

Ministry of Industry, Cairo, Egypt.

Eilenrieder, G.; Geromazzo, H. and De Bojarski, A.B. (1980).

Thermal inactivation of trypsin-inhibitors in aqueous extracts of soybeans, peanuts, and kidney beans, presence of substances that accelerate inactivation.

Am. Association of Cereal Chemists, 57.

Eka, O.U. (1978).

Chemical evaluation of nutritive value of soya paps and porridges, the Nigerian weaning food, Food Chemistry 3(3):199-206.

El-Hashimy, F.S. (1976).

Chemical and biological evaluation of some Egyptian leafy vegetables.

Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.

El-Malky, W.A. (1974).

Study on processing of some baby foods.

M.Sc. Thesis, Fac. of Agric., Ain Shams Univ.

El-Masry, E. (1973).

Preparation of supplementary meals for infants.

M.Sc. Thesis, Fac. of Agric., Alex. Univ., Egypt.

El-Tarras, M.F.; Abdel-Moely, E.M.; Ahmed, A.K.S. and Amer, M.M. (1976).

Studies on rancidity of oils and fats. XV. Stability of some infant foods.

Egyptian J. of pharmaceutical Sci., 17(2):161-172.

Ewida, E.H. and El-Nahry, F.I. (1988).

Saudian protein food mixtures for feeding infants and young children in Saudi Arabia.
Bulletin of the Nutrition Institute of the Arab Republic of Egypt 8(1):34-51.

Fahmy, A.A. (1969).

Studies on chemical and physicochemical changes of hydrogenated oils during storage.
M.Sc. Thesis, Fac. of Agric., Ain Shams Univ.

FAO (1957).

Protein Requirements.
Food Agric. Organ., Nutrition Studies, No. 16.

--- (1970).

Amino acid content of food and biological data on protein.
Nutrition Studies No. 24. (c.f. Food Sci. 43:1113, 1978).

--- (1973).

Energy and protein requirements.
FAO nutrition Meeting Rep., Ser. No. 52. FAO, Rome,

FAO/WHO Committee (1965).

Protein Requirements. World Health Organization.
Techn. Rep. Ser. 301. Food & Agriculture Organization W.H.O. Expert Group of the U.N. Geneva, Switzerland.

FAO/WHO Committee (1973).

"Energy and protein requirements".
FAO Nutrition Meeting Rep., Ser. No. 52.

---/--- ----- (1974).

Human nutritional requirements.
Handbook, (c.f. Nutr. Rev. 33(5), 1573, 1975).

---/--- ----- (1976a).

Food standards programme. Food Additives No.
26. Codex Alimentarius Commission, Eleventh
Session.

---/--- ----- (1976b).

Recommended international standards for food
for infants and children. Joint FAO/WHO food
standards programme CAC/RC 72/74. Food and Agric.
Orga. of the United Nations, World Health Organi-
zation. Geneva.

---/--- ----- (1985).

Food and Agriculture Organization of the United
Nations/World Health Organization. Energy and
protein requirements. Report of a Joint FAO/WHO/
UNU Expert consultation. Technical Report series
No. 724.

FAO/WHO/UNICEF (1966).

PAG Development of Algeria weaning food mixtures.
Doc., 10193.

---/---/----- (1970).

PAG statement No. 3 protein Advisory Group. United
Nations. New York, 10017, USA.

Fashakin, J.B.; Awoyefa, M.B. and Furst, P. (1986).

The application of protein concentrates from
locally available legumes in the development
of weaning foods.

Zeitschrift für Ernährungswissenschaft 25(4):
220-227. (c.f. FSTA 19(12), 1987).

Fernandez, R.; Elias, L.G.; Braham, J.E. and Bressani, R. (1982).

Trypsin-inhibitors and hemagglutinins in beans (phaseolus vulgaris), and their relationship with the content of tannins and associated polyphenols. J. Agric. Food Chem. 30:734.

Fetuga, B.L.; Rabatunde, G.M. and Oyenuga, V. (1973).

J. Sci. Food Agric. 24:1505.
(c.f. Ostrowski, 1980).

Filer, L.J. and Martinez, G.A. (1963).

Caloric and iron intake by infants in the United States.
Clin. Pediat., 2:470. (c.f. Nutr. Abst. & Rev. 34, 508, 1964).

----- and ----- (1964).

Intake of selected Nutrients by infants in the United States on evaluation of 4,000 representative six month olds.
(c.f. Nutr. Abst. & Rev., 35, 756, 1965).

Fomon, S.J. and Filer, L.J. (1967).

Amino acid requirements of normal growth.
In: Nyan, W.L. Ed. Amino Acid Metabolism and Genetic Variation. New York, Mc Graw Hill, p. 391.

Food and Nutrition Board (1958).

Recommended Dietary Allowances for Artificially Fed Infants.
(c.f. M.Sc. Thesis by Ebtesam El-Masry, 1973).

----- and ----- (1974).

National Academy of Sciences, National Research Council, U.S.A.
(c.f. Davidson et al., 1979).

----- and ----- (1980).

National Research Council U.S.A. (FNB, NRC).
Recommended Daily Allowances. (9th Ed.).
National Academy of Sciences, Washington, D.C.,
U.S.A.

Frazer, N.C. and Faster, E.M. (1950).

Laboratory Manual for Food Microbiology.
Burgess Publishing, Minnesota.

Fritz, J.C.; Ela, G.W.; Robert, T.; Boehne, J.W. and Hove, E.L. (1970).

Biological availability in animals of iron from common dietary sources.
J. Agric. Food Chem. 18:647.

Fromm, G. and Strahtmann, U. (1970).

Microbial content of baby foods, canned and deep frozen vegetables-a comparison study.
(c.f. FSTA 3(9), 550, 1971).

Gallardo, F.; Araya, H.; Pak, N. and Tagle, M.A. (1974).

Toxic factors in Chilean legumes. II. Trypsin-inhibitor activity.
Archivos Latinoamericanos de Nutricion (Caracas) 24:101.

Georgieva, F.; Chaneva, M.; Slavova, S. and Trifonova, L. (1988).

Heating regimes and nutritive values of canned fish and vegetable products for infant feeding.
Khranitelna promishlenost 37(4):26-28.
(c.f. FSTA 21(7), G9, 1989).

Golenkova, S.P. (1975).

Effect of stabilization on the content of amino acids in pumpkin puree with rice.

Konservnoi i Ovoshchesuchil noi pramyshlennosti, 33:34-41. (c.f. FSTA 8(7), 462, 1976).

Guggenbein, K. and Szmelcman, S. (1965).

Protein-rich mixture based on vegetable foods available in Middle Eastren Countries.

J. Agric. Food Chem. 51:148.

Guttikar, N.M.; Panemangalare, M.; Daraiswamy, R.T. and Swaminthan, M. (1966).

The metabolism of nitrogen digestibility coefficient and biological value of the protein.

J. Nutr. Diet., India 18:113.

Habash, L.A. (1977).

Some chemica, microbiological and physical studies on the production of some baby foods.

M.Sc. Thesis, Fac. of Agric., Cairo Univ., Egypt.

Hafez, Y.S. and Elgadri, S.S. (1986).

Proposed weaning formula for developing countries.

Nutr. Rep. Inter. 34(5):915-919.

Hansen, L.P.; Hosek, R.; Callan, M. and Jones, F.T. (1981).

The development of high-protein rice flour for early childhood feeding.

Food Technology 35(11):38-42.

Harper, J.M.; Hegazy, E.M.; Gabril, G.N. and Morsi, M.K. (1983).

Weaning food project, Food Technology Report. prepared for: Nutrition Institute and USAID, August 20, 1983.

Harris, D.A.; Burns, R.A. and Rida Ali (1988).

Evaluation of infant formula protein quality
Comparison of invitro with invivo methods.
J. Association of Official Analytical
Chemists 71(2):353-357.

Hawk, P.D.; Oser, B.L. and Summerson, W.H. (1949).

Practical physiological chemistry.
J. Nutr. 12:962.

Hegested, D.M. (1968).

Amino acid fortification and the protein problem.
Am. J. Clin. Nutr. 21:688.

Higashi, A.; Ikeda, T.; Uehara, I. and Matsada, I. (1982).

Effect of low-content zinc and copper formula
on infants nutrition.
European J. Pediatrics 3:237-240.

Ho, J.H. and Kim, S.H. (1970).

Protein rich food mixtures for Korean infants.
Korean, J. Nutr. 3:95. (c.f. FSTA 4(10), 488,
1972).

Hofvander, Y. and Underwood, B.A. (1987).

Processed supplementary foods older infants and
young children, with special reference to develop-
ing countries.
Food and Nutr. Bull. 9(1):1-7.

Hulse, J.H.; Rachie, K.O. and Billingesley, L.W. (1977).

Nutritional Standards and Methods of Evaluation
for Food Legume Breeders.
Ottawa, IDRC, Pub. CRDI, 65.

Hussein, M.A.H. (1987).

Production and evaluation of baby food mixtures from some raw and extruded starchy and proteinaceous materials.

Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.

Iotova, N.; Todorova, M. and Ivanova, S. (1988).

New, cereal-based dietetic foods.

Khranitelna Promishlenost 37(5):16-18. (c.f. FSTA 21(7), 1989).

Jamalian, J. (1978).

Favism-inducing toxins in broad beans (Vicia faba). Determination of vicine content and investigation on other non-protein nitrogeous compounds in different broad bean cultivars.

J. Sci. Food Agric. 29:136.

Jansen, G.R. and Howe, E.E. (1964).

World problem in protein nutrition.

Am. J. Clin. Nutr. 15:262.

Jelliffe, D.B. (1968).

Infant nutrition in the subtropics and tropics. (2nd Ed.) WHO, No. 29.

Kakade, M.L. and Evans, R.J. (1965).

Growth inhibition of rats fed navy bean fractions.

J. Agric. Food Chem. 13:450.

----- and ----- (1966).

Effect of soaking and germinating on the nutritive value of navy beans. J. of Food Sci. 31:781.

J. Food Sci. 31:781.

Kakade, M.L.; Rackis, J.J.; McGhee, J.E. and Puski, G.
(1974).

Trypsin inhibiting and hemagglutinating activities
in seed of some Legume species.
(c.f. Chem. Abst. 88(9), 1978).

Kakade, M.L.; Simons, N. and Liener, I.F. (1969).

An evaluation of natural vs. synthetic substrates
for measuring the antitryptic activity of soybean
samples.
Cereal Chem., 46:518.

Kapsiotis, G.D. (1969).

In: Protein Enriched cereal foods for world needs.
Ed. Max. Milner, Pub. Am. Assoc. Cereal Chem.
"AACC" Minnesota, U.S.A. p. 255.

Kas'yanov, G.I.; Petrichenko, L.K. and Lychkinal, L.V.
(1987).

Canned fish foods for infants.
Rybone Khozyaistvo No. 12:57-60. (c.f. FSTA 21(5),
1989).

Ketiku, A.O. and Olusanya, J.O. (1986).

Nutrient composition of multimixes for use as
weaning foods in Nigeria.
Food Chem. 21(1):47-56.

Khan, M.A. and Kissana, A.S. (1985).

Nutritional evaluation of some commercial baby
foods consumed in Pakistan.
J. Sci. Food Agric. 36(12):1271-1274.

King, K.W.; Jansen, G.R. and Anson, M.L. (1968).

Letter to the editor.

Am. J. Clin. Nutr. 21:525.

K.I.T. (Koninklijk Instituut voor De Tropen) (1983).

Ready-made weaning food mixtures in developing countries.

Caribs. Neerlandica/Mensen in Nood, the Netherland

Korte, R. (1972).

Heat resistance of phytohemag glutinins in weaning food mixtures containing beans (Phaseolus vulgaris).

Ecol. Food Nutr. 1:303-307.

Krasheninina, P.F.; Ushakov, V.V.; Mazo, V.K.; Gmoshinskii I.V.; Marokko, I.N. and Shirina, L.I. (1989).

Production of a dry product for children.

USSR Patent SU 1494898. (c.f. FSTA 22(5), 1990).

Kumarasinghe, P. and Abrahamsson, L. (1981).

Study on home-made weaning foods suitable for sri lanka.

Indian J. of Nutr. and Diet. 18(12):442-454.

Kuzdzal-Savoie, S. (1988).

Lipid profiles of some infant formulae available in France.

Technique Laitiere & Marketing No. 1034:36-40.
(c.f. FSTA 22(5), 1990).

Liener, I.E. (1962).

Toxic factors in legumes and other elimination.

Am. J. Clin. Nutr. 11:281.

----- (1982).

Factors affecting the nutritional quality of soya products.

American Soybean Association C/P quer 7. Madrid, 33.

Lumen, B.O. and Bels, P.S. (1981).

Inhibitors of trypsin and chemotrypsin in winged bean (Psophocarpus tetragonolobus) tubers.
J. Agric. Food Chem., 29:884.

Malleshi, N.G.; Daodu, M.A. and Chandrasekhar, A. (1988).

Development of weaning food formulations based on malting and roller drying of sorghum and cowpea
Inter. J. Food Sci. Tech. 24(5):511-519.

Marero, L.M.; Payumo, E.M.; Aguinaldo, A.R. and Homma, S. (1988).

Nutritional characteristics of weaning foods prepared from germinated cereals and legumes.
J. of Food Sci. 53(5):1399-1402.

Marero, L.M.; Payumo, E.M.; Aguinaldo, A.R.; Homma, S. and Igarashi, O. (1991).

Vitamin E constituents of weaning foods from germinated cereals and legumes. J. of Food Sci. 56(1):270-271.

Marinescu, J.; Lazar, T.; Baloscan, G.; Marin, E.; Zlota, R. and Agatestein, H. (1972).

Manufacture of "Sehior-food" canned foods for children, Lucrari de cercetara 10:97-102. (c.f. FSTA 6, (6), 1974).

Masha, A.C. and Svanbery, U. (1983).

Preparation of weaning foods with high nutrient density using flour of germinated cereals.
Food and Nutr. Bull. 5(2):10-14. (c.f. FSTA 16 (12), G44, 1984).

Masklarska, T. and Todorova, M. (1987).

Infant foods from bananas.

Khranitelna Promishlenost 36(6):26-27. (c.f.
FSTA 21(6), J42, 1989).

Mathew, S.P. and Pellett, P.L. (1986).

Protein quality of home made weaning food mixtures
I. Biological evaluation and quality.

Ecology of Food & Nutr. 19(1):31-40. (c.f.
FSTA 19(9), G14, 1987).

Mayer, A.M. and Pol Jakoff, M. (1963).

The Germination of Seeds.

International Series of Monographs on pue and
Applied Biology Plant. Physiology. Division 3(3).

Meneely, G. (1973).

Qual plant-plant Foods Hum. Nutr. 23(1/3).
c.f. Effect of processing on the sodium, potassium
and calcium : phosphorus content of foods.
J. Agric. Food Chem. 1583, 31:415-420.

Miller, D.S. and Bender, A.E. (1955).

Evaluation of protein quality.

Britsh J. Nutr. 9:382.

Miller, D.S. and Dema, I.S. (1958).

Proc. Nutr. Soc. 17:1.

(c.f. Sidky, 1986).

**Miric, M.; Miletic, I.; Lalic, Z. and Aksentijevic, S.
(1989).**

The composition and nutritional value of industri-
ally produced baby food. 11, Lipids and proteins.
(c.f. FSTA 21(1), A68, 1989).

Mitchell, H.H. (1924).

A method of determining the biological value of protein.

J. Biol. Chem., 58:873.

Mitchell, H.H.; Rynbergen, H.J.; Anderson, L. and Dibble, M.V. (1968).

Cooper's nutrition in health and disease.

J. B. Lippincott Company, Philadelphia, Toronto.

Mitkov, S.; Davidova, I. and Nikolova, T. (1969).

Chemical composition and organoleptic quality of preserved baby food containing meat.

Proceedings of the European Meeting of Meat Research Workers, 15:1189-1199.

Moore, S.; Spachman, D.H. and Stein, W. (1958).

Chromatography of amino acids on sulphonated polyslurenersins.

Anal. Chem. 30:1185-1190.

Morcos, S.R. and Gabrial, G.N. (1983).

Protein-rich food mixtures for feeding the young in Egypt. II. Chemical and biological evaluation. Qualitas Plantarum Plant Foods for Human Nutrition 32(1):75-81.

Morcos, S.R.; Elhawary, Z. and Gabrial, G.N. (1981).

Protein-rich food mixtures for feeding the young in Egypt. I. Formulation.

Food Sci. and Nutr. Res. Dept., National Research Centre, Dokki, Cairo, Egypt.

Moyls, A.W. (1958).

Apple base juices for infant.

Rep. Can Committee on Fruit and Vegetable Preservation. Can Dept. Agric., 40:1.

(c.f. Sidky, 1986).

Mrudula Kalpalathika, P.V.; Nanjundaswamy, A.M. and Patwaradha, M.V. (1988).

Studies on canned strained baby foods based on vegetables. I. Carrots.

J. Food Sci. Tech., India 25(3):137-141.

(c.f. FSTA 21(11),A10, 1989).

Myron, A.B. (1969).

Protein enriched cereal foods for world needs.

Am. Assoc. Cereal Chem. Minnesota.

Narayana, S.D.; Daniel, A.V.; Swaminathan, M. and Parpia, M. (1973).

Nutritive value of protein enriched cereal foods.

Nutr. Rep. Inter. India, 18:111.

Narayana, R.M.; Joseph, K.; Swaminathan, M. and Subrahmanyam V. (1961).

Development of precooked balanced protein food suitable for weaned infants.

Proc. Symp. Protein, Mysore, India, 9:286.

Narayana, R.M.; Rajagopalan, R. and Swaminathan, M. (1964).

A processed protein food based on a blend of peanut flour and full fat soybean flour.

J. Nutr. Diet. (Cimilatare-India) 9(1):1-3.

Narayana, R.M.; Rajagopalan, R. and Swaminathan, M. (1965).

A vegetable protein mixture based on peanut and soybean flour.

J. Am. Oil Chemists, 42(7):628.

Narayana, S.D. ; Amborase, V.D.; Rajalakshmi, M.L.;
Machadera, S.D. and Hussin, B. (1970).
Effect of supplementation with limiting amino
acids of PER of low cost protein foods.
Nutr. Rep. Inter. India 30:225.

Nasralla, E.M.K. (1978).
Evaluation of the protein value of some concent-
rated feedstuff.
M.Sc. Thesis, Fac. of Agric., Cairo Univ., Egypt.

Natters, L.A.; Mehta, T.; Mitchell, M.E. and Finney, P.L.
(1987).
Formulation and nutritive value of weaning food
from germinated food grains.
Nutr. Res. 7(12):1309-1320.

Ohlson, M.A. (1963).
Hand book of experimental and theraputic diets.
Brugess Pub. Company, Minneapolis 15 min.

Osborne, T.B. and Mendel, L.B. (1917).
The use of soybean as food.
J. Biol. Chem. 32:369-387.

Ostrowski, H.T. (1980).
Quantities and quality of protein extracted from
pasture herbage using heat precipitation or
ultrafiltration.
J. Sci. Food Agric. 31:177-187.

PAG (1970).
Guideline No. 7, FAO/WHO/UNICEF Protein Advisory
Group, Guideline for Human Testing of Supplement-
ary Food Mixture, 10 June, 1970. In: Food From
Waste. Pub. by Brich et al. (1976) pp. 5.

Pak, N.; Beriner, L.; Duffau, G.; Uacay, J. and Soriano, H.
(1974).

Infant feeding formula based on chickpeas. Its
use in the treatment of acute diarrhoea with
dehydration preliminary communication.
Indian J. Nutr. Diet. 12:42.

Patwardhom, V.N. (1962).

Pulses and beans in human nutrition.
J. Clin Nutr. 11:12.

Pavitt, F. (1988).

Energy density and consistency of traditional
African weaning foods.
Dissertation Abstracts International, B49(1):9
order No. BRD X 80485, 194 pp. (c.f. FSTA 20(6),
G12, 1989).

Pralhad Rao, N. and Vijayaraghavan, K. (1976).

Spin-off benefits in nutrition programmes.
Indian J. Med. Res. 64:1107-1118.

Produb, Z.; Pazoha, Z. and Geshak, J. (1970).

Powder puree from vegetables and cereals and
from fruits and cereals for children.
Spoz. 20:81-99. (c.f. Nutr. Abst., 41, 7912, 1971).

Protein Advisory Group (PAG) (1970).

Guideline for human testing of supplementary
food mixtures.
PAG New Bull. No. 10:80.

Proudfit, F.T. and Robinson, C.H. (1966).

Normal and therapeutic nutrition (12th Ed.).
The MacMillan Company, New York, USA.

Pushpamma, P. and Devi, C.A. (1979).

Nutritional quality of sorghum and legume based food mixtures for infants and pre-school children. Nutr. Rep. Inter. 19(5):635-641.

Rao, S.V.; Vijayalakshmi, D.; Kurien, S.; Prasannappa, G.; Chandrasekhara, H.N.; Swaminathan, M. and Chandrasekhard, M.R. (1976).

Nutritive value of weaning foods and malted milk powder.

J. Food Sci. Tech., India, 13(4):209-213.

Rettenbacher, M. (1990).

Extrusion of cereals for use in cereal-based foods for infants.

Ernährung 14(5):256-258. (c.f. FSTA 23(1),M60, 1991).

Richardson, M. (1982).

Thermal Stability of trypsin-inhibitor activity in winged bean (*Psophocarpus tetragonolobus*). J. Agric. Food Chem. 30:1140.

Roman, A.V.; Bender, A.E. and Morton, I.D. (1987).

Formulation and processing of a weaning food based on rice, cowpeas and skim milk powder. Human Nutrition.

Food Sci. Nutr. 41F(1):15-22.

Roy and Reo (1971).

Evidence, isolation, purification and some properties of a trypsin-inhibitor in *Lathyrus sativus*. J. Agric. Food Chem. 19(2):257.

Sadre, M.; Payan, R.; Donose, G. and Hedayo, H. (1972).

Protein food mixture for tran.

J. Amer. Diet. Asso. 60:131. (c.f. FSTA 4(8),G9, 1974).

Schaaf, H.J. (Heinz Schaaf OHG Nahrungsmittel-extrusionstechnik) (1989).

Process for manufacture of cereal products snacks and instant porridge type products.

German Federal Republic Patent Application DE 37 442 A1. (c.f. FSTA 21(10),V13, 1989).

Scrimshaw, N.S. (1959).

International of nutrition infection.

Amer. J. Med. Sci. 237-267.

Scrimshaw, N.S.; Wilson, P. and Bressani, R. (1960).

International of nutrition and infection.

J. Trop. Pediatrics. 6:37.

Sebrell, W.H. (1969).

In: Protein Enriched Cereal Foods for World needs.

Ed. Max. Milner, Pub. Amer. Assoc. Cereal Chemists Minnesota, U.S.A. p 32.

----- (1971).

Nutrition and Future. In: Proceedings of 6th Symposium on Nutrition Health in Near East. Fac. of Agric. Sci., Amer., Univ., Beirut.

Sedky, K.A.H. (1968).

Sulphur metabolism with special references to nutrition.

Ph.D. Thesis, Queen Elizabeth College, Univ. of London.

Segura, V.C.; Mahecha, L.G.; Moreno, S.B.E. and Rodriguez, G.S. (1988).

Development of an infant food product based on rice.

Archivos Latinoamericanos de Nutricion 38(2): 278-287. (c.f. FSTA 23(4), M76, 1991).

Select, F.L. (1990).

Preparation and evaluation of some infantile foods.

M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt.

Shahwan, A.A.K. (1976).

Chemical and technological studies on preparation of baby food from cereal and cereal products.

M.Sc. Thesis, Fac. of Agric., Cairo Univ., Egypt.

Sheikh, I.A.; Arshad, M.; Aslam, M. and Adil, R. (1986).

Preparation and nutritional evaluation of weaning food based on wheat, rice and soybean (soylac).

Pakistan J. of Scientific & Industrial Research 29(2):151-154.

Sidky, H.M. (1986).

Formulation and evaluation of baby food mixtures.

Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.

Sindair, H.M. and Dorothy, F.H. (1969).

Hutchison's Food and The Principles of Nutrition. 12th Ed.

Sister, L. and Raddy, P.R. (1984).

Development and evaluation of low-cost indigenous food mixez for preschoolers of Trichur, Kerala, I.A. Survey of nutritional adequacy among selected preschoolers and formulation of diets.

Indian J. Nutr. Diet. 21(7):241-251.

Skovgaard, N. (1969).

Standard for the microbiological quality of dried foods. In: the Microbiology of Dried Foods. Proceedings of the sixth International symposium on Food Microbiology. pp. 457-479. (c.f. FSTA 2(7),H23, 1970).

Smith, A.K. and Cirde, S.J. (1980).

Soybeans chemistry and technology (2nd pri). The AVI Pub. Co., West Port, Connectivt.

Solomadina, L.V.; Krasheninine, P.F.; Stolyarova, A.V.; Kasyanov, V.F.; Nikitenko, E.I.; Ladodo, K.S. and Gribakin, S.G. (1989).

Production of infant food. USSR Patent, SUI 486, 143. (c.f.FSTA 22(2), V22, 1990).

Sprang, F.J. (1969).

Bacteriological requirements for dried infant and weaning foods. In: The microbiology of dried foods. Proceedings of the sixth International Symposium on Food Microbiology. pp. 496-510. Bilthoven, The Netherlands. International Association of Microbiological Societes. (c.f. FSTA 2(7),H18, 1970).

Stanimirovic, D. Stanimirovic, S.; Miric, M.; Lalic, Z. and Radosevic, P. (1986).

The composition and nutritional value of industrially produced baby food. I. Carbohydrates and minerals.

Hrana: Ishrana. 27(4):189-193. (c.f. FSTA 20(9), G15, 1988).

Stegel, A.; Bumivatana, A. and Lineback, D.R. (1965).

Development, acceptability and nutritional evaluation of high protein soy supplemented rice noodles for that children.

Cereal Chem. 52:801.

Svanberg, U. (1990).

Dietary bulk in weaning foods and its effect on food and energy intake.

International Development Research Centre pp. 272-287.

Swaminathan, M. (1975).

Supplementary food based on oil seeds for infant and children.

Central Food Tech., Res. Inst., Mysore, India, J. of Scientific and Industrial Res. 34(6):329-335.

Tan, N.H. and Wong, K.C. (1982).

Thermal stability of trypsin-inhibitor activity in winged bean (Psophocarpus tetragonolobus).

J. Agric. Food Chem. 30:1140.

Tandon, O.B.; Bressani, R.; Scrimshaw, N.S. and Le Beau, F. (1957).

Nutritive value of beans.

Nutrients in Central American Beans.

J. Agric. Food Chem. 5:137.

Tannous, R.I.; Cowan, J.W.; Rinnu, T.; Asfour, R.Y. and Sabry, Z. (1965).

Mixtures for feeding infants and pre-school children in Middle East. 1. Development and evaluation.

Am. J. Clin. Nutr. 17:143.

- Torija I.M.E.; Carballido, A. and Barragan Ruiz, M.R. (1984)**
Studies on baby foods. VI. Crude and dietary fibre contents.
Anales de Bromatologia 36(1):137-149. (c.f. FSTA 18(2),A36, 1986).
- Tung, T.; Huang, P. and Linchen, H. (1961).**
Protein-rich foods for infants in Taiwan.
Taiwan i- Hsuoh-Hui, TS-China, 60:82-473.
- Turbin, N.V.; Gruzder, L.G. and Shebrak, E.A. (1975).**
Biological value of protein from grain crops.
USSR. (c.f. Chem. Abst. 82,13812g, 1975).
- Ulloa, J.A.; Valencia, M.E. and Garcia, Z.H. (1988).**
Protein content concentrate from chickpea: nutritive value of a protein concentrate from chickpea (Cicer arietinum) obtained by ultrafiltration and its potential use in an infant formula.
J. Food Sci. 53(5):1396-1398.
- Ulmer, K. (Gebruder Buhler AG) (1987).**
Process for manufacture of a product from starch containing raw materials.
European Patent EP 0129235 B₁ (1987) [Priority CH 83-3375 (83062)(Buhler, Uzwil, Switzerland].
(c.f. FSTA 20(4),V39, 1988).
- Vaidehi, M.P. (1988).**
Protein quality of nutri-mixbased on dehydrated sweet potato flour, soybean, defatted ground nut and milk powder.
Nutri. Rep. Inter. 37(4):723-728. (c.f. FSTA 20(9),J12, 1988).

Valencia, M.E.; Troncoso, R.; Higuera, I. (1988).

Linear programming formulation and biological evaluation of chickpea-based infant foods. Cereals Chem. 65(2):101-104.

Vogel, A.R. (1975).

A text book of practical organic chemistry. 3rd Ed. English Language. Book Society and Longman Group LTD.

Weihrauch, J.L.; Kinsella, J.E. and Watt, B.K. (1976).

Comprehensive evaluation of fatty acids in foods. VI. Cereal products.

J. Amer. Diet. Assoc. 68(4):335-340. (c.f. FSTA 9(3), M337, 1977).

Weyere, H. (1961).

Dried vegetables and fruits. A New way of supplementary infants diet.

Med. U. Ernährung 2:231. (c.f. Nutr. Abst. & Revs. 33, 165, 1963).

Wilson, D.; Fisher, H. and Enqua, E. (1974).

Principles of Nutrition. (3rd Ed.).

John W. Ley & Sons, Inc. New York, London, Sydney Toronto.

Winton, L. and Winton, W. (1958).

The Analysis of Foods.

New York, John Wiley & Sons, INC. Chapman & Hall, LTD.

Wolf, W.J. (1977).

Legumes: Seed composition and structure. Processing into protein products and protein properties. In: Food Protein. Ed. by Whitaker, J.R. and Tannenbaum, S.R. AVI Pub. Co. INC West Port, Connecticut.

WHO (1972).

From energy and protein requirements.
Tech. Rep. Ser. 503. (c.f. Habash, 1977).

--- (1988).

Nutrition and food safety - United States of
America.

International Digest of Health Legislation 37(3):
580-585. (c.f. FSTA 20(12), V10, 1988).

Zein, R.I. (1980).

Chemical and technological studies on baby foods.
M.Sc. Thesis, Fac. of Agric. Kafr El-Shikh,
Tanta Univ., Egypt.

Zimmerman, G.; Weismann, S. and Yannai, S. (1967).

The distribution of protein, lysin, methionine
and antitryptic activity in the cotyledons of
some leguminous seeds.

J. Food Sci., 32:129.