CONTENTS

		Page
INTRODUCTION		1
REVIEW OF LITERATUR	· 2	8
	and physical properties of fish	0
muscles:	and physical properties of fish	8
* Chemical comp	ogition	0
_		
trimethylamin	ozen storage on degradation of	^
_	•	9
* Color of fish	ges during frozen storage.	14
	storage at -20 on the functional	17
	of fish proteins:	20
* Protein solu	_	20
* Apparent vis	_	23
* Emulsifying	•	25
* Emulsion Sta	• •	28
	frozen storage on the	20
	ility of natural actomyosin (NAM).	30
	electrophoretic patteren of	30
_	e natural actomyosin during frozen	
storage.	,darar accomycorn daring frozen	32
-	properties of natural actomyosin	
	croscopic observation.	38
zuro zv. Brecoron m	.croscopic observacion.	30
MATERIALS AND METHOD)S:	41
	_,	
A) MATERIALS		41
B) CHEMICAL ANALYSIS	3 :	42
B.1. Moisture, crude	protein, and ash.	42
B.2. Crude fat.		42
B.3. pH value.		42
B.4. Determintion of	dimethylamine.	43
B.5. Formaldehyde co	ntent.	43

C) PHYSICAL MEASUREMENT:	44
C.1. Texture	44
C.2. Color	44
D) FUNCTIONAL PROPERTIES:	45
D.1. Protein solubility.	45
D.2. The total protein of mince homogenate.	45
D.3. Apparent viscosity.	46
D.4. Emulsifying capacity.	46
D.5. Emulsion stability.	46
E) NATURAL OF ACTOMYOSIN (NAM):	47
E.1. Extractability of NAM.	47
E.2. Emulsifying properties:	48
E.2.a. Emulsion activity Index (EAI).	48
E.2.b. Emulsion stability Index (ESI).	49
E.3. Polyacrylamide gel electrophoresis of NAM (S1).	49
E.4.a. Extractability of aggregates (P1 Fraction).	50
E.4.b. Polyacrylamide gel of electrophoresis of (S2).	50
D) Transmission electron microscopy.	51
F) Statistical analysis.	51
RESULTS AND DISSCUSSION:	54
Part I: Chemical and physical properties:	54
* Chemical composition.	54
* pH value.	56
* Dimethylamine.	60
* Formaldehyde.	68
* Texture.	76
* Color.	83
Part II: Effect of frozen storage at -20°C on the	
functional properties of proteins of minced	
fish.	94
* Changes in protein solubility.	94
* The total protein content of the homogenate.	10:

* Changes in apparent viscosity.	108
* Changes in emulsifying caapacity.	115
* Changes in emulsion staability.	122
Part III: Effect of frozen storage at -20°C on the	
natural actomyosin (NAM):	129
A.1. Extractability of natural actomyosin (S1	
Fraction).	129
A.2. Electrophoretic pattern of natural	
actomyosin (S1 fraction).	135
A.3. Emulsifying properties of natural	
actomyosin (S1 Fraction).	144
B.1. Extractability of precipitated aggregates	
(P1 Fraction).	152
B.1.a. Breaking down the secondary interactions	s.153
* Treatment with 2 % SDS.	153
* Treatment with 2 % SDS + 8M Urea.	153
B.1.b. Breaking down secondary interaction and	
disulfide bonds.	156
* Treatment with 2 % SDS + 5 % ß-mercapto_	_
ehanol (ME).	156
B.2. Polyacrylamide Gel Electrophoresis (S2	
Fraction).	158
* Protein extracted with 2 % SDS.	158
* Protein extracted with 2 % SDS + 8 M Urea.	160
* Protein extracted with 2 % SDS + 5 % ß-	
mercaptoethanol (ME).	163
Part IV: Transmission electron microscopy (TEM):	167
* Muscle morphology.	167
 Morphology of natural actomyosin (S1 	
Fraction).	167
 Morphology of precipitate residual of 	
natural actomyosin (Pl Fraction).	173
SUMMARY.	180
ARABIC SUMMARY	