

REFERENCES

REFERENCES

Abd-Elmoty, A.K.I.; A.H. El-Bogdady, and A.A. Faltas,
(1986):

Performance and physiological responses on growing LSL
chickens fed diets substituted with dried poultry
manure.

Egypt, Poult. Sci., 6: 65-83.

Abdo, M.S.; S.A. Mansour, and A.M. El-Nahla, (1983):

Effect of some feed additives on blood constituents of
growing Hubbard chickens.

Med. J., 31 (3):221-231.

Afifi, M.A., (1972):

Sunflower seed meal as a substitute for soyabean meal
in broiler rations.

Archiv für Geflügelkunde, 36 (4):129-134.

Alexander, J.C., and D.C.Hill, (1952):

The effect of heat on the lysine and methionine in
sunflower seed oil meal.

J. Nutrition, 48: 149-160.

A.O.A.C. "Association of Official Agricultural Chemists",
(1980):

Official Methods of Analysis. 10th Ed. Washington,
D.C.

Assy H.M., and N.S. Erian, (1980):

Biochemical studies on the fatty components of two sunflower varieties.

J. Agric. Sci. Mansoura Univ., 5: 263-270.

Attia, F.M.; N.F. Abdel Hakim, and S.E. El-Sharkawy, (1978):

Effect of diet, season and sex on blood cholesterol and glucose in Fayoumi chickens.

Agric. Res. Rev., 56:35-46.

Bell, D.J., (1957):

The distribution of glucose between the plasma water and the erythrocyte water in hen's blood.

Quart. J. Exp. physiol., 42:410.

Bell, D.J., (1971):

Plasma glucose in "Physiology and biochemistry of the domestic fowl".

Vol.2(D.J.Bell and B.M. Freeman, Ed.), London Academic press, P. 913-919.

Bell, D.J.; W.M. McIndoe, and D. Gross, (1959):

Tissue components of the domestic fowl. 3- The non-protein nitrogen of plasma and erythrocytes.

Biochem. J., 71:355.

Bell, G.A.; J.N. Davidson and D. Easlie-Smith, (1972):

Text book of physiology and biochemistry.

Longman, Edinburgh, Eighth Edition.

Bell, J.M., (1955):

Nutritional value of rapeseed oil meal.

Can. J. Agr. Sci., 35:242-251.

Brandt, L.W.; R.E. Clegg, and A.C. Andrews, (1951):

The effect of age and degree of maturity on the serum proteins of the chicken.

J. Biol. Chem., 105-111.

Campbell, E.A., (1959):

Effect of esterogen on blood volume and hemoglobin in immature pullets.

J. Physiol., 197: 1181.

Cherms, F.L., Jr.; F.H. Wilcox, and C.S. Shaffner, (1960):

Genetic studies of serum cholesterol level in the chicken.

Poult. Sci., 39: 889-892.

Clandinin, D.R., and A.R. Robbee, (1950):

The effect of methods of processing on the nutritive value of sunflower meal.

Poultry Sci., 29: 753.

Clandinin, D.R., (1958):

Sunflower seed oil meal processed plant protein.

Foodstuffs, Ed., A.M. Alt schull, Academic Press, New York.

Cornelius, C.E., (1957):

New concept and methods in the laboratory diagnosis of canine liver disease.

Gaines Veterinary Symposium, Kankakee, 111: 23.

Crawley, S.W.; D.R. Sloan, and K.K.Jr. Hale, (1980):

Yields and composition of edible and inedible by-products of broilers processed at 6,7 and 8 weeks of age.

Poult. Sci., 59: 2243-2246.

Cuca, M.; E.Avila, and E.Sasa, (1973):

Threonine supplementation to sunflower diets for chicks.

Poultry Sci., 52: 2016-2017.

Daghir, N.J., and S.L. Balloun, (1961):

Influence of fat and choline on liver and serum cholesterol in the chick.

Poult. Sci., 40: 1712-1717.

Daghir, N.J.; M.A.Raz, and M. Uwayjan, (1980):

Studies on the utilization of full-fat sunflower seed
in broiler rations.

Poultry Sci., 59: 2273-2278.

Day, H.C., and E. Levin, (1954):

The nutritional value of sunflower seed meal.

Sci., 101: 438-439.

Deaton, J.W.; F.N. Reece, and W.J. Tarver, (1969):

Hematocrit, hemoglobin and plasma protein levels of
broilers reared under constant temperatures.

Poult. Sci., 48: 1993-1996.

Deaton, J.W.; J.L. McNaughton, and D. Burdick, (1979):

High-fibre sunflower meal as a replacement for soyabean
meal in layer diets.

Br. Poult. Sci., 20: 159-162.

Dogan, K., and M. Zincirlioglu, (1985):

Protein quality of rapeseed oil meal and possibilities
of its use in broiler rations.

Doga Bilim Dergisi, Di (Veterinerlik ve Hayvancilik),
9(1): 35-44.

Dixon, M. and E.C. Webb, (1964):

Text book of enzymes. Longman, London. Second Edition.

Ducks, H.H., (1960):

The physiology of domestic animal.

Comstock Publishing Associates Ithaca, New York. P.
49.

Duncan, D.B., (1955):

Multible range and multible F test.

Biometrics, 11: 1-42.

Ehsan El-ansary; A. El-Naggar, and M.P. Asar, (1986):

Effect of glucagon hormone on carbohydrate metabolism
of fasting chickens.

Alex. J. Agric. Res. 31(1): 59-68.

El-Ansary, E.; M.A. Kosba;; M.M. Ali, and A. El-Nagar,
(1981):

Carbohydrate metabolism in two strains of chickens
selected for divergent body weight. 1. Blood glucose
concentration and effect of glucagon hormone adminis-
tration.

Alex. J. Agric. Res., 29(3): 1115-1123.

El-Bogdady, A.H.; A.A. Faltas, and A.K.I. Abd- Elmoty,
(1987):

Effect of starvation and transport on some physiologj-
cal parameters of LSL chickens.

Egypt Poult. Sci., 7: 26-36.

El-Husseiny, O.; A.I.Eissa, and S. Hashish, (1982):

Effect of dietary methionine and cystine levels and proportions on growth rate, certain hepatic enzymes, plasma and liver composition of broiler chicks.

Annals of agric. Sci., Moshtohor, 18: 107-122.

Essary, E.O.; C.E. Holmes, and W.L. Beane, (1964):

Influence of feeding to levels of protein and energy in broiler rations on live weight, feed conversion, dressing percentage and certain carcass evaluation.

Poult. Sci., 43:1316.

Essary, E.O.; L.E. Dawson; E.L. Wisman, and C.E. Holmes, (1965):

Influence of different levels of fat and protein in broiler rations on live weight, dressing percentage and specific gravity of carcasses.

Poult. Sci., 44: 304-304.

Ewing, W.R., (1947):

Poultry Nutrition.

W. Ray Publisher, South Pasadena, California, U.S.A.

Falts, A.A.; A.K.I. Abd-Elmoty, and Kh.A. Mohamed, (1987):

The effect of exposure at 40 C on the physiological response of the domestic fowl maintained at 16 C.

Minia J. Agric. Res. & Rev. 9(1): 383-398.

Featherston, W.R., (1969):

Nitrogenous metabolites in the plasma of chicks adapted to high protein diets.

Poult. Sci., 48: 646-652.

Freeman, B.M., (1971):

Metabolic energy and gaseous metabolism. In "Physiology and biochemistry of the domestic fowl".

Vol.1 (D.J. Bell and Bell and B.M. Freeman, Ed.), London, Academic press, P.279-391.

Gheyasudin, S.; C.M. Carter, and K.F. Mattil, (1970):

Effect of several variables on the extractability of sunflower seed proteins.

J. of food Sci., 35:453-456.

Gheyasudin, S.; C.M. Carter, and K.F. Mattil, (1972):

Effect of chlorogenic, quinic and caffeic acids on the solubility and color of protein isolates, specially from sunflower.

Cereal Chem., 49: 508-512.

Grau, C.R., and H.J. Almquist, (1945):

The value of sunflower seed protein.

Proc. Soc. Expt. Biol. Med., 60:373-374.

Grimbergen, A.H.M., and E.J.M.V. Kuiken, (1963):

Elektrophoretische untersuchungen von geflugelblutseren. Arch. Geflugel Kunde, 27:325.

Guyton, A.C., (1981):

Text Book 1- Medical Physiology. W. B. Saunders Company (sixth edition), London.

Hamdy, S.; M.A. Kosba; A.Z. Khalil, and Abo-Egla, (1980):

Carcass quality of chickens as affected by environmental temperature and diet.

Alex. J. Agric. Res. 28(3):39-48.

Hazelwood, R.L., and F.W. Lorenz, (1959):

Effects of fasting and insulin on carbohydrate metabolism of the domestic fowl.

Am. J. Physiol. 197:47.

Huston, T.M., and T. Subhas, (1968):

The influence of environmental temperature upon adrenal activity of bursectomized chicks.

Poult. Sci., 47:1760.

Jaap, R.G.; M.M. Renard, and R.D. Buckingham, (1950):

Dressed and eviscerated meat yields from chickens at 12 weeks of age.

Poult. Sci., 29: 874.

Kahle, H.S., and L.R. Gray, (1956):

Utilization and disposal of poultry by-products and wastes.

Agric. Mktg. Serv., Mktg. Res. Rep. No. 143, USDA, Washington, D.C.

Klain, G.J.; D.C.Hill; H.D. Branion, and J.A. Gray, (1956):

The value of rapeseed oil meal and sunflower seed oil meal in chick starter rations.

Poultry Sci., 1315-1326.

Kondra, P.A.; J.F. Richards, and C.C. Hodgson, (1960):

The effect of strain, sex and ration on meat yield in chicken broilers.

Poult. Sci., 39: 1265.

Kratzer, F.H.; P.N. Davis; D.E. Williams, and B.J. Marshall, (1954):

Factors influencing the growth of chicks and poult fed rations containing rapeseed oil meal.

J. Nutrition, 53:407-418.

Kumar, A.; D.N. Seth, and J.S. Rawat, (1974):

Distribution of serum protein fraction and non-protein nitrogen constituents in White Leghorn birds in relation to age, sex and reproduction.

Indian J. Anim.Sci., 44:771.

Labib, A.I., (1983):

Poultry nutrition course for post-graduate students.

Agric. Col., Zagazig Univ. Benha Branch.

Lautner, V., and Z. Zenisek, (1964):

Estimation of energy value of components of feed mixtures chickens.

Nutr. Abstrs. Reviews, 35: 540.

Lee, P.K., and Y.F. Yang, (1980):

Raw unhulled sunflower seed as feedstuff for broiler chicks. J. of the Taiwan Livestock Research, 13 (2): 49-57.

Lee, P.K., and M.C. Lee, (1982):

Effect of feeding locally produced sunflower oil as protein supplement on the performance of the broiler chicks.

Journal of the Taiwan Livestock Research, 15(2): 9-24.

Leeson, S., and J.D. Summers, (1980):

Production and carcass characteristics of broiler chickens. Poult. Sci., 59: 786-798.

Leveille, G.A.; J.W. Shockley, and H.E. Sauberlick, (1962):

Influence of dietary protein level and amino acids on plasma cholesterol of the growing chick.

J. Nutr., 76: 321.

Lortscher, L.L.; G.F. Sachsels; O.Jr. Whilhelmy, and R.B Jr.

Filbert, (1957):

Processing poultry by-products in poultry slaughtering plants.

Agric. Mktg. Serv., Mktg. Res. Rep. No. 181, USDA, Washington, D.C.

Mandev, I., (1974):

Sunflower meal as the chief source of protein starter feeds for broiler chicken.

Zhivot. Novudni Nauki 11:80.

Marinov, B.; L. Angelova, and I. Mandev, (1973):

Sources of protein in mixed feeds for chickens. 1-Sunflower seed oil meal.

Zhivot Novudni Nauki, 10:81.

Marinov, B.; D. Chotinski, and D. Mircheva, (1985):

Testes on sunflower oil meal from hybrid seeds as an ingredient in mixed feeds for broilers.

Zhivotnov dni Nauki, 22(8): 31-36.

Marian, J.E., and J.G. Woodroof, (1965):

Composition and stability of broiler carcass as affected by protein and fat.

Poult. Sci., 44:1396.

Mastakas, G.C., (1971):

Full-fat and defatted say flowers for human nutrition.
J. Am. Oil Chem. Soc., 48: 815-818.

McGinnis, J.; P.T. Hse, and J.S. Carver, (1984):

Nutritional deficiencies of sunflower seed oil meal
for chicks.
Poultry Sci., 27: 389-393.

Mehrez, A.Z.; M.El-S. El-Moghazy; M.M. El-Shinnawy; M.
Khalifa, and F.A. Abd-Alla, (1982):

Effect of varying dietary protein level and source of
animal protein on chick performance: 1-Growth and some
blood parametars of growing chicks.
J. Agric. Sci., Mansoura Univ., 7:180-196.

Milovanovic, M.; D. Ciric; V. Petrovic, and D. Delibasic,
(1972):

Relationship between dressing percentage and some
carcass cuts in broilers.
Anim. Breed. Abstr., 42: 4598.

Morgan, G.W., JR, and B. Glick, (1972):

Aquantitative study of serum proteins in bursectomized
and irradiated chickens.
Poult. Sci., 51:771-778.

Morrison, A.B.; D.K. Clandinin, and A.R. Robbee, (1953 a):
The effects of processing variables on the nutritive
value of sunflower seed oil meal.
Poultry Sci., 32: 492-496.

Morrison, A.B.; D.K. Clandinin, and A.R. Robbee, (1953 b):
The supplementary value of sunflower seed oil meal in
practical chick starting rations.
Poultry Sci. 32: 542-547.

Morrison, F.B., (1959):
Feeds and Feeding.
Twenty-second Editions, The Morrison Publishing Com-
pany, U.S.A.

Naber, E.C., (1976):
The cholesterol problem, the egg and lipid metabolism
in laying hen.
Poult. Sci., 55: 14-30.

Negm, H.M.; M.A. Kosba; S. Hamdy, and H.S. Zewel, (1980):
The effect of strain and sex on some economic traits
of chickens.
J. Agric. Res., Tanta Univ.

Newell, G.W., (1954):
Percentage yield of parts of cut-up broilers.
Poult. Sci., 33: 1074.

Nishida, T.; A. Yeno, and F.A. Kummerow, (1960):

Metabolism of cholesterol. 4-C¹⁴ in bile duct cannulated chicks and rats.

Circulation Res., 8:742.

N.R.C. "National Research Council", (1984):

Nutrient requirement of poultry.

8th Revised Ed. National Academy of Science, Washington, D.C.

Okumura, J., and I. Tasaki, (1969):

Effect of fasting, refedding and dietary protein level on uric acid and ammonia content of blood, liver and kidney in chickens.

J. Nutr., 97: 316-320.

O'Neil, J.B., (1948):

Mustard seed oil meal and sunflower seed oil in chick starter rations.

Sci. Agric., 28:127-130.

Orr, L., and T. Moran, (1975):

Effects of age and sex on carcass quality and yield in broilers.

Anim. Breed. Abst., 45(1): 976.

Ota, H., and E.H. McNally, (1961):

Poultry respiration calorimetric studies of laying hens. U.S. Dept. of Agric., A.R.S., 42.

Pettit, J.H.; S.J. Slinger; E.V. Evans, and F.N. Marcellus, (1944):

The utilization of sunflower seed oil meal, wheat distillers dried grains and rapeseed oil meal in poultry rations.

Sci. Agric., 24:201-213.

Plavnik, I., and S. Hurwitz, (1983):

Organ weights and body composition in chickens as related to the energy and amino acid requirements: Effects of strain, sex and age.

Poult. Sci., 62:152-163.

Pudelkiewicz, W.J.; R.V. Boucher; E.W. Callenbach, and R.C. Miller, (1959):

Some physiological responses of New Hampshire chickens to a mixture of penta and hexachloronaphthalenes.

Poult. Sci., 38:424.

Rad, F.H., and K. Keshavarz, (1976):

Evaluation of the nutritonal value of sunflower meal and the possibility of sunflower meal for soybean meal in poultry diets.

Poultry Sci., 55: 1757-1765.

Rako, A.; F. Dumanovsky, and K. Mikulec, (1964):

On the relation between the laying capacity and the activity of some enzymes, the level of serum proteins and blood sugar in hens.

Poult. Sci., 43: 201-204.

Rasheed, A.A., and J.A. Ondfield, (1963):

Comparitive data on growth, feed intake, feathering and meat quality.

Poult. Sci., 42:1001-1009.

Renner, R.; D.K. Clandinin; A.B. Morrison, and A.K. Robbee, (1953):

The effect of proces temperatures on the amino acid content of sunflower seed oil meal.

Poultry Sci., 32: 922.

Rico, A.G.; J.P. Braun; P. Benard; A.A. Hassan, and C.A. Ali, (1977):

Tissue distribution and blood levels of gamma glutamyl transferase in the horse.

Equine Vet. J., 9(2): 100-101.

Rico, V.O., (1978):

Performance of Cobb broiler chicks fed with commercial mash supplemented with dried unprocessed sunflower (Helianthus sp.) seeds.

TCA Res. J., 1(1): 63-73.

Rose, R.J.; R.N. Coit, and J.L. Sell, (1972):

Sunflower seed meal as a replacement for soybean meal protein in laying hen rations.

Poultry Sci., 51: 960-967.

Rotter, B.; W.Guenter, and B.R. Boycatt, (1985):

Sudden death syndrome in broilers dietary fat supplementation and its effect on tissue composition.

Poult. Sci., 64(6): 1128-1136.

Salander, R.C., and Hans Fisher, (1956):

Plasma amino nitrogen and creatine values of growing chick and laying hen.

Proc. Soc. Exp. Biol. & Med., 92:538.

Scott, M.L.; M.C. Nesheim, and P.Y.Young, (1969):

Nutrition of the chicken.

Ed. Humprey Press. Geneva, New Yourk.

Selim, A.D.; F.E. Abdel-Salam; A.A. Aboulseoud; A.K. Abou-Raya, and M.A. Ghany, (1976):

The effect of variable dietary energy and protein with holding the calorie protein on growing chicks.

F.S.T.A. 8(12):220.

Setchell, B.P., (1959):

Nature, 184: 1242, Cited by El-Amrousi et al. (1965).

Vet. Med. J. Giza, 10:291-294.

Setchell, B.P., (1961):

Aust. J. Agric. Res. 12(5): 944. Cited by El-Amrousi

et al. (1965). Vet. Med. J. Giza, 10: 291-294.

Shelef, L.A., and L.R. Morton, (1976):

Soybean protein foods.

Food Technol., 30(4): 44-48.

Sherlock, (1975):

Diseases of liver and biliary system.

5th ed. Oxford Blackvell acienlfre publisher.

Shim, K.F.; Y.M. Sin, and P.G. Pee, (1979):

Different protein and energy levels on the growth and
serum protein of broilers.

Singapore J. of primary Industries, 7 :29.

Siller, W.G., (1959):

Avian nephritis and visceral gout.

Lab. Investig., 8: 1319.

Singh, G., (1981):

The serum protein pattern of chickens during the early
growing period using crossed immunoelectrophoresis.

Poult. Sci., 60:674-678.

Singh, K.S., and C.M. Prasad, (1979):

Feeding value of sunflower and groundnut cakes for broilers. Animal Feed Science and Technology, 4(2): 143-159.

Smith, C.J., (1969):

Environmental influences of the blood glucose level of young chicks.

Poult. Sci., 48:1873.

Smith, C.J., (1972):

Blood glucose levels in young chickens. The influence of high regimes.

Poult.. Sci., 51: 268-273.

Smith, K.J., (1968):

A review of the nutritional value of sunflower meal.

Foodstuffs, 8:20.

Smith, O., (1976):

Whey extrusion cooking cereal.

Food World, 21: 4-9.

Smith, R.E., (1966):

Importance of an accurate reference diet in the evaluation of protein for chick growth using plasma amino acid titers. J. Nutr., 89: 271.

Sosulski, F.W., and G. Sarwar, (1973):

Amino acids composition of oil seed meals and protein isolates.

Can. Inst. Food Sci. Tech. J., 6: 1-5.

Staron, T., (1977):

Les nouvelles sources de proteines alimentaires.

Recueil de travaux et de conferences. I.N.R.A., Luce, C.F., Gassmann.

Steele, R.G.D., and J.H., Torrie, (1980):

Principals and procedures of statistics.

2nd ed. McGraw Hill Book Co. Inc., New York.

Sturkie, P.D., (1954):

Avian physiology (1st ed.) Cornell University press, Ithaca, New York.

Sturkie, P.D., (1961):

The effect of age and reproductive state on plasma uric acid levels in chickens.

Poult. Sci., 40:1650.

Sturkie, P.D., (1965):

Avian physiology. Comstock Publishing Associates, a division of Cornell University Press, Ithaca, New York.

Sullivan, T.W.; F.L. Chermes, and M.L. Sunds, (1958):

Giblets yields and dressed weight of four classes of poultry.

Poult. Sci., 37:736.

Summers, J.D., and S. Leeson, (1979):

Composition of poultry meat as affected by nutritional factors.

Poult. Sci., 58: 536-542.

Tanabe, Y., and F.H. Wilcox, (1960):

Effect of age, sex and line on serum alkaline phosphatase of the chicken.

Proc. Soc. Exp. Biol. & Med. 103:68.

Tapper, D.N., and M.R. Kare, (1956):

Distribution of glucose in blood of chicken.

Proc. Soc. Exp. Biol. and Med. 92:120.

Tasaki, I., and J. Okumura, (1964):

Effect of protein level of diet on nitrogen excretion in fowls.

J. Nutr., 83:34-38.

Temperton, H.; F.J. Dudley, and G.J. Pickering, (1965):

Phosphorus requirements of poultry. 4. The effects on growing pullets of feeding diets containing no animal protein or supplementary phosphorus.

Br. Poult. Sci., 6:125-133.

Thomas, O.P.; R.S. Martin; J.P.H. Wessels, and J.B.B. Human, (1965):

Sunflower meal as a source of protein for chickens rations. S. Afr. J. Agric. Sci., 8: 1061.

Titus, K.W., (1971):

The scientific feeding of chicken.

5th ed. The Interstate Printers and Publishers. Inc. Danville, Illinois.

Tosovsky, J.; R. Cihak, and M. Tosovska, (1976):

The composition of growth and carcass quality of Ross and Hypeco broilers.

Anim. Breed. Abst., 45: 5645.

Valdivie, M., (1976):

La harina de girasol en la alimentacion avicola.

Rev. Div. Agrop., 6:38.

Valdivie, M.; E. Aragon, and H. Jordan, (1976):

Broiler diets with high contents of seed meal and different energy: protein in rations.

Cuban J. Agric. Sci., 10: 297.

Valdivie, M.; O. Sardinas, and J.A. Garcia, (1982):

The utilization of 20% sunflower seed meal in broiler diets. Cuban J. Agric. Sci., 16: 167.

Vaughan, J.G., (1970):

The structure and utilization of oil seeds.

Chapman and Hall Ltd., London, England.

Waldroup, P.W.; C.M. Hillard, and R.J. Mitchell, (1970):

Sunflower meal as a protein supplement for broiler diets. Feedstuffs, 42:41.

Ward, M.A., and Peterson, R.A., (1973):

The effect of heat exposure on plasma uric acid lactate dehydrogenase chloride, total protein and zenk in the broiler.

Poult. Sci., 52:1671.

West, E.S.; E.T. Wibert; H.S. Manson, and J.V. Bruggen, (1968):

Text Book of Biochemistry. The aucmillan, Comp. New York, London.

Williams, F.I., and S. Rodbard, (1960):

Increased circulating plasma volume following di-
benzylamine. Am. J. Physiol., 198:169.

Yanovich, V.G., (1983):

Fat supplements in diets for animals.

Zhivotnovodstvo, 5:39-42.