

References

- Alam, M.M.; O.G. Ali; A. Masood and A.M.Kham (1977).** Studies on the chemical change induced the infection of the root-knot nematode (*Meloidogyne incognita*) in tomato and eggplant and the stunt nematode (*Tylenchorhynchus brassica*) in cabbage and cauliflower roots. (C.F. Hort. Abstr. 47 (6) : 5661).
- A.O.A.C (1970)** Official methods of Analysis of the Association of official Agriculture chemists 11th Ed., Washington D.C., U.S.A.
- A.O.A.D (1995):** Yearbook of Agricultural Statistics; Vol. (15). Arabic Organization for Agricultur Development.
- Bailey, D.M. (1941):** The seedling test method for root-knot nematode resistance. Proc. Amer. Soc. Hort. Sci. 38:573-575.
- Bajaj, K.L. and R. Mahajan (1977).** Phenolic compounds in tomato susceptible and resistant to *Meloidogyne incognita* (Kofoid & white) Chitwood (nematode). Nemat. Mediterr. 5(2): 329-333.
- Bajaj K.L.; Y.K. Arora and R. Mahajan (1983).** Biochemical differences in tomato cultivars resistant and susceptible to *Meloidogyne incognita*. Helm. Abstr. 52(4): 1541.
- Bajaj, K.L.; P. Singh and K. Mahajan (1986).** Changes induced by *Meloidogyne incognita* in superoxide dismutase, Peroxidase and oxidase activity in tomato roots: Helm. Abstr. B 55(1): 96.
- Barham, W.S. and J.W. Sasser (1956).** Root-Knot nematode resistance in tomato. Proc. Assoc. Sou. Agric. 53. 173-176.

- Barham, W.S. and N.N. Winstead (1957).** Inheritance of resistance to root-knot nematodes in tomatoes. Proc. Amer. Soc. Hert. Sci. 69: 372-377.
- Barrons, K.C. (1939).** Studies of the nature of root knot resistance. J.Agric. Res. 58: 263-271.
- Bhatt, G.M. (1971):** Heterotic performance and combining ability in a diallel cross among spring wheats (*T. aestivum* L.). Austr. J. of Agric. Res. 22:359-369.
- Canto-Saenz, M. (1983):** The nature of resistance to *Meloidogyne incognita* (Kofoid & White 1919) Chitwood 1949, pp. 160-165. In Proc. Third Res. and Plann. Conf. on Root-knot Nematodes, *Meloidogyne* spp., March 22-26, 1982, ed. C.C. Carter, International *Meloidogyne* Project, Lima, Peru. 233 pp.
- Castagnone-Sereno, P.; M. Bongiovanni, and A. Dalmasso (1994).** Reproduction of virulent isolates of *Meloidogyne incognita* on susceptible and Mi-resistant tomato. Journal of Nematology. 26(3): 324-328.
- Castagnone-Sereno, P.; M. Bongiovanni and Apalloix and A. Dalmasso (1996).** Selection for *Meloidogyne incognita* virulent against resistance from tomato and pepper and specificity of the virulence/resistance determinants. European Journal of plant pathology 102:585-590.
- COPR (1983). (Centre for Overseas Pest Research)** Pest control in tropical tomatoes overseas development administration London, 130 pp.
- Desouki, I.A.M.; M.A. El-Sherif and A.A. Hassan (1982).** Evaluation of some sources of root knot nematode resistance in tomatoes. Egypt. J. Hort. 9:233-253.
- Dropkin, V.H. (1954).** Infectivity and gall size in tomato and cucumber seedlings infected with *Meloidogyne incognita* var *acrita* root knot nematode. Phytopathology 44:43-49.

- Dropkin, V. H. (1972).** Pathology of *Meloidogyne* -galling, gaint cell formation, effects on host physiology. Eu. Mediterr. Plant Port. Organ. (OEPP/EPPO) Bull. 6:23-32.
- Dropkin, V.H. and R.E. Webb (1967).** Resistance of axenic tomato seedlings to *Meloidogyne incognita acrita* and to *M. hapla*, Phytopath. 57 (6): 584-587.
- Eisenback, J.D.; H. Hirschmann; J.N. Sasser and A.C. Tviontaphyllou (1981).** Aguide to the four most common species of root-knot nematodes with a pictorial key. A coop, pub. Deps. plant pathol. and Genetics. North carolina state Univ. and U.S. Agency Int, Dev., Raleigh, North Carolina N.C. 48 PP.
- El-Gindi, D.M. and F.F. Moussa (1979).** Distribution and identify of *Meloidogyne species* and thier biotypes in Egypt. Proceeding of the second Research planning conference on Root knot nematodes, *Meloidogyne spp.* November 26-30, 1979 Athens, Greec.
- El-Sherif, M.A.; M.H. Belal and T.Badra (1980).** Activity of polyphenol oxidase in tomato healthy and infected by *Rotylenchus reniformis* and *Meloidogyne incognita*. Anz, Schadl. Pflanz. Umwelt. 53 (1): 10-13.
- Fassuliotis, G. (1970).** Resistance of *cucumis spp.* to the root knot nematode, *Meloidogyne incognita acrita*. J.Nematol., 2,174-178.
- Fassuliotis, G. (1979).** Plant breeding for root knot nematodes spp. 425-453. In : F. Lamberti and C.E. Taylor, eds. Root-knot nematodes *Meloidogyne species* Systematics, biology and control. New York: Academic press.
- Fassuliotis, G. and D.P. Bhatt. (1982).** Potential of tissue culture for breeding root-knot nematode resistance into vegetables. J. Nematol. 14: 10-14.
-

- Fatunla, T. and A. Salu (1977).** Breeding in tomatoes. J. Agric. Sci. 88(1): 187-191.
- Freeaz, L.C.C.B. and M.G.C. Churata-Masca (1985).** Evaluation of tomato cultivars inoculated with the root knot nematode *Meloidogyne incognita*. Helm. Abstr. B. 54(1): 113.
- Ganguly, S. and D.R. Dasgupta (1982).** Characterization of polyphenol oxidase from susceptible and resistant varieties of tomato inoculated with the root knot nematode, *Meloidogyne incognita*. Ind. J. Nemat. 12(2): 320-329.
- Gardner, C.O. (1963)** Estimates of genetic parameters in crosses fertilizing plants and thier implication in plant breeding, P. 225-252. In: W.D. Hanson and H.F. Robinson (eds.) Statistical genetics and plant breeding. Nat. Acad. of Sci. Nat. Res. Counc. Pub. 982.
- Ghosh, P.K. and M.M. Syamal (1994).** path analysis and correlation studies in tomato *lycopersicon esculentum* Miller. Orissa Journal of Horticulture 22(1/2) 41-45. (C.F. PL. Breed. Abster (1996) 66(11) 1602.
- Ghosh, P.K.; M.M. Syamal; N. Rai and A.K. Joshi (1995).** Improvement of hybrid tomatoes. Advances in plant sciences 8(1) 207-213 (C.F. pL. Breed. Abstre. 1997. 67(8) 1144.)
- Giebel, J. (1974).** Biochemical mechanisms of plant resistance to nematodes. J. Nemat. 6:175-184.
- Gilbert, J.C. and D.C. McGuire (1956).** inheritance of resistance to severe root knot from *Meloidogyne incognita* in commercial types tomatoes. Proc. Amer. Soc. Hort. Sci. 68: 437-442.
- Gomez, K.A and A.A. Gomez (1984)** Staitical procedures for agricultural research 2nd. John Wiely and Sons. New. York.

- Goodey, T. (1951).** Laboratory methods for work with plant and soil nematodes. Minn. Agric. and Fish. Tech. Bull. No.2 P.25.
- Griffing, B (1956).** Concept of general and specific Combining ability in relation to diallel crossing systems. Austral. J. Biol. Sci 9:463-493 (J. Amer. Soc. Hort. Sci. 107(2) :243-247.
- Guinez, S.A. (1983).** Behaviour of seven tomato cultivars in soil infested with a high population of *Meloidogyne incognita*. Pl. Breed. Abstr. 53(7): 5896.
- Hadisoeganda, W. W. and J.N. sasser (1982).** Resistance of tomato, bean, southern pea and garden pea cultivars to root-knot nematodes based on host suitability. PL. Dis. Rept. 66(2): 145-150.
- Haroon, S.A.; A. Abdul-Baki and R.N. Huettel (1993).** An in vitro test for temperature sensitivity and resistance to *Meloidogyne incognita* in tomato. J.Nematol. 25:83-8.
- Harrison, A.L. (1960).** Breeding of disease resistant tomatoes with special emphasis on resistance to nematodes. In: compbell soup company "proceedings of Plant Science Seminar", PP. 57-75. Camden, N.J.
- Hartman, K.M. and J.N. Sasser (1985).** Identification of *Meloidogyne species* on the basis of differential host test and perinea-lpattern morphology. pp. 69-78. In: Barker, K.R, Carter, C.C. and Sasser, J.N. (Eds), Advanced treatiseon *Meloidogyne*. Raleigh, North Carolina U.S.A., North Carolina Univ. craphics.
- Hashem, E.K (1977).** Evaluation of some tomato cultivars under local conditions. M.Sc. thesis. Fac. Agric., Alex. Univ., Egypt.
- Hassan, H.M (1990).** Studies on the nature of resistance to root-knot nematode in tomato. Ph. D. Thesis, Fac. Agric., Alex. Univ., Egypt.

- Hassan, N. and S.K. Saxena (1977).** Effect of inoculating tomato with root-knot nematode, *Meloidogyne incognita* on the phenolic content and polyphenol oxidase activity. Symp. on physiology of host-pathogen interaction. Nov. 25-26, 1977, Madras. India.
- Hassan, A.A.; M.A. Elshrif; S.E. Moustafa and G.S. Shohla (1980).** Screening tomatoes *Lycopersicon esculentum* for resistance to root-knot nematodes under field and greenhouse conditions. Res. Bulletin, Fac. Agric., Ain shams Univ. 1404: 10 pp. (C.F. Hort. Abstr. 51, 7840).
- Hernandez, T.P.; J.C. Miller; M.J. Giamalva and W. Birchfield (1965).** Inheritance of resistance to root-knot nematode group in tomatoes. Proc. Amer. Soc. Hort. Sci 87: 412-414. (C.F. Pl. Breed. Abstr. 36:217)
- Hirshmann, H. (1985).** The genus *Meloidogyne* and morphological characters differentiation its species. pp. 74-93., In: J.W. Sasser and C.C. Carter, eds. An advanced treatise on *Meloidogyne*. Vol. 1: Biology and control. North. carolina state Univ. Graphics.
- Houssny, H.H and B.A. Oteifa (1956).** preliminary field tests for evaluating some tomato varieties for resistance to root-knot nematodes (*Meloidogyne spp.* PL. Dis. Rept. 40:974-976.
- Hung, C.L. and R.A. Rohde (1973).** Phenol accumulation related to resistance in tomato to infection by root-knot and lesion nematode. J. Nematol. 5(4): 253-258.
- Ibrahim, I.K.A.; M.A. Rezk and M.A. ElSaedy (1976).** Occurrence and distribution of plant parasitic nematodes in northern Egypt. Alex. J. Agric. Res. 24:93-100.
- Johnson, A.W. and G. Fassuliotis (1984).** Nematode Parasites of vegetable crops. pp. 323-372, In: Nickle, W.R.ed. plant and insect nematodes. Marcel. Dekker Inc., New York and Basel.

- Kalloo, R.K.J and D.S. Bhatti (1979).** Inheritance studies on resistance to root-knot nematodes in tomato *Lycopersicon esculentum*. (C.F. Pl. Breed. Abstr. 49(5): 4181.
- Kaloshian, I.; J. Yaghoobi; T. Liharska; J. Hontelez; D. Hanson; P. Hogan; T. Jesse; J. Wijbrandi; G. Simons; P. Vos; P. Zabel and Vm. Williamson (1998).** Genetic and physical localization of the root-knot nematode resistance locus Mi in tomato. Molecular-and-General-Genetics. 257:3, 376-385; 43 ref.
- Kamar, M.E.A (1967).** Studies on the inheritance of root-knot nematode resistance. The effect of root-knot nematodes-*Fusarium* and their combination on yield and quality of tomato. M.Sc. Thesis, Fac. Agric., Alex. Univ., Egypt.
- Kandakova, E.I.; B.V. Kvasnikov and S.I. Ignatova (1979).** The resistance of greenhouse tomato cultivars and hybrids to *Meloidogyne incognita*. Helm. Abstr. 48(1): 184.
- Kumari, N.V. and M. Subramanian (1994)** Genetic variability in tomato. Maras. Agriculture Journal 81(12) 651-653. (C.F PL. Breed. Abster. 1995).
- Lamberti, F. (1979).** Economic importance of *Meloidogyne* spp. in subtropical and Mediterranean climates, P341-357. In: F. Lamberti and C. F. Taylor (eds). Root-knot nematodes (*Meloidogyne species*): Systematics, biology, and control. Academic, New York.
- MAFF (1983).** Control of pest and diseases of protected crops tomatoes. Min. of Agric. Fisheries and Food. London, MAFF Pub., Booklet 2243-110 PP.
- Masood. A. and S.I Hussain (1979).** Phenolic and orthodihydroxy phenolic changes and their role in the resistance and susceptibility of three tomato varieties to *Meloidogyne incognita*. Helm. Abstr. B48(3): 1119.
-

- McClure, M.A.; T.H. Kruk and I. Misaghi (1973).** A method for obtaining quantities of clean *Meloidogyne* eggs. J. Nematol. 5:230.
- McFarlane, K.C.; E. Hartzler and W.A. Frazier (1946).** Breeding tomatoes for nematode resistance and for high vitamin C content in Hawaii, Proc. Amer. Soc. Hort. Sci. 47:262-271.
- Medina Fiho, H.P. and M.A. Stevens (1980).** Tomato breeding for nematode resistance : survey of resistant varieties for Horticultural characteristics and genotype of acid phosphates. Acta Hort. 100: 383-393.
- Narayana, Y.D. (1979).** Studies on the nature of resistance in tomato (Var. NTDR-1) to root-knot nematodes. Helm. Abstr. B.48(4): 1578.
- Narayana, Y.D. and D.D.R. Reddy (1980).** The role of nitrogen, amino acids and phenols in resistance of tomato to root-knot nematodes Nemat. Mediterr. 8(1): 51-57.
- Netscher, C. and R.A. Sikora (1990).** Nematode parasites of vegetables, pp. 237-283. In : M. Luc, R.A. Sikora and J. Bridge, eds. plant parasitic nematodes in subtropical and tropical Agriculture. C.A.B. International Insti. Parasitology. U.K.
- Oheker, G.B. (1965).** Nature of the root-knot resistance introduced into *Lycopersicon esculentum* by interspecific crosses with *L. Peruvianum*. Diss. Abstr. 25:4870. (C.F. Pl. Breed. Abstr. 36:5246.)
- Okophyi, N.S. (1983).** Polyphenol oxidase isoenzymes in roots of tomatoes resistance to *Meloidogyne incognita*. Helm. Abstr. B. 52 (2): 669.
- Okophyi, N.S. and A.V. Sadykin (1978).** The resistance of tomatoes to the gall nematodes. Pl. Breed. Abstr. 48(7): 7128.
-

- Okophyi, N.S.; N.N. Zaginailo and A.V. Sadykin (1987).** A method of Selection nematode resistant tomato varieties and lines. Pl. Breed. Abstr. 57(4): 3324.
- Oostenbrink, M. (1966).** Major characteristics of the relation between nematodes and plants. Meded. Landbouwhoges. Wageningen 66-4. 46 pp.
- Oteifa B.A. (1964).** A taxonomic guide to the common genera of soil and plant nematodes with a supplement of current know economic parasitic species of U.A.R. Contribution of Nat. Res. Cent. Nematol. Unit. Giza. 32 PP.
- Oteifa, B.A. and M. EL Gindi (1957).** Studies on root knot nematodes, *Meloidogyne spp.* in Giza, Egypt. I. Morphology of the root-knot nematode. *Meloidogyne javanica*. Fac. of Agric. Bull. No. 100 Cairo Univ.
- Pchelintseva, M.T. (1979).** Evaluation of a collection of tomato varieties for resistance to root-knot nematodes in the field in Tajikistan. Dushanbe Tajik SSR 45-46. (CF. Helm. Abstr. 50(1): 198, 1981)
- Potenza, C.L.; S.H. Thomas; E.A. Higgins and C. Sengupta-Gpaian (1996).** Early root reponse to *Meloidogyna incognita* in resistant and susceptible Alfalf cultivars. Journal of nematology 28 (1): 175-181
- Pujari, C.V.; R.S. Wagh and P.N. Kale (1995).** Gentie variability and heritability in tomato. Journal of Maharshtra Agricultural Universities 20(1) 15 -171. (C.F. Pl. Breed. Abstr. 1997 67(3) 356.)
- Rai, N.; M.M. Syamal; A.K. Joshi and A.K. Rajput (1997).** Genetics of yield and yield components in tomato *Lycopersicon esculentum* miller; Indian journal of agricultural Research 31(1) 1/6-50 (C.F. Pl. Breed. Abstr (1998) 68(4): 542.
- Raijadhav, S.B and P.N. Kale (1986).** Heterosis in root-knot nematode resistant tomato lines. Pl. Breed. Abstr. 56(5): 4359.
-

- Raijadhav, S.B.; K. G. Choudhari and P.N. Kale (1997).** Correlation studies in tomato. *Journal of Maharashtra agricultural Universities.* 21(2) 252-253. (C.F. Pl. Breed. Abstr. (1998) 68 (5): 682.
- Rajkumar, P. and K. Krishnappa (1986).** Response of tomato cultivars to three races of root-knot nematode *Meloidogyne incognita* Helm. Abstr. B 55(1): 199.
- Riggs, R.D. (1960).** Studies on resistance in tomato to root-knot nematodes. Pl. Breed. Abstr. 36:2057.
- Riggs, R.D. and N.N. Winstead (1959).** Studies on resistance in tomato to root-knot nematode and occurrence of Pathogenic Biotypes Phytophology Vol (49) 716-724
- Roberts, P.A. and J. Thomason (1987).** Variability in reproduction of isolates of *M. incognita* and *M. Javanica* on resistant tomato genotypes. Pl. Dis. 70 (6): 547-551. (C.F. Pl. Breed. Abstr. 57(2): 1560).
- Rohini. H.M.; K. Ekanayake and M. Di. Vito (1984).** Response of lines and varieties of tomato to root-knot nematode infection. Nemat. Medditerr. 12(2) 207-212. (C.F. Pl. Breed. Abstr. 55(7): 5745, 1985.).
- Sadykin, A.V. and N.S. Okophyi (1985).** Rapid method of evaluating the nematode resistance of tomato. Pl. Breed. Abstr. 55(12) 10029.
- Sasser, J.N. (1977).** World wide dissemination and importance of the root-knot nematodes *Meloidogyne* spp. J.Nematol. 9: 26-29.
- Sasser, J.N.; C.C. Carter and K.M. Hartman (1984).** Standardization of host suitability studies and reporting of resistance to root-knot nematodes. A coop pub. Dept. PL. Path., North carolina state Univ. and US agency for Inter Development Raleigh, Nc., U.S.A
-

- Sidhu, G. and J.M. Wedster (1973).** Identification of a resistance gene in tomato *Lycopersicon esculentum* against root-knot nematode. *Meloidogyne incognita*. Can. J. Genet. Cytol. 15 (3) : 663.
- Sikora, R.A.; K. Sitaramaiah and R.S. Singh (1973).** Reaction of root-knot nematod resistant tomato cultivars to *Meloidogyne javanica* in India. PL. Dis. Rept. 57(2): 141-143.
- Singh, B. and B. Choudhury (1973).** The Chemical characteristics of tomato cultivars resistant to rooot knot nematodes *Meloidogyne spp.* Nematol. 19(4):443-448.
- Singh, B.; M.K. Baner Jee and K. Singh (1974).** Inheritance of resistance to root-knot nematodes in tomato. SABRAO. Journal 6(1): 75-78. (C.F. Pl. Breed. Abstr. 44:8228).
- Singh. A.; P.K. Singh; J. Dixtt and J.P.S. Gautam (1995).** Heterosis and inbreeding depression in tomato. Horitcultural Journal. (C.F. Pl. Breed. Abstr. 1996).
- Snell, F.D. and C.T. Snell (1953).** Clorimetric methods of Analysis including some turbidimetric methods. D. van. Nostrand compony, Inc. Tornto-New York. London vol III orgnic-1, pp. 606.
- Tanveer, M. and A.T. Saad (1971).** Reaction of some cultivated crops to two species of root-knot nematodes. Pl. Dis. Rcpt. 55(12): 1082-1084.
- Taylor, D.P. (1975).** Observations on a resistant and a susceptible variety of tomato in a field heavily infested with *Meloidogyne* in senegal. Cahievs Orstom. Seria Biologic (France) 10 (3): 239-245. (C.F. Abstr. On trop. Agtic. 2(6): 09001, 1976).
- Taylor, A.L. and J.N. Sasser (1978).** Biology, identification and control of root-knot nematodes (*Meloidogyne spp.*) Coob Pub. Dep. Plant Pathol., North Carolina State Univ., and U.S. Agency Int. Dev. Raleigh, N.C. III PP.

- Thomason, J. and P.G. Smith (1957).** Resistance in tomato to *Meloidogyne javanica* and *M. incognita acrita*. PL. Dis. Rept. 41:180-181.
- Uppal, G.S.; Tarsem Lal and D.S. Cheema (1997).** Performance of tomato hybrids with regard to yield and quality characters. Journal of research, punjab Agricultural University 34(1) 45-56. (C.F. PL. Breed. Abstr.(1998) 68(6): 842).
- Veech, J.A. (1981).** Plant resistance to nematode pp. 377-403. In: B.M. Zeeckerman and R.A. Rohde, eds., Plant prastic Nematode Vol.(3) Academic press.
- Vito, M. di.; F. Saccard; A. Carella and N.La. Gioia (1986).** New lines of tomato resistant to root-knot nematodes (*Meloidogyne spp.*). Pl. Breed. Abstr. 56(12): 11158.
- Walters, S.; C. Wehner, and R.Barker (1995).** Inheritance of resistance to root-knot nematode (*Meloidogyne arnaria*) in cucumber (*cucumis stivus*). Abstract Hortscience, Vol. 30(4).
- Watts, V. H. (1947).** The use of *Lycopersicon peruvianum* as a source of nematode resistance in tomatoes. Proc. Amer. Soc. Hort. Sci. 49:233-234.
- Williamson, V.N; K.N. Lambert and J.Y. Ho (1993).** Root-knot nematode resistance in tomato. Molecular Biology 349/M57.
- Wilski, A.; J. Giebel and J. Glowinkowska (1971).** Phenolic compounds in roots of tomatoes susceptible and resistant to *Meloidogyne spp.* proce Naukowe Instytutu Ochrony Roslin, 13(1): 155-162.
- Zacheo. G.; G. Pricolo and T. Bleve-Zacheo (1988).** Effect of temperature on resistance and biochemical change in tomato inoculated with *Meloidogyne incognita*. Nemat. Mediterr. 16(1):107-112. (C.F. Pl. Breed. Abstr. 59(3): 2348,1989).