

REFERENCES

Abdel-Al, Z. E. ; Khalf-Allah, A.M. and Shenouda, G.S.(1973): Effect of visual selection and inbreeding on some quantitative characters of summer squash. Alexandria J. Agric. Res. 21 : 277-282 .

Abd El-hafez, A.A. ; Shehata, S. A. S ; Baha El-Din, S.A. ; El-Doweny , H. H. and Awad, M.M.W. (1990) : Inheritance of downy mildew disease and its nature of resistance in cucumber .

Annals of Agric. Sci. Moshtohor 28 (5) 1081-1097 .

Abdel Migid, A.A. (1984) : Study on interspecific hybridization in squashes (*Cucurbita Sp.*) . Ms.C. Thesis, Fac. of Agric. Moshtohor Zagazig Univ. Benha branch.

Al Diab, M.A.G. and Kasrawi, M. A. (1996) : Breeding behavior of vine habit of summer squash (*Cucurbita pepo*) . Dirasat . series B, Pure and Applied Sci. 23 (1) 1-7 [En, ar,16 ref] . (C. F. Plant breed. Abstr., 67 (7) 7340 : 1997) .

Amaral-Junior, A. T. ; Casali, V. W. D. ; Craz, C. D. ; Silva, D. J. H. da; Silva, L. F. C. da; Da-Silva , D. J. H. and Da-Silvag, L. F. C. (1994) : Estimates of phenotypic , genotypic and environmental correlations between seven morphological and agronomic characters in eight pumpkin accessions . Bragantia 53 (2) 163 – 166 (C. F. data base of Agricola-internet) .

A. O. A. C. (1970) : Official Methods of Analysis of the Association of Official Agriculture Chemists.11 th Ed.,Washington,D. C., U. S. A.

- Arima, H. K. and Rodriguez-Amaya, D. B. (1988) :** Carotenoid composition and vitamin A value of commercial Brazilian squashes and pumpkins. *J. Micronut. Anal.* (4) : 177 – 191 .(C. F. data base of Agricola-internet).
- Bhatt, G. M. (1971) :** Heterotic performance and combining ability in a diallel cross among spring wheats (*T. aestivum* L.). *Austr. J. of Agric. Res.* 22 : 359-369 .
- Borchers, E. A. and Taylor, R. T. (1988) :** Inheritance of fruit Bitterness in a cross of *Cucurbita mixta* × *C. pepo* . *Hor. Sci.* 23 (3) 603 – 604.
- Borghi, B. ; Maggidre , T. ; Bogging, G. and Bonali, F. (1973) :** inbreeding depression and heterosis in *Cucurbita pepo* evaluated by of diallel analysis . *Genetica . Agrouria* 27 : 415 – 432 (C. F. Plant Breed. Abstr. 45 – 1634 , 1975) .
- Branchard, M.; Chateau, M.; Megnegneau, B. and Debeaujon, I. (1988) :** Somatic embryogenesis and plant regeneration from cotyledon callus culture of *Cucumis melo* . In : INRA(Ed) Cucurbitaceae 88, Proceedings of the Eucarpia meeting on Cucurbit Genetics and Breeding, Avignon – Montfaret, France, May 31 – June 2, 1988 Paris, France (pp 133-136) .
- Carle, R. B. (1997) :** Bisex sterility governed by a single recessive gene in *Cucurbita pepo* , L. Report – Cucurbita .Genetics cooperative 20 :46-47 [En, 2 ref] .
- Castetter, E. F (1930) :** Species crosses in the genus *Cucurbita* . *Amer . J. Bot .* , 17:41-57 .

- Chee, P.P. (1990) :** High frequency of somatic embryogenesis and recovery of fertile cucumber Plants . Hor. Sci., 25:792-793 .
- Chee, P.P. (1991) :** Somatic embryogenesis and plant regeneration of squash *Cucurbita pepo* L. Cv. Yc 60 . Plant Cell Reports. 9 (11) 620-622.
- Chee , P. P. (1992) :** Initiation and Maturation of somatic embryos of squash *Cucurbita pepo*. Hor. Sci. 27 (1) 59 – 60 .
- Chrambrach, A. and Rodbard, D. (1971) :** Polyacrylamide gel electrophoresis . Science. 172 : 449 – 451.
- Cohen, R. ; Leibovich, G. and Shtienberg , D. P. H. (1993) :** Variability in the reaction of squash (*Cucurbita pepo*) to inoculation with *Sphaerotheca fuliginea* and methodology of breeding for resistance . Plant Pathology 42 (4) 510 – 516 .
- Culpepper, C. W. and Moon, H.H. (1945):** Differences in the composition of the fruits of *Cucurbita* varieties at different ages in relation to culinary Use. J. Agric. Res. 71: 111-136 .
- Cutler, H. C. and Whitaker, T.W. (1956):** *Cucurbita mixta* , its classification and relationships . Bull. Torrey . Bot. Cl . 83: 253-260 .(C. F. Whitaker, T. W. and Davis, G. N. 1962. Cucurbits, Interscience Pub., INC. New York).
- Damarany, A.M.; Aboul Nasr, M.H. and Abdalla, M.M.A. (1995) :** Yield and yield components of some *Cucurbita Spp.* Cultivars and hybrids under Assiut conditions . 1. Summer squash (*Cucurbita Spp.L.*) . Assiut J. Agri. Sci. 26 : 151-58.

- Davis, M.J.; Boker, R. and Hanan, J.J. (1979) :** Colonal multiplication of cornation by micro propagation . J. Amer. Soc. Hort. Sci . 102 : 48-53 .
- Dematte, M. E. S. P. ; Camargo, L. S. ; Alves, S. ; Scaranati, H. J. ; Pettinell, A. ; Veiga, A. A.; Rocha, T. R. and Nagal, V. (1975) :** Study of progenies of varieties of *C. moschata* Duchesne . 618 Facultade de Medicine, veterinariae Agronomia, Jaboticabal , Brazil (C.F. Plant . Breed. Abstr. 46 : 3861 , 1976) .
- De vaulx, R. D. and Pitrat, M. (1985) :** Interspecific cross between *Cucurbita pepo* and *C. martinizii* . Report, Cucurbit , Genetics, Cooperative . 8 : 35 .
- Devi, D. S. ; Balakrishnan, R. ; Paulas, D. ; Subbiah, R. and Natarajan , S. (1989) :** Evaluation studies in pumpkin (*Cucurbita moschata* Poir) . South . Indian Horti. 37 (5) 274 – 276.(C. F. Plant Breed. Abstr. 62 (1) 684 , 1992) .
- Dhingra, O. D., and J. B. Sinclair. (1985) :** Basic Plant Pathology Methods. CRC press, Inc. U. S. A. PP. 355.
- Doijode, S. D. and Sulladmath, U. V. (1981) :** Genetics of heterosis and inbreeding depression for certain quantitative characters in pumpkin (*C. moschata* Poir) Egypt . J. Genet. Cytol. 11 : 135 – 141.

- Dubois, M., Gilles, K. A, Hamilton, J. K., Robers , P. K. and Smith, F. (1956):** Colorimetric method for determination of sugars and related substances. *Analyt. Chem.* 28 : 350 .
- Earle, E. D. and Langhams, R. W. (1975) :** Carnation propagation from shoot tip cultured in liqued medium. *Hortscienc* 10 : 608 – 610 .
- El-Murabaa, , A. and Rushdi, M. (1966) :** The development of summer squash strains resistant to powdery mildew . *Alex. J. Agric. Res.* 14 : 23 – 32 .
- Erwin, A. T. and Haber, E. S. (1929) :** Species and varietal crosses in cucurbits . *Bull. Iowa. Agric. Exp. Sta.* 263 : 344 – 372 .
- Evans, D. A. ; Sharp, W. P. and Flirk, L. (1981) :** In plant tissue culture methods and application in agriculture. Thorpe T. A. (ed.), Academic press. New York , P. 45 .
- Fang, X. ; Yin, Y. ; Xu, H. and Gu, X. (1994) :** Breeding the new cucumber cultivar zhong Nong. *China Vegetables* . 6 : 1 – 3 (C. F. Plant Breed. Abstr. 66 (3) 1902 : 1996) .
- Francis, F. J. (1962) :** Relationship between flesh color and pigment content in squash. *Pro . Amer . Soc. Hort . Sci.* 18 : 408 – 414 .
- Gill, H. S.; Singh, J. P. and Ron, S. (1972) :** Studies on heterosis in summer squash (*C. pepo*) . *Progressive Hort.* 3 : 5-15 .
- Gomez, K. A. and Gomez, A. A. (1984) :** Statistical procedures for agricultural research . 2nd ed. John Wiely and Sons New York .

- Hassan, A. M. ; Sherif, H. S; Nazeem, H. R. and Abdel Migid, A. A. (1984) :** Genetic behaviour of some economic characters in squashes (*Cucurbita sp*) . Annals of Agric. Sc. Moshtohor 22 (1) 175-187 .
- Hayase, H. (1950) :** *Cucurbita* crosses $\times$ 1 : Cytogenetical studies on interspecific hybrids of *C. maxima* $\times$ *C. pepo* in Japanese . Res. Bull. Hokkaido Nat. Agric. Sit. 72 : 48-58 . (C. F. Whitaker, T. W. and Davis, G. N. 1962. Cucurbits, Interscience Pub., INC. New York) .
- Hayase, H. (1957):** Studies on *Cucurbita* crosses XI : cytogenetical studies on interspecific hybrids of *C. maxima* $\times$ *C. pepo* in Japanese. Res Bull. Hokkaido Natl. Agric. Sta., 70 , 15-29 (C. F. Whitaker, T. W. and Davis, G. N. 1962. Cucurbits, Interscience Pub., INC. New York).
- Helal, R. M. ; Zaki , M. S. and Fadl, F. A. (1978) :** Physiological studies on the nature of resistance to powdery mildew in cucumber. Res. Bull., Ain Shams Univ. Cairo Egypt . No. 953 , 12 pp.
- Helal, R. M. (1999):** Production and breeding of *Cucurbits* . In Arabic. Academic Bookshop Pub. Cairo, Egypt 283 pp.
- Holmes, A. D. and Spelman, A. F. (1946) :** Composition of squashes after winter storage. Food Res. 11 : 345-350 .
- Hong, K. H. ; Om, Y. H. and Park, H. G. (1994) :** Interspecific hybridization between *Cucurbita pepo* and *C. moschata* through ovule culture. J. of the korean society for Horti. Sci. 35 (5) 438-448 .(C. F. Plant-Breed. Abstr. 65 (6) 6399 , 1995).

- Jindal, S. ; Arora, Y. K. and Bajaj, K. L. (1979) :** Studies on the phenols of resistant and susceptible varieties of muskmelon in relation to powdery mildew disease. *Plant Biochemical. J.* 6 (1) 48-53 .
- Juretic, B. and Jelaska, S. (1991) :** Plant development in long – term embryogenic callus lines of *Cucurbita pepo* . *Plant Cell Rep.* 9 : 623-626.
- Kasrawi, M. A. (1994) :** Heterosis and reciprocal differences for quantitative traits in summer squash (*Cucurbita pepo* L.). *J. of Genetics and Breeding* 48 (4) 399-403 .
- Kitat, F. M. (1959) :** Colohiploidy in *Citrus auratifilia* , Swingle, C. *linetta* , Russo and *C. sinensis* , osbeck. Ph. D. Thesis Fac. Of Agric. Alex. Univ. pp. 127 .
- Kristkova, E. and Lebeda, A. (1997) :** Possibilities of using wild *Cucurbita species* in breeding via interspecific hybridization
Moznosti Vyuziti Planev rostoucich druhu rodu cucurbita ve slechteni prostrednictvim mezidruhove hybridizace zahradnictv 24 (3) 113-120 (*C. F. Plant Breed. Abstr.* 68 (6) 5969 , 1998).
- Kubicki, B. and Walczak, B. (1976) :** Variation and heritability of beta carotene content in some cultivars of the *Cucurbita species*. *Genet. Pol.* 17 : 531-544 .

- Kwack, S. N. and Fujieda, K. (1985) :** Breeding high female lines through interspecific hybridization of cucurbita . Report, Cucurbit – Genetics – Cooperative – USA No. 8 : 78-79 .
- Kwack, S. N. and Fujieda, K. (1987) :** Seed abortion and techniques for obtaining hybrids in interspecific crosses of *Cucurbita* . J. of the japanese – society for –Horti. Sci. 55 (4) 455-460 .
- Kwack , S. N. and Fujieda, K. (1988a) :** Growth habit in populations of an interspecific cross *Cucurbita pepo* × *C. moschata*. J. of the Korean Society for Hori. Sci. 29 (1) 9-12 . (C. F. Plant Breed. Abstr. 59 (7) 4258 , 1989).
- Kwack, S. N. and Fujieda , K. (1988b) :** Somatic embryogenesis in cultured unfertilized ovules of *Cucurbita moschata*. J. Jpn. Soc. Hort. Sci. 57 (1) 34-42 .
- Lebeda, A. and Kristkova, E. (1994) :** Field resistance of *Cucurbita species* to powdery mildew (*Erysiphe cichorocearum*) . Zeitschrift fur pflanzenkankheiten eiten und pflanzenschutz. 101 (6) 598 – 603 (C. F. Plant Pathology. 74 (7) 4379 , 1995).
- Lebeda, A. and Kristkova, E. (1996) :** Genotypic variation in field resistance of *Cucurbita pepo* cultivars to powdery mildew (*Erysiphe cichoracearum*) . Genetic Resources and Crop Evolution . 43 (1) 79-84 .
- Lee, C. Y. ; Smith, N. L. and Robinson, R. W. (1984) :** Carotenoids and vitamin A. value of fresh and canned winter squashes . Nutr Rep. Int. 29 : 129-133 .

- Leibovich, G.; Elkind, Y.; Dinoor, A. and Cohen, R. (1995) :** Quantitative genetic analysis of *sphaerotheca fuliginea* sporulation in *Cucurbita pepo*. Plant Breed. 114 (5) 460-462 .
- Lewis, E. P. and Merrow, S. B. (1962) :** Influence on the estimation of B- carotene by other carotenoids in butternut squashes at harvest and during storage .J. Agric. Food chem. 10 : 53-56 .
- Malepszy, S. and Nadolska - Orczy, K. A. (1983) :** In vitro culture of *cucumis sativus* , I . Regeneration of plantlets from callus formed by leaf explants Z. pflanzenphysiol . Bd. 111 (5) 273-276 .(C. F. Plant Breed. Abstr. 53 (9) 6998 , 1983).
- Merghany., M.M. (1989):** Genetics and nature of resistance to powdery mildew in interspecific hybrids in *Cucumis melo* . Ph. D. Thesis, Cairo Univ. Fac. of Agric., 522 .
- Metwally, E. I. and El-Fadly, G.A. (1991):** In vitro differentiation of the immature interspecific embryos derived from the crosses between *Cucurbita pepo* and *Cucurbita ficifolia* 4th N1. Conf. of pests and dis. of Veg. and fruits in Egypt . PP : 598-606 .
- Metwally, E. I. ; Haroun, S.A. and El-Fadly, G.A (1996):** Interspecific cross between *Cucurbita pepo* L. and *Cucurbita martinii* through in vitro embryo culture. Euphytica 90: 1-7 .
- Munger, H.M.(1990):** Availability and use of interspecific populations involving *Cucurbita moschata* and *C. pepo*. Report Cucurbit Genetics Cooperative. 13,49 .

- Murashige, T. (1974):** plant propagation through tissue culture . Ann. Reve . Plant Physiol . 25: 135-166 .
- Murashige; T. and Skoog, F.(1962):** Arevised medium for rapid growth and bioassays with tobacco tissue cultures . Physiol . Plant . 15: 473-497 .
- Nesmith, D. S. and Hoogenboom, G. (1994) :** Variation in the onset of flowering of summer squash as a function of days and heat units . J. of the Amer. Soci. For Horti. Scie. 119 (2) 249-252 .
- Noel, M.; Comeau, D. and Lence, P. (1992) :** Embryogenese somatique de la courgette (*Cucurbita pepo* L.) a partir de cotyledons . In. Comptes -rendus du 1er forum jeunes chercheurs de l, IAPTC - france - Du gene a fentreprise' - Amiens, 22-23 Juin 1992 (pp 14-18) . (C. F. data base of Agricola-internet).
- Pangalo, K. I. and Goldhausen, M. K. (1939) :** Interspecific hybridization in the genus *Cucurbita* . C. R. (Doklady) Acad. Sci. U. S. S. R., 24, : 61-64 .
- Phillips, D. J. (1986) :** Carnation shoot tip culture . Tech. Bull. Colorado State University, 102 .
- Pinto, B. G. and Wall, J. R. (1972) :** Genetic control of leucine aminopeptidase and esterase isozymes in the interspecific cross *cucurbita lundelliana* × *C. moschata* . Virginia. J. of. Sci. 23 : (3) 115 .

- Plader, W. and Rakoczy-Trojanwska ,M. (1994) :** Obtaining of hybrids within the family cucurbitaceae by in vitro culture of immature embryos . 111. Characteristics of hybrids of *Cucurbita maxima* × *C. ficifolia* and *C. maxima* × *C. foetidissima* . *Genetica Polonica* . 35 : ½ , 11 – 22 .
- Qi, X. Q. ; Dong, H. R. and Hu, S. L. (1989) :** Electrophoretic analysis of peroxidase and isoesterase of three main cultivated species in the genus cucurbita . *Acta. Horti. Sinica* 16 (4) 299-304 .
- Rakoczy-Trojanowska, M. and Malepszy, S. (1989) :** A method for increased plant regeneration from immature F₁ and Bc₁ embryos of *Cucurbita maxima* Duch. × *C. pepo* L. hybrids . *Plant Cell, Tiss and Organ Culture* 18 : 191-194 .
- Reuveni, R. (1983) :** Resistance reaction of *Cucumis melo* to inoculation with *pseudoperonospora cubensis*. *Ann. Appl. Biol.* 102 : 533-537.
- Rhodes, A. M. (1964) :** Inheritance of powdery mildew resistance in the genus cucurbita. *Plant Dis. Res.* 48 : 54-55 .
- Robinson, R. W. ; Boettger, M. A. and Shail, J. W. (1978) :** Gynoecious sex expression in cucurbita resulting from an interspecific cross. *Cucurbit Genetics Cooperative* . 1 : 31-32 .
- Robinson, R. W.; Munger, H. M. ; Whitaker, T. W.; and Bohn, G. W. (1976) :** Genes of the cucurbitaceae. *Hortscience* 11 : 554-568 .
- Sayed-Ahmed, S. M. K. (1996) :** Regeneration and gene transfer in squash. M. Sc. Thesis Fac. of Agric Cairo . Univ., 95 pp.

- Schloms, M. (1958) :** Embryologische untersuchung auf frage unterschiedlichen Entwicklung reziproker Bastardembryonen aus Kreuzungen Zwischen Riesenkurbis (*C. maxima*) und Gartenkiirbis (*C. pepo* L.) .Flord 146 ; 586-651 (C. F. Whitaker, T. W. and Davis, G. N. 1962. Cucurbits, Interscience Pub., INC. New York)
- Seo, Y. K. ; Paek, K. Y.; Hwang, J. K. and Cho, Y. H. (1991) :** Micropropagation of rootstock and production of interspecific hybrids by embryo culture in *cucurbita* . J. of the korean. Society for . Horti. Sci. 32 (3) 137-145 (C. F. Plant Breed. Abstr. 63 : 6654, 1993) .
- Sharp, W. R. ; Sandable , M. R. and Maraffu, S. B. (1980) :** The physiology of in vitro a sexual embryogenesis . Hortic. Rev. 2 : 268-310.
- Shelepov, V. V. and Kulchitskaya, V. A. (1970) :** Detection of the effect of heterosis in the forage pumpkin. Shelektsiya i semenovodstvo. Resp mezhved tomat nauchn sb. No 16 , 87-90 .(C. F. Plant Breed. Abstr. 44 : 1176 , 1974).
- Smith, H. H. (1952) :** Fixing transgressive vigor in *Nicotiana rustica* . In Heterosis, Iowa state college Press. Amer. Iowa, U.S.A. .
- Smith, M. K. and Drew, R. A. (1990) :** Current applications of tissue culture in plant propagation and improvement. Australian. J. of plant. Physiology. 17 (3) 267-289 .

- Snell, F. D. and Snell, C. T. (1953) : Colorimetric methods of analysis. including some turbidimetric methods D. Van Nostrand company, Inc. Toronto- New York . London vol. 111 organic 1, pp. 606.
- Steel, R. G. D. and J. H. Torrie (1980) : Principles and procedures of statistics . 2nd Ed. Mc Graw-Hill Inc., New York. 633 P.
- Takashima, S. (1954) : The growth rate of pollen tubes in interspecific hybrids of *Cucurbita*. 11 : The growth rate of pollen tubes in intraspecific matings in Japanese . Jap. J. Genet. 29, 36-39 .
- Thambural, S. and Kamalantham, S. (1973) : Sex expression in *Cucurbita moschata* pair and *Benincasa hispida* cogn . Madras Agri. J. 60 (9/12) 1673-1680 (C. F. Plant Breed. Abstr. 43 (8) 5011, 1974) .
- Tronickova, E. (1971) : The courgette Diamant. Zaharad. Listy. 64 (9) 271-272 (C. F. Plant. Breed. Abstr., 43 (9) 7413, 1973) .
- Umiel, N. and Gobelmoii; W. H. (1971) : Analytical procedure for detecting carotenoides of carrot (*Daucus carota* L.) roots and tomato (*Lycopersicon esculentum*) fruits. J. Amer. Soc. Hor. Sci. 96 , 702-704 .
- Wall, J.R. (1954): Interspecific hybrids of cucurbita obtained by embryo culture . Proc.Amer. Soc. Hort. Sci. 63,427-430 .
- Wall, J. R. and Whitakr, T.W. (1971): Genetic control of leucin aminopeptidase and esterase iszymes in the interspecific cross *Cucurbita ecuadorensis* × *C. maxima* . Biochem . Genet . 5: 223-229 .

- Weeden, N. F. and Robinson, R.W. (1986): Allozyme segregation ratios in the interspecific cross *Cucurbita maxima* × *Cucurbita ecuadorensis* suggest that hybrid breakdown is not caused by minor alterations in chromosome structure. *Genetics. Society of America* 144: 593-609.
- Weilung, F. (1955): Über die interspezifische Kreuzbarkeit verschiedener Kürbisarten. *Zeitschrift für Pflanzenzüchtung* 25 : 33-57 (C. F. Whitaker, T. W. and Davis, G. N. 1962. *Cucurbits*, Interscience Pub., INC. New York).
- Whitaker, T. W. and Davis, G.N (1962) : *Cucurbits* (Botany, Cultivation and utilization). Leonard Hill, London and Interscience publ, New York (Worlds crops Books).
- Yamane, Y. (1952): Studies on species hybrids in the genus *Cucurbita*. I, *C. maxima* × *C. moschata* and its reciprocal cross. Rep. Kihara Inst. Biol. Res. (Seikenziho), 5, 49-99 (C. F. Whitaker, T. W. and Davis, G. N. 1962. *Cucurbits*, Interscience Pub., INC. New York).
- Yamane, Y. (1953): Studies on species hybrids in the genus *cucurbita*. III, F₁ hybrids of *C. moschata* × *C. Pepo* with special reference to the varieties Kogi Ku (*C. moschata*) and Somen (*C. pepo*). Biol. J. Okayama Uni. 0. 1 : 202-208. (C. F. Whitaker, T. W. and Davis, G. N. 1962. *Cucurbits*, Interscience Pub., INC. New York).
- Ziv, M. and Godasi, G. (1986): Enhanced embryogenesis and plant regeneration from cucumber (*Cucumis sativus* L.) callus by activated charcoal in solid/ Liquid double layer cultures. *Plant Sci.* 47: 115-122.