

REFERENCES

- A.A.C.C. 1962.
Approved Methods of American Association of Cereal Chemists. 7th Ed.
The American Association, St.. Paul Minn.
(C.F. Egypt. J. Food Sci., 10 (1-2): 3 1982).
- Abd- El- Gawad, H.F. 1968.
Microbiological and biochemical studies on the deterioration of rice bran.
Ph.D. Thesis, Fac. of Agric., Ain Shams Univ.
- Aly, M.M. 1972.
Deterioration of stored wheat grains by some fungi.
M.Sc. Thesis, Fac. of Agric., Ain Shams Univ.
- Anonymous 1949.
Report of the senior plant pathologist.
Egyptian Ministry of Agriculture (in Arabic)
(C.F. S.M. Fathi, Deterioration of corn grain in storage by fungi, Ph.D. Thesis, Fac. of Agric., Ain Shams Univ.).
- 1983.
The Agricultural Economic Report.
Ministry of Agriculture, U.A.R. (in Arabic).
- A.O.A.C. 1960.
Official methods of analysis, 7th Ed.
The Association, Washington, D.C. p. 247.
- 1965.
Official Methods of Analysis. Association
of Official Agricultural Chemists 10 Ed. p. 584.

A.O.A.C. 1975.

Association of Official Analytical Chemists
Official Methods of Analysis, 12th Ed.
The Association, Washington, D.C. p. 413.

-----, 1980.

Association of Official Analytical Chemists
Official Methods of Analysis. "Determination
of Crude Fat ". 14th Ed.
The Association, Washington, D.C., p. 376.

-----, 1980.

Association of Official Analytical Chemists
Official Methods of Analysis. "Determination
of sugars". 14th Ed.
The Association, Washington, D.C., p. 516.

Assawah, M.W., and H. Elarosi 1960.

Fungi associated with wheat, barley and maize grains.
Egyptian, J. Botany, 3: 153- 164.

Azab, Y. and M. Fahim 1960.

Apreliminary report on the effect of corn grains
treatment.
Agric. Science Magazine, (in Arabic), 13: 93- 99.

Barnett, H.L. 1955.

Illustrated Genera of Imperfect Fungi.
Burgess Publishing Company.

Barron, G.L., and R.W., Lichwardt 1959.

Quantitative estimation of fungi associated with
deterioration of stored corn in Iowa.
Iowa St. Coll. J. Sci., 34: 147- 155.
(C.F. S.M. Fathy, Deterioration of corn grain in storage
by fungi, Ph.D. Thesis, Fac. of Agric. Ain Shams Univ.).

Baruah, P., and H.K., Barush 1978.

Polyphenoloxidase activity of certain fusaria during infection.

Iranian Journal of Plant Pathology, 14 (1-4): 3-8.

(C.F. Review of Plant Pathology, 59 (8): 363, 1980).

Bothast, R.J.; R.A., Anderson; K. Warner and W.F., Kwolek 1981.

Effect of moisture and temperature on Microbiological and sensory properties of wheat flour and corn meal during storage.

Cereal Chem., 58: 309- 311.

Bottalico, A. and V., Piglionica 1978.

Presence of zearalenone in maize and wheat in Italy.

Sulla presenza di zearalenone in cariossidi di frumento e di mais in Italy, 59- 72.

(C.F. Review of Plant Pathology, 57 (12) 485, 1978).

Bottomley, R.A.; C.M. Christensen; and W.F., Geddes 1950.

Grain storage studies, 9: The influence of various temperatures, humidities and oxygen concentrations on mold growth and biochemical changes in stored yellow corn.

Cereal Chem., 27: 271- 296.

----- 1952.

Grain storage studies, 10: The influence of aeration, time and moisture content on fat acidity, non-reducing sugars and mold flora of stored yellow corn.

Cereal Chem. 29: 53- 64.

Brodnik, T. 1975.

Influence of toxic products of Fusarium graminearum and Fusarium moniliforme, on maize seed germination and embryo growth.

Seed Science and Technology, 3 (3-4): 691-696.

Cappellini, R.A. and J.L., Peterson 1966.

Production, in vitro, of certain pectolytic and cellulolytic enzymes by fungi associated with corn stalk rot.

Bull. Torrey Bot., Club, 93 (1): 52- 55.

(C.F. Rev. Appl. Mycol., 45 (8): 375, 1966).

Cheremisinov, N.A. 1959.

The range of causal agents of diseases and the routes for fungal infection of maize seed (Zea mays L.).

Bot. Zh. S.S.S.R., 44 (71): 916- 928.

(C.F. Rev. Appl. Mycol., 39 (4): 427, 1960).

Christensen, C.M. 1957.

Deterioration of stored grains by fungi.

Botan. Rev., 23: 108- 134.

----- 1964.

Effect of moisture content and length of storage period upon germination percentage of seeds of corn, wheat and barley free storage fungi.

Phytopathology, 54 : 1464 - 1466.

----- 1967 a.

Some changes in No. 2 corn stored two years at moisture contents of 14.5 and 15.2 and temperatures of 12°, 20° and 25°C.

Cereal Chem. 44: 95- 99.

Christensen, C.M. 1967 b.

Increase in invasion by storage fungi and in fat acidity values of commercial lots of soybeans stored at moisture content of 13.0- 14.0. Phytopathology, 57: 622- 624.

Christensen, C.M. 1969.

Influence of moisture, temperature and time of storage upon invasion of rough rice by storage fungi.

Phytopathology, 59: 145- 148.

----- 1970.

Moisture content, transfer and invasion of stored sorghum seeds by fungi.

Phytopathology, 60: 280- 283.

----- 1972.

Storage tests with popcorn.

Plant Disease Reporter, 56 : 1097- 1099.

----- and C.E., Dorworth 1966.

Influence of moisture content, temperature, and time on invasion of soybeans by storage fungi.

Phytopathology, 56: 412- 418.

----- and D.R., Gordon 1948.

The mold flora of stored wheat and corn and its relation to heating of moist grain.

Cereal Chem., 25: 40- 51.

----- and H.H., Kaufmann 1965.

Deterioration of stored grains by fungi.

A. Rev. Phytopath., 3: 69- 84.

Christensen, C.M., and L.C., Lopez 1965.

Relation of moisture content and length of storage
to changes in the microflora and germination
percentage of rough rice.
Phytopathology, 55: 953- 956.

----- and P., Linko 1963.

Moisture contents of hard red winter wheat as
determined by meters and by oven drying and inf-
luence of small differences in moisture content
upon subsequent deterioration in storage.
Cereal Chem., 40 : 129- 137.

Clayton , C.N. 1942.

The germination of fungus spores in relation to
controlled humidity.
Phytopathology, 32: 921- 943.

Cook, R.J. 1973.

Disease of grain maize.
ADAS Quarterly Review , 11: 113- 117.
(C.F. Review of Plant Pathology, 53: (8): 606, 1974).

Delp, C.J. 1954.

Spore germination (Water and Germination).
Phytopathology, 44: 615- 626.
(C.F. Vincent W. Cochrane, 1958, Physiology of fungi).

D'Ercole, N., M.G., Fantino and A., Canova 1978.

Research on the presence of Fusarium moniliforme
Sheld on maize (Zea mays L.) Kernels.
Istituto di pathologi Vegetale : 47- 57.
(C.F. Review of Plant Pathology, 57 (12): 485, 1978).

Dimitrijevic, M. and D., Jurkovic 1979.

Mycoflora on maize seeds.

(C.F. Review of Plant Pathology, 58 (4): 143, 1979).

Dubinvica, B.N. 1964.

On diseases of maize cobs.

Kukuruz, 9 (8): 56.

(C.F. Rev. Appl. Mycol., 44 (6): 287, 1965).

Fangari, I.M. 1975.

Studies on the late wilt disease of maize.

Further studies on the mechanism of resistance and susceptibility to the disease.

Ph.D. Thesis, Fac. of Agric. Cairo, University.

Fathi, S.M. 1966.

Some studies on seed and seedling diseases of corn.

M.Sc. Thesis, Fac. of Agric., Ain Shams, Univ.

Futrell, M.C. 1973.

New concepts in chemical seed treatment of agronomic.

Crops Biological Abstracts, 55: 27858.

Gangopadhyay, S. and Wyllie, T.D. 1973.

Biochemical comparison of nine soybean varieties in relation to their susceptibility to Macrophomina phaseolina.

Indian J. Mycol. and Pl. Pathol., 3: 131- 140.

(C.F. M.M. Mostafa, studies on some rots of soybean plant, Ph.D. Thesis, Fac. of Agric. Al-Azhar Univ.).

Ghosal, S., D.K., Chakrabarti and R.S., Sriwastava 1979.

Effect of 3 polyphenolic compounds against ear rot of corn incited by Fusarium moniliforme shield.

Experientia, 35 (1): 83- 84.

(C.F. Review of Plant Pathology, 58 232, 1979).

- Golubevuk, M., H., Sorgar-Domenigg, L.S. Cuendet, C.M., Christensen, and W.F., Geddes 1956.
Grain storage studies 19. Influence of mold infestation and temperature on the deterioration of wheat during storage at approximately 12 % moisture.
Cereal Chem. 33: 45- 52.
- Goodman, J.J., and M. Clyde Christensen, 1952.
Grain storage studies. XI Lipolytic activity of fungi isolated from stored corn.
Cereal Chem., 29: 299- 309.
- Goodman, R.N., Z., Kiraly and M.Z., Aitlin 1967.
The biochemistry and physiology of infections plants diseases.
D. Van. Nostrand Company, Inc. Toronto. New York, London. pp: 853- 902.
- Gottlieb, D. 1950.
Biological effects on spore germination.
Boton. Rev. 16: 229- 257.
(C.F. Vincent W. Cochrane, 1958, Physiology of fungi).
- Grisenko, G.V. 1972.
On the forms of parasitism of facultative pathogens on maize.
Nikol. i. Fitopatol., 5 (3): 251- 257.
(C.F. Review of Plant Pathology, 51 (1): 59, 1972).
- Gulya, T.J., C.A., Martinson and L.H., Tiffany 1979.
Ear- rotting fungi associated with opaque- 2 maize.
Plant Dis. Rep., 63 (5): 370- 373.

Hall, C.W., 1957.

Drying Farm Crops.

Agricultural Consulting Assoc., Reynolds Burgohio.
Avi. Publishing Co. Westport. pp: 93-115.

Hansen, H.N. 1926.

A simple method of obtaining single spore culture.
Science, 64: 384, 1959.

Hummal, B.C.W., L.S., Cuendet, C.M. Christensen, and
W.F., Gedder 1954.

Grain storage studies. 13: Comparative changes in
respiration, viability, and chemical composition
of mold-free and mold contaminated wheat upon
storage.

Cereal Chem., 31 : 143- 150.

Ibrahim, I.A. and S.A., Farag 1965.

A study on some fungi isolated from grains of
Egyptian maize varieties.

Alex. J. Agric., Res., 13: 401 - 413.

Karadzhova, I. 1978.

Fusarium species on maize.

Restitelna Zashchita, 26 (10): 28- 32.

(C.F. Review of Plant Pathology, 59 (8): 373, 1980).

Kennedy, B.W. 1964.

Moisture content, mold invasion, and seed viability
of stored soybeans.

Phytopathology, 54: 771- 774.

Koehler, B. 1938.

Fungus growth in shelled corn as affected by
moisture

J. Agric. Res., 56: 291- 307.

Leon-Gallegos, H.M. 1966.

The differential effect of mercuric chloride on growth of certain fungi associated with corn seed. Proc. Indiana Acad. Sci., 76: 217- 220.
(C.F. Rev. Appl. Mycolo., 47 (9): 470, 1968).

Lichtwardt, R.W., G.L., Barron and L.H., Tiffany 1958.

Mold flora associated with shelled corn in Iowa. Iowa St. Coll. J. Sci., 33: 1- 11.
(C.F. S.M.F. Ahmed. Deterioration of corn grains in storage by fungi, Ph.D. Thesis, Fac. of Agric., Ain Shams, Univ.).

Lopez, L.C., and C.M., Christensen 1967.

Effect of moisture content and temperature on invasion of stored corn by Aspergillus flavus. Phytopathology, 57: 588- 590.

Marasas, W.F.O., N.P.J., Kriek, V.M., Viggins, P.S.,

Steyn, D.K., Towers, and T.J., Hastie 1979.
Incidence, geographic distribution, and toxigenicity of Fusarium species in South African Corn. Phytopathology, 69 (11): 1181- 1185.

Medina Candanedc, D.M. and M., Zenteno Zevada 1976.

Mycoflora of corn ears in Mexico II. Biol. Soc. Mex. Micel., 10 : 71- 77.
(C.F. Microbiology Abstracts : Section A, 12 (10) : 60, 1977).

Milatovic ~~Y~~anka, I. 1967.

The causes of maize ear- rot in the territory of Croatia. Poljopr. Znanst. Smotra, 24 (7): 3- 9.
(C.F. Review of Plant Pathology, 50 (12): 667, 1971).

Milovoj-Milharcic, L. and J., Macek 1975.
Moulding of maize ears in Slovenia.
Agronomski Glasnik, 37 (1-4): 185- 190.
(C.F. Review of Plant Pathology, 55 (8): 687, 1976).

Mislivec, P.B., and J., Tuite 1970 .
Species of Penicillium occurring in freshly- har-
vested and in stored dent corn kernels.
Mycologia, 62: 67- 74.

Mohamed, A.A.A. 1968.
Studies on banana fruit rot during development,
harvesting and ripening in A.R.E.
Ph.D. Thesis, Fac. of Agric., Cairo Univ.

Mohamed, H.A., and S.M., Fathi 1965.
Nigrospora cob rot of corn in the United
Arabe Republic.
Pl. Dis. Reptr., 49 (3): 244- 246.

-----, W.E., Ashour, A.R. Sirry, and S.M.,
Fathi 1967.
Fungi carried by corn seed and their importance in
causing corn diseases in the United Arab Republic.
Pl. Dis. Reptr., 51 (1): 53- 56.

----- 1968 a.

Factors affecting severity of seedling blight of
corn in The United Arab Republic.
Plant Dis. Reptr., 52: 79- 83.

----- 1968 b.

Fungi causing seedling blight of corn in The
United Arab Republic.
Plant Dis. Reptr., 52: 84- 86.

Moreno, M.E., and C.M., Christensen 1970.

Effect of moisture and fungi on the viability of stored maize.

Reuta Lat. Am. Microbiol. 12: 115- 121.

(C.F. Review of Plant Pathology, 50 (2): 110, 1971).

-----; G.J.; Ramirez; M., Mendoza; and G.

Valencia 1982.

Effect of fungicides on storage of maize seeds previously attacked by storage fungi.

Turrialba, 32 (2): 97- 101.

(C.F. Review of Plant Pathology, 62 (5): 178, 1983).

-----; and G.V., Vidal 1981.

Preserving the viability of stored maize seed with fungicides.

Plant Disease, 65: 260- 261.

Mostafa, M.M. 1980.

Studies on some rots of soybean plant.

Ph.D. Thesis, Fac. of Agric. Al- Azhar Univ.

Nagel, C.M.; and G., Semeniuk 1947.

Some mold- induced changes in shelled corn.

Pl. Physiol., 22: 20- 33.

Olafson, J.H.; C.M., Christensen and W.F., Geddes 1954.

Grain storage studies. 15: Influence of moisture content, commercial grade, and maturity on the respiration and chemical deterioration of corn.

Cereal Chem., 31: 333 - 340.

Osman, M.A.; M.F., Abdel Rahim and I.M., El-Assiuty 1979.

Late-wilt disease of maize. Part 2: Relation of phenols, sugars, Zinc and Copper to late- wilt disease resistance of maize.

Ann. Agric. Res. Bull., p. 1144, Ain Shams Univ.

Papavizas, C.C.; and C.M., Christensen 1960.

Grain storage studies. 29: Effect of invasion by individual species and mixture of species of *Aspergillus* upon germination and development of discolored germs in wheat.

Cereal Chem., 37 : 197- 203.

Pelhate, J. 1981.

Mycoflora of maize seeds. II- Ear storage.

Cryptogamie Mycologie, 2 (1): 61- 84.

(C.F. Review of Plant Pathology, 61 (1): 17, 1982).

Pencic, Viktorija, 1969.

The effect of *Nigrospora* (*Khuskia*) *oryzae* on the development of maize.

Arh. Poljopr. Nauke Teh., 22 (11): 118- 158.

(C.F. Review of Plant Pathology, 50 (9): 526, 1971).

Piglioni, V.; A., Siniscalco; and F., Gigante 1976.

Heat treatment of maize kernels internally infected by *Fusarium moniliforme*.

Phytopathologia Mediterranea, 15 (2-3): 98-102.

(C.F. Review of Plant Pathology, 57 (1): 20 , 1978).

Popov, A. 1974.

Fusarioses of maize.

Rastitelna Zashchita, 22 (6): 37- 40.

(C.F. Review of Plant Pathology, 54 (8): 614, 1975).

Qasem, S.A., 1959.

Studies on deterioration of stored by fungi.

Dis. Abstr., 20: 1952 - 1953.

----- and C.M., Christensen 1958.

Influence of moisture content, temperature, and time on the deterioration of stored corn by fungi.

Phytopathology, 48: 544- 549.

Qasem, S.A.; and C.M., Christensen 1960.

Influence of various factors on the deterioration
of sotred corn by fungi.

Phytopathology, 50: 703-709.

Radwan, I.A.M.I. 1980.

Studies on storage diseases of gralic in the A.R.E.
Ph.D. Thesis, Fac. of Agric., Zagazig University.

Rohringer, R. and Samborski, D.J. 1967.

Aromatic compounds in host parasite interaction.
Ann. Rev. Phytopathol., 5: 77- 86.

Rubin, B.A. and Artsikhovaskaya, Y.V. 1963.

Biochemistry and physiology of plant immunity.
Ann. Rev. Phytopathol., 4: 349- 364.

Salazar, F.F.; and G.E., Vargas 1977.

Methods of evaluation and levels of resistance in
maize to ear rot caused by Fusarium moniliforme.
Agronomia Costarricense, 1 (2): 93- 99.
(C.F. Review of Plant Pathology, 59 (7): 321, 1980).

Saur , D.B.; and C.M., Christensen 1966.

Comparison of hard red spring and winter wheats
in storage.
Phytopathology, 56: 619- 623.

----- 1968.

Germination percentage, storage fungi isolated
from, and fat acidity values of export corn.
Phytopathology, 58: 1356- 1359.

----- 1969.

Some factors affecting increase in fat acidity
values in corn.
Phytopathology: 59: 108- 110.

Semeniuk, G. 1954.

Microflora, pp. 77- 151. In. J.A. Anderson and A.W. Alcock, (Eds.), Storage of Cereal Grains and Their Products.

An. Assoc. Cereal Chem., St. Paul., Minn.

(C.F. S.M. Fathi, Deterioration of corn grains in storage by fungi, Ph.D. Thesis, Fac. of Agric. Ain Shams Univ.).

Shaw, M. and Colotelo, N. 1961.

The physiology of hostparasite relations:

VII. The effect of stem rust on the nitrogen and amino acids in wheat leaves.

Can. J. Bot., 39: 1351- 1372.

Singh, D. and T., Singh 1977.

Location of Fusarium moniliforme in kernels of maize and disease transmission.

Indian, Journal of Mycology and Plant Pathology;
7 (1): 32- 38.

Sinha, R.N. and W.E., Muir 1973.

Grain storage, part of a system.

The Avi. Publishing Company, Inc. Westport.

(C.F. Review of Plant Pathology, 54 (2): 104, 1975).

----- and H.A.H., Wallace 1966.

Association of granary mites and seed borne fungi in stored grain and outdoor and indoor habitats.

Ann. Entmol. Soc. Am., 59: 1170.

(C.F. Review of Plant Pathology, 45 (12): 621, 1966).

Sirry, A.R.; M.F., Higazi and M.M., Aly 1973 a.

Deterioration of wheat under commercial long storage.

Res. Bull. (31) High Polytechnical Inst. Agric.

Dept. Moshtohor, A. R. E.

Siu, R. G. H., 1951.

Microbial decomposition of cellulose.
New York; Reinhold Publ. Corp. p. 531.

Smiljakovic, H., 1975.

Parasitic lodging of maize.
Agronomski Glasnik, 37 (1-4): 171-176.
(C.F. Review of Plant Pathology, 55 (6):
504, 1976).

-----, and M. Draganic 1977.

Contribution to the study of the aetiology of
root, stalk and ear rot of maize in Serbia.
Zastita Bilja, 28, (139): 97- 106.
(C.F. Review of Plant Pathology, 58 (4):
143, 1979).

Snell, F.D., and C.T., Snell 1953.

Colorimetric methods of analysis including some
turbidimetric and nephelometric methods.
D. Van. Nostrand Company, Inc. Toronto- New York
London. Vol. III. Organic. 1, p. 606.

Sorger- Domenigg, H., L.S., Cuendet; C.M., Christensen
and W.F., Geddes 1955.

Grain storage studies 17: Effect of mold growth
during temporary exposure of wheat to high moisture
contents upon the development of germ damage and
other indices of deterioration during subsequent
storage.
Cereal Chem., 32: 270- 285.

Staples, R.C. and Ledbetter, M.C. 1958.

A study by microautoradiography of the distribution of tritium labeled glycine in rusted pinto bean leaves.

Contrib. Boyce Thompson Inst., 19: 349- 359.

(C.F. Goodman et al. 1967).

Strobel, G.A. and D.E., Mathre 1970.

Outlines of Plant Pathology.

Van Nostrand Reinhold Company N.Y. 10001.

Tanaka, S. and Katsuki, H. 1953.

On the chemical constituents on plant susceptible to the plant disease .

Chem. Soc. Japan, 73: 259.

(C.F. M.M. Mostafa, Studies on some rots of soybean plant. Ph.D. Thesis, Fac. of Agric. Al-Azhar Univ.).

Tolba, M.K.; and A.M., Salama 1958.

Effect of temperature on the absorption and utilization of sucrose by mycelial mats of Fusarium oxysporium.

Egyptian.J., Bot., 1: 81- 91.

Warren, H.L., 1977.

Survival of Colletotrichum graminicola in corn kernels.

Phytopathology, 67 (2): 160- 162.

Welly, R.E.; S.A., Qasem and C.M., Christensen 1963.

Tests of corn stored four years in a commercial bin. Cereal Chem., 40: 277- 282.

- Wilson, D.M.; L.H., Huang and E., Jay 1975.
Survival of Aspergillus flavus and Fusarium moniliforme in high moisture corn stored under modified atmospheres.
Applied Microbiology, 30 (4): 592- 595.
- Windels, C.E.; M.B., Windels and T., Kommedahl 1976.
Association of Fusarium species with picinic Beetles on corn ears.
Phytopathology, 66 (3): 328- 331.
- Zeinab, A. El- Awamry; M.M., Hassen, and A. El-Zainy, Tahany 1980.
Purification and some properties of Fusarium moniliforme allantoinase.
Egypt. J. Bot., 23 (1): 35- 41.
- Zeleny, L. 1954 .
Chemical, physical, and nutritive changes during storage, pp. 46-76. In J.A. Anderson, and A.W. Alcock, (Eds.), Storage of Cereal Grains and Their Products, Am. Assoc. Cereal Chem., St. Paul. Minn.
- , and D.A., Coleman 1938.
Acidity in cereals and cereals products, its determination and significance.
Cereal. Chem., 15: 580- 595.
- Zenteno- Zevada, M.; and M., Ullora Sosa 1977.
Mycoflora of maize ear (Zea mays L.). I.
Rev. Latinoam. Microbiol., 19 (1): 27- 31.
(C.F. Microbiology Abstracts: Section C., 8 (11) : 48, 1979).