

REFERENCES

- Abdel-ware, A. A. (1993): Studies on growth and development of *Tilapia nilotica* as affected by different environmental factors. M.Sc. Thesis, Faculty of Agriculture, Al-Azhar University, Egypt.
- Akiyama, T. and Nose, T. (1981): Changes in body weight, condition factor and body composition of fingerling Chum salmon with various sizes during starvation. Bull. Res. Nat. Inst. Aquacult. 1:71-78.
- Al-Amoudi, M. M., El-Nakkadi, A. M. N. and El-Nouman, B. M. (1992): Evaluation of optimum diet requirement of vitamin C for the growth of *Oreochromis spilurus* fingerlings in water from the Red Sea. Aquaculture, 105:165-173.
- Anadu, D. I. and Ebeonwm, B. I. N. (1994): Application of liquid manure in Common carp production. Aquaculture magazine. 20:56-62.
- Anderson, C. E. and Smitherman, R. O. (1978): Production of normal male and androgen sex/reserved tilapia aurea and T. nilotica fed a commercial catfish diet in ponds, Pp. 34-39. In R. O. Smitherman, W. L. Shelton and J. H. Grover (eds.) culture of exotic fish symposium proceedings. Fish culture section. American Fisheries Society, Auburn, Alabama.
- Anderson, J., Jackson, A. J., Maltby, A. J. and Capper, B. S. (1984): Effect of dietary carbohydrates and fiber on the tilapia (*Oreochromis niloticus*). Aquaculture, 37:303-314.
- A.O.A.C. (1990): Official Methods of Analysis. Association of Official Analytical Chemists. Washington, D. C.
- Arzel, J. Lopez, F. Metailler, R. Stephan, G. Viau, M. Gandemer, G. and Guillaume, J. (1994): Effect of dietary lipid on growth performance and body composition of brown trout (*Salmo trutta*) reared in seawater. Aquaculture, 123:361-375.
- Avella, M., Berhaut, J. and Bornancin, M. (1993): Salinity tolerance of two tropical fishes, *Oreochromis aureus*, and *O. niloticus*. 1. Biochemical and morphological changes in the gill epithelium. J. Fish Biol., 42:243-254.
- Balarin, J. D. and Hatton, J. P. (1979): Tilapia- A guide to their biology and culture in Africa. Institute of Aquaculture, University of Stirling, Stirling, Scotland (Book).
- Ballestrazzi, R. and Lanari, D. (1996): Growth, body composition and nutrient retention efficiency of growing sea bass (*Dicentrarchus labrax* L.) fed fish oil or fatty acid Ca salts. Aquaculture, 139:101-108.
- Barash, H., Plavnik, I. and Moav, R., (1982): Integration of duck and fish farming : Experimental results. Aquaculture, 27:129-140.
-

- Barash, H. and Schroeder, G. L., (1984): Use of fermented cow manure as a feed substrate for fish polyculture in Stagnant water ponds. *Aquaculture*, 36:127-140.
- Bardach, J. E., Ryther, J. H. and McLarney, W. D. (1972): *Aquaculture - The farming and husbandry of freshwater and marine organisms*. Wiley-Interscience, New York.
- Beamish, F. W. H., Hilton, J. W., Niimi, E. and Slinger, S. J. (1986). Dietary carbohydrate and growth, body composition and heat increment in rainbow trout (*Salmo gairdneri*). *Fish. Physiol. Biochem.*, 2:85-91.
- Behrends, L. L., Maddox, J. J., Madewell, C.E. and Pile, R.S. (1980): Comparison of two methods of using liquid swine manure as an organic fertilization in the production of filter feeding fish. *Aquaculture*, 20:147-153.
- Belay, M. (1995): Induzierte Tetraploidie bei Afrikanischen Buntbarschen (*Oreochromis niloticus*). Diplomarbeit, Institut für Tierzucht und Haustiergenetik and der Georg-August-Universität, Göttingen, Germany.
- Benfey, T. J. (1991): The physiology of triploid salmonids in relation to aquaculture. In: V. A. Pepper (Editor), proceedings of Atlantic Canada workshop on methods for the production of non-maturing salmonids : 9-21 February, 1991, Dartmouth, N. S. Canadian Technical Report of fisheries and Aquatic Science No. 1789, pp. 73-80.
- Benfey, T.J. and Sutterlin, A. M. (1984): The haematology of triploid atlantic landlocked salmon *Salmo salar* L. *J. Fish Biol.*, 24:333-338.
- Billard, R. (1995): The major carps and other cyprinids. pp. 21-55 in: C.E. Nash and A.J. Novotny(Editors). *Production of aquatic animals (fishes)*. World Animal science, C8.
- Bitterlich, G. (1985): Digestive processes in Silver carp (*Hypophthalmichthys molitrix*) studied in vitro. *Aquaculture*, 50:123-131.
- Bitterlich, G. and Gnaiger, E. (1984): Phytoplanktivorous or omnivorous fish? Digestibility of zooplankton by silver carp, *Hypophthalmichthys molitrix* (val). *Aquaculture*, 40:261-263.
- Borlongan, I. G. and Benitez, L. V. (1992): Lipid and fatty acid composition of milkfish (*Chanos chanos* Forsskal) grown in freshwater and seawater. *Aquaculture*. 104:79-89.
- Bowen, S. H. (1982): Feeding, digestion and growth-qualitative considerations. Pages 141-156 in R.S.V. Pullin and R. H. Lowe-McConnell (editors). *The biology and culture of tilapias*. International Center of Living Aquatic Resources Management, ICLARM conference proceedings 7, Manila. Philippines.
- Bowman, D. (1977): Comparacion entre *Tilapia aurea* Steindachner *Tilapia mossambica* peters en estanques de El Salvador. *FAO fish Rep.* 159 (1):78-90 (In Spanish with English summary).

- Boyd, C. E. (1979): Water quality in warmwater fishponds. Auburn University Agriculture Experiment station, Auburn, Alabama, pp 359.
- Brämick, U. Puckhaber, B., Langholz, H.-J. and Hörstgen-Schwark, G. (1995): Testing of triploid Tilapia (*Oreochromis niloticus*) under tropical pond conditions. *Aquaculture*, 137:343-353.
- Brown, P. B. and Robinson, E. H. (1989): Comparison of practical catfish feeds containing 26 or 30% protein. *Prog. Fish-Cult.*, 51:149-151.
- Brown, M. L. and Murphy, B. R. (1991): Relationship of relative weight (Wr) to proximate composition of juvenile Striped bass and hybrid Striped bass. *Trans. Am. Fish. Soc.*, 120:509-518.
- Brummett, R. E. and Alon, N. C., (1994): Polyculture of Nile tilapia (*Oreochromis niloticus*) and Australian red claw crayfish (*Cherax quadricarinatus*) in earthen ponds. *Aquaculture*, 122:47-54.
- Buckley, J. T. and Groves, T. D. D. (1979): Influence of feed in the body composition of finfish. In J. E. Halver and K. Tiews (editors), *Fish nutrition and fish feed technology. Proceedings of a symposium, 20-23 June 1978, at Hamburg, Germany.* Heenemann, Berlin, vol.II, pp. 335-343.
- Butler, D. G. (1968): Hormonal control of gluconeogenesis in the north American eel (*Anguilla rostrata*). *General and comparative Endocrinology*, 10:85-91.
- Cassani, J. R. and Caton, W. E. (1986): Growth comparisons of diploid and triploid grass carp under varying conditions. *Prog. Fish-cult.*, 48:184-187.
- Catacutan, M. R. and Coloso, R. M. (1995): Effect of dietary protein to energy ratios on growth, survival and body composition of juvenile Asian seabass, *Lates calcarifer*. *Aquaculture*, 131:125-133.
- Catacutan, M. R. and Coloso, R. M. (1997): growth of juvenile Asian seabass, *Lates calcarifer*, fed varying carbohydrate and lipid levels. *Aquaculture*, 149:137-144.
- Chang, S. L. and Liao, I. C. (1996): Triploidy induced by heat shock in *Oreochromis aureus*. p. 273-279. In R. S. V. Pullin, J. Lazard, M. Legendre, B. J. Amon Kothias and D. Pauly (eds.). *The third international Symposium on Tilapia in Aquaculture.* ICLARM Conf. Proc. 41. 575 p.
- Cherfas, N. B., Gomelsky, B., Ben-Dom, N., Peretz, Y. and Hulata, G. (1994): Assessment of triploid common carp (*Cyprinus carpio* L.) for culture. *Aquaculture*, 127:11-18.
- Chervinsky, J. (1982): Environmental physiology of tilapia. In R.S.V. Pullin and R. H. Lowe-McConnell (Editors). *The biology and culture of tilapias.* International Center of Living Aquatic Resources Management, ICLARM conference proceedings 7, Manila. Philippines. pp. 119-128

- Cho, C. Y., Cowey, C. B. and Watanabe, T. (1985): Fish nutrition in Asia-methodological approaches to research and development. Ottawa:IDRC.
- Cho, C. Y., Slinger, S. J. and Bayley, H. S. (1976): Influence of level and type of dietary protein and level of feeding on feed utilization by rainbow trout. *J. Nutr.*, 106:1547-1556.
- Chou, B.-S. and Shiau, S.-Y. (1996): Optimal lipid level of growth of juvenile hybrid tilapia, *Oreochromis niloticus* × *Oreochromis aureus*. *Aquaculture*, 143:185-195.
- Chourrout, D. and Nakayama, T. (1987): Chromosome studies of progenies of tetraploid female rainbow trout. *Theor. Appl. Genet.*, 74:687-692.
- Chua, T. E. and Teng, S. K. (1982): Effects of food ration on growth, condition factor, food conversion efficiency, and net yield of estuary grouper, *Epinephelus salmonides* Maxwell, cultured in floating Net-cages. *Aquaculture*, 27:273-283.
- Cisse, A. (1996): Effect of the varying protein-energy levels on food consumption, growth and body composition of *Sarotherodon melanotheron* (Ruppel, 1852). P. 193-197. In R. S. V. Pullin, J. Lazard, M. Legendre, J. B. Amon Kothias and D. Pauly (eds.) the third international Symposium on Tilapia in Aquaculture. ICLARM Conf. Proc. 41, 575 P.
- Clark, A. E. Watanabe, W. O., Olla, B. L. and Wicklund, R. I. (1990): Growth, feed conversion and protein utilization of Florida red tilapia fed isocaloric diets with different protein levels in seawater pools. *Aquaculture*, 88:75-85.
- Clement, S. and Lovell, R. T. (1994): Comparison of processing yield and nutrition composition of cultured Nile tilapia (*Oreochromis niloticus*) and Channel catfish (*Ictalurus punctatus*). *Aquaculture*, 119:299-310.
- Coche, A.G., (1982): Cage culture of tilapias. In R.S.V. Pullin and R. H. Lowe-McConnell (editors). The biology and culture of tilapias. International Center of Living Aquatic Resources Management, ICLARM conference proceedings 7, Manila Philippines. Pp. 205-246.
- Cohen, D. and Ra'anani, Z. (1983): The production of the freshwater prawn *Macrobrachium rosenbergii* in Israel. III. Density effect of all-male tilapia hybrids on prawn yield characters in polyculture. *Aquaculture*, 35:57-71.
- Collis, W. J. and Smitherman, R. O. (1978): Production of tilapia hybrids with cattle manure of a commercial diet. pp 43-54. In R. O. Smitherman, W. L. Shelton and J. H. Grover(editors). Culture of exotic fishes symposium proceedings. Fish culture section, American Fisheries Society, Auburn, Alabama.
- Cortes Maldonado, R. C. and Pagan-Font, F. A. (1979): Evaluation of polyculture in plastic fishponds utilizing the catfish (*Ictalurus punctatus*) and blue tilapia (*Sarotherodon aureus*) fed with bird food. *Proc. Assoc. Isl.Mar. Lab. Caribb.*, 14:36.

- Craig, R. J. and Wolters, W. R. (1988): Source of variation in body size traits, dressout percentage and their correlations for the crayfish, *Procambarus clarkii*, Aquaculture, 72:49-58.
- Crawford, K. W., Dunseth, D. R., Engle, C. R., Hopkins, M. L., McCoy, E. W. and Smitherman, R. O. (1978): Marking tilapia and chinese carps. Fish culture section, American Fisheries Society, Auburn, Alabama. pp., 257.
- Cruz, E. M. and Laudencia, I. L. (1980): Polyculture of milkfish (*Chanos chanos Forskal*), all-male Nile tilapia (*Tilapia nilotica*) and Snakehead (*Ophicephalus striatus*) in freshwater ponds with supplemental feeding. Aquaculture, 20:231-237.
- Cruz, E. M. and Ridha, M. (1991): production of the tilapia *Oreochromis spilurus* Gunther stocked at different densities in sea cages. Aquaculture, 99:95-103.
- Dadzie, S. (1982): Species combination in tilapia culture. Aquaculture, 27:295-299.
- Dato-Cajegas, C. R. and Yakupitiyage, A. (1996): The need for dietary mineral supplementation for Nile tilapia, *Oreochromis niloticus*, cultured in a semi-intensive system. Aquaculture, 144:227-237.
- Degani, G. (1985): The influence of 17- α -methyltestosterone on body composition of eels (*Anguilla anguilla* L.). Aquaculture, 50:23-30.
- Degani, G. (1986): Effect of combined dietary 17- α -estradiol and 17- α methyltestosterone on growth and body composition of European eels (*Anguilla anguilla* L.). Aquaculture, 59:169-175.
- Degani, G., Dosoretz, C. and Levanon, D. (1982): Feeding *Sarotherodon aureus* with fermented cow manure. Bamidgeh, 34:119-129.
- Degani, G., Dosoretz, C. and Levanon, D. (1984): The influence of cow manure on growth rates of *Oreochromis niloticus* and *Clarias lazera* in Israel in small outdoor tanks. Bamidgeh, 36:114-120.
- Diana, J. S., Lin, C. K. and Schneeberger, P. J. (1991): Relationships among nutrient inputs, water nutrient concentrations, primary production, and yield of *Oreochromis niloticus* in ponds. Aquaculture, 92:323-341.
- Dioundick, O. B. and Stom, D. I. (1990): Effects of dietary α -cellulose levels on the juvenile tilapia, *Oreochromis niloticus* (peters). Aquaculture, 91:311-315.
- Don, J. and Avtalion, R. R. (1988): Production of viable tetraploid tilapias using the cold shock technique. Bamidgeh, 40:17-21.
- Duncan, D. B. (1955): Multiple range and multiple F test. Biometrics, 11:1-42.
- Eid, A. (1991): Sheep manure and poultry waste as feed materials for Nile tilapia, (*Oreochromis niloticus*). Zagazig J. Agric. Res., 18:1503-1510.

- Craig, R. J. and Wolters, W. R. (1988): Source of variation in body size traits, dressout percentage and their correlations for the crayfish, *Procambarus clarkii*, Aquaculture, 72:49-58.
- Crawford, K. W., Dunseth, D. R., Engle, C. R., Hopkins, M. L., McCoy, E. W. and Smitherman, R. O. (1978): Marking tilapia and chinese carps. Fish culture section, American Fisheries Society, Auburn, Alabama. pp., 257.
- Cruz, E. M. and Laudencia, I. L. (1980): Polyculture of milkfish (*Chanos chanos Forskal*), all-male Nile tilapia (*Tilapia nilotica*) and Snakehead (*Ophicephalus striatus*) in freshwater ponds with supplemental feeding. Aquaculture, 20:231-237.
- Cruz, E. M. and Ridha, M. (1991): production of the tilapia *Oreochromis spilurus* Gunther stocked at different densities in sea cages. Aquaculture, 99:95-103.
- Dadzie, S. (1982): Species combination in tilapia culture. Aquaculture, 27:295-299.
- Dato-Cajegas, C. R. and Yakupitiyage, A. (1996): The need for dietary mineral supplementation for Nile tilapia, *Oreochromis niloticus*, cultured in a semi-intensive system. Aquaculture, 144:227-237.
- Degani, G. (1985): The influence of 17- α -methyltestosterone on body composition of eels (*Anguilla anguilla* L.). Aquaculture, 50:23-30.
- Degani, G. (1986): Effect of combined dietary 17- α -estradiol and 17- α methyltestosterone on growth and body composition of European eels (*Anguilla anguilla* L.). Aquaculture, 59:169-175.
- Degani, G., Dosoretz, C. and Levanon, D. (1982): Feeding *Sarotherodon aureus* with fermented cow manure. Bamidgeh, 34:119-129.
- Degani, G., Dosoretz, C. and Levanon, D. (1984): The influence of cow manure on growth rates of *Oreochromis niloticus* and *Clarias lazera* in Israel in small outdoor tanks. Bamidgeh, 36:114-120.
- Diana, J. S., Lin, C. K. and Schneeberger, P. J. (1991): Relationships among nutrient inputs, water nutrient concentrations, primary production, and yield of *Oreochromis niloticus* in ponds. Aquaculture, 92:323-341.
- Dioundick, O. B. and Stom, D. I. (1990): Effects of dietary α -cellulose levels on the juvenile tilapia, *Oreochromis niloticus* (peters). Aquaculture, 91:311-315.
- Don, J. and Avtalion, R. R. (1988): Production of viable tetraploid tilapias using the cold shock technique. Bamidgeh, 40:17-21.
- Duncan, D. B. (1955): Multiple range and multiple F test. Biometrics, 11:1-42.
- Eid, A. (1991): Sheep manure and poultry waste as feed materials for Nile tilapia, (*Oreochromis niloticus*). Zagazig J. Agric. Res., 18:1503-1510.

- Eid, A. and Ghonim, S. I. (1994): Dietary zinc requirement of fingerling *Oreochromis niloticus*. *Aquaculture*, 119:259-264.
- Edwardson, W. (1976): Energy demands of Aquaculture: A world-wide survey. *Fish farming Int.*, 3(4):10-13.
- Edwards, P., Kaewpaitoon, K., Little, D. C. and Siripandh, N. (1994): An assessment of the role of buffalo manure for pond culture of tilapia. II. Field trial. *Aquaculture*, 126:97-106.
- Ekpo, I. and Bender, J. (1989): Digestibility of a commercial fish feed, wet algae, and dried algae by tilapia nilotica and Silver carp. *Prog. Fish-cult.*, 51:83-86.
- Eleraky, W., Saleh, G. and Gropp, J. M. (1995): A short note on the vitamin E requirement of tilapia nilotica (*Oreochromis niloticus*) - effect on health and growth. *J. Appl. Ichthyol.* 11:375-377.
- El-Husseiny, O. M., Arafa, S. A., Shehata, T. M. and Abd El-sami, A. M. (1993): Effect of dietary energy sources on tilapia performance. 4-th Symp. Anim. Poult. Fish Nut., El-Fayoum, 63-93.
- El-Sayed, A. M. and Teshima, S. (1992): Protein and energy requirements of Nile tilapia, *Oreochromis niloticus*, fry. *Aquaculture*, 103:55-63.
- Engle, C. and Skladany, M. (1992): The economic benefit of chicken manure utilization in fish production in Thailand. *CRSP Research Reports*, 92-45.
- Ensminger, M. E., Oldfield, J. E. and Heinemann, W. W. (1990): Feeds and nutrition. The Ensminger publ. Co., Clovis, California, 1169-1197.
- Eves, A., Turner, C., Yakupitiyage, A., Tongdee, N. and Ponza, S. (1995). The microbiological and sensory quality of septage-raised Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 132:261-272.
- Eya, J. C. and Lovell, R. T. (1997): Available phosphorus requirements of food-size channel catfish (*Ictalurus punctatus*) fed practical diets in ponds. *Aquaculture*, 154:283-291.
- Fagbenro, O. A. (1992): Quantitative dietary protein requirements of *Clarias Isberiensis* (Sydenham 1980) (Clariidae) fingerlings. *J. Appl. Ichthyol.*, 8:164-169.
- FAO (1995): Aquaculture production statics (1984-1993). Fisheries Circular. No. 815 Revision 7. Rome, Italy.
- FAO (1996): FAO training series 21/1. Simple methods for aquaculture, Management for freshwater fish culture. Ponds and water practices, Rome, Italy.
- Fauconneau, B., Alami-Durante, H., Laroche, M., Marcel, J. and Vallot, D. (1995): Growth and meat quality relations in carp. *Aquaculture*, 129:265-297.

- Fernando, C. H. (1982): Zooplankton and fish production, with special reference to the tilapias. In L. Fishelson and Z. Yaron (Editors), International symposium on tilapia in aquaculture, Proceeding, PP. 95-105, Nazareth, Israel.
- Flajshans, M., Linhart, O. and Kvasnicka, P. (1993): Genetic studies of tench (*Tinca tinca* L.): induced triploidy and tetraploidy and first performance data. *Aquaculture*, 113:301-312.
- Fox, M. G. (1989): Effect of prey density and prey size on growth and survival of juvenile Walleye (*Stizostedion vitreum*). *Can. J. Fish Aquat. Sci.*, 46:1323-1328.
- Fox, M. G., Flowers, D. D. and Watters, C. (1992): The effect of supplementary inorganic fertilization on juvenile Walleye (*Stizostedion vitreum*) reared in organically fertilized ponds. *Aquaculture*, 106:27-40.
- Gabel, S. J., Avault, J. W. and Romaine, R. P. (1981): Polyculture of Channel catfish (*Ictalurus punctatus*) with all-male tilapia hybrids (*Sarotherodon mossambica* male $\times$ *Sarotherodon hornorum* female). *J. world maricult. Sos.* 12(2):153-162.
- Galbreath, P. F., and Thorgaard, G. H. (1994): Saltshwater performance of all-female triploid Atlantic salmon. *Aquaculture*, 138:77-85.
- Galbreath, P. F., Jean, W. St., Anderson, V. and Thorgaard, G. H. (1994): Freshwater performance of all-female diploid and triploid Atlantic salmon. *Aquaculture*, 128:41-49.
- Geiger, J. G. (1983): A review of pond zooplankton production and fertilization for the culture of larval and fingerling striped bass. *Aquaculture*, 35:353-369.
- Geiger, J. G. and Parker, N. C. (1985): Survey of Striped pass hatchery management in the Southeastern United States. *Prog. Fish-cult.*, 47:1-13.
- Geri, G., Poli, B. M., Gualtieri, M., Lupi, P. and Parisi, G. (1995): Body traits and chemical composition of muscle in the Common carp (*Cyprinus carpio* L.) as influenced by age and rearing environment. *Aquaculture*, 129:329-333.
- Gershanovich, A. D., Markevich, N. M. and Dergaleva, Z. T. (1984): Using the condition factor in ichthyological research. *Journal of Ichthyology*, 24:78-90.
- Gill, H. S. and Weatherley, A. H. (1984): Protein, lipid and caloric contents of bluenose minnow, *Pimphales notatus Rafinesque*, during growth at different temperatures. *J. Fish Bio.*, 25:491-500.
- Gjerde, B. and Gjedrem, T. (1984): Estimation of phenotypic and genetic parameters for carcass traits in Atlantic salmon and rainbow trout. *Aquaculture*, 36:97-110.
- Graham, M. S., Fletcher, G. L. and Benfey, t. j. (1985): Effect of triploidy on blood oxygen content of atlantic salmon. *Aquaculture*, 50:133-139.

- Green, B. W. (1992): Substitution of organic manure for pelleted feed in tilapia production. *Aquaculture*, 101:213-222.
- Green, B. W., Phelps, R. and Alvarenga, H. (1989): The effect of manures and chemical fertilizers on the production of *Oreochromis niloticus* in earthen ponds. *Aquaculture*, 76:37-42.
- Green, B. W., Teichert-Coddington, D. R. and Phelps, R. P. (1990): Response of tilapia yield and economics to varying rates of organic fertilization and season in two central American countries. *Aquaculture*, 90:279-290.
- Grover, J. J. Olia, B. L., Brien, M. and Wicklund, R. I. (1989): Food habits of Florida red tilapia fry in manured seawater pools in the bahamas. *Prog. Fish-cult.*, 51:152-156.
- Guerrero, R. D. and Guerrero, L. A. (1988): Feasibility of commercial production of sex reversed Nile tilapia fingerlings in the Philippines. In Pullin, R. S. V., Bhukasawan, T., Tonguthai, K., Maclean, J. L. (eds.) 2nd Int. Symp. On Tilapia in Aquaculture. ICLARM Conf Proc. 15, Dept. of Fisheries. Bangkok, and ICLARM, Manila. pp 183-186.
- Gunasekera, R. M., Shim, K. F. and Lam, T. J. (1997): Influence of dietary protein content on the distribution of amino acids in oocytes, serum and muscle of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 152:205-221.
- Guo, X., DeBrosse, G. A., and Allen, S. K. (1996): All-triploid Pacific oysters (*Crassostrea gigas* Thunberg) produced by mating tetraploids and diploids. *Aquaculture*, 142:149-161.
- Guziur, J. and Markiewicz, F. (1987): The effect of intensive mineral fertilization of ponds on the state of carp, *Cyprinus carpio* L. *Acta Hydrobiol. Krakow*, 29:509-521.
- Haard, N. F. (1992): Control of chemical composition and food quality attributes of cultured fish. *Food Res. Int.*, 25:289-307.
- Habitzky-Biester, H. (1987): Vergleichende Untersuchungen Zu Erbruttungs-und Aufzuchtmethoden bei Afrikanischen Buntbarschen (*Oreochromis niloticus*). Dissertation, Universität Göttingen.
- Hafez, A. F. (1991): Studies on some productive traits in fish. Ph.D, Thesis, Faculty of Agriculture at Moshtohor, Zagazig University (Banha branch), Egypt.
- Halver, J. E. (1976): Formulating practical diets for fish. *J. Fish Res. Bd. Con.*, 33:1032.
- Halver, J. E. (1989): Fish nutrition. 2nd edn. New york, Academic press.
- Hanley, F. (1991): Effect of feeding supplementary diets containing varying levels of lipid on growth, food conversion, and body composition of Nile tilapia *Oreochromis niloticus*. *Aquaculture*, 93:323-334.

- Harvey, W. R. (1990): User's guide for LSMLMW. mixed model least-squares and maximum likelihood computer program. Ohio state University, Columbus, USA:
- Helal, H. and Culver, D. A. (1991): N:P ratio and plankton production in fish ponds. *Verh. Int. Verein. Limnol.*, 24:1508-1511.
- Henken, A. M., Brunink, A. M. and Richter, C. J. J. (1987): Differences in growth rate and feed utilization between Diploid and triploid African catfish, *Clarias gariepinus* (Burchell 1822). *Aquaculture*, 63:233-242.
- Hepher, B. and Pruginin, Y. (1981): Commercial fish farming, with special reference to fish culture in Israel. J. Wiley, New york, 261 pp.
- Herodek, S., Tatrai, I. and Olah, J. (1989): Feeding experiments with silver carp (*Hypophthalmichthys molitrix* val.) fry. *Aquaculture*, 83:331-344.
- Hillestad, M. and Johnsen, F. (1994): high-energy/low-protein diets for Atlantic salmon: effect on growth, nutrient retention and slaughter quality. *Aquaculture*, 124:109-116.
- Hoffman, L. C., Prinsloo, J. F., Theron, J. and Casey, N. H. (1995): A chemical comparison between the golden and normal coloured strains of the A African sharptooth catfish, *Clarias gariepinus* (Burchell 1822). *J. Appl. Ichtyol.* 11:71-85.
- Hogendoorn, H. and Koops, W. J. (1983): Growth and production of African catfish, *Clarias lazera*. 1. Effects of stocking density, pond size and mixed culture extensive field conditions. *Aquaculture*, 34:253-263.
- Holland, R. A. B. (1970): Factors determining the rate of gas uptake by intracellular hemoglobin. In: D. Hershey (Editor), blood oxygenation. Plenum Press, New York, NY, pp. 1-23.
- Hopkins, K. D. and Cruz, E. M. (1982): The ICLARM-CLSU integrated animal-fish farming project: final report. ICLARM technical Report 5. International Center for Living Aquatic Resources Management, Manila, and the Freshwater Aquaculture center, Central Luzon State University, Nueva Ecija, Philippines, 96 pp.
- Hung, S. S. O., Conte, F. S. and Hallen, E. F. (1993): Effect of feeding rates on growth, body composition and nutrient metabolism in striped bass (*Morone saxatilis*) fingerlings. *Aquaculture*, 112:349-361.
- Hung, S. S. O., Liu, W., Li, W., Storebakken, T. and Cui, Y. (1997): Effect of starvation on some morphological and biochemical parameters in white sturgeon, *Acipenser transmontanus*. *Aquaculture*, 151:357-363.
- Hussain, M. G., Chatlerji, A., McAndrew, B. J. and Johnstone, R. (1991): Triploidy induction in Nile tilapia, *Oreochromis niloticus* L. using pressure, heat and cold socks. *Theor. Appl. Genet.*, 81:6-12.

- Hussain, M. G., Penman, D. J., and McAndrew, B. J. (1996): Effect of triploidy on sexual maturation and reproduction in Nile tilapia *Oreochromis niloticus* L. p. 320-325. In R. S. V. Pullin, J. Lazard, M. Legendre, B. J. Amon Kothias and D. Pauly (eds.). The third international Symposium on Tilapia in Aquaculture. ICLARM Conf. Proc. 41. 575 p.
- Hussain, M. G., Rao, G. P. S., Humayun, N. M., Randall, C. F., Penman, D. J., Kime, D., Bromage, N. R., Myers, J. M. and McAndrew, B. J. (1995): Comparative performance of growth, biochemical composition and endocrine profiles in diploid and triploid tilapia *Oreochromis niloticus* L. *Aquaculture*, 138:87-97.
- Hussein, A. A. (1990): Biochemical studies on fish by-products. M. Sc. Thesis, Faculty of Agriculture, Zagazig University, Egypt.
- Ince, W. B. and Thorpe, A. (1976): The effect of starvation and force-feeding on the metabolism of the northern pike, *Esox lucius* L. *J. Fish biol.*, 8:79-88.
- Jauncey, K. (1982): The effect of varying dietary protein level on the growth, food conversion, protein utilization and body composition of juvenile tilapia *Oreochromis mossambicus*. *Aquaculture*, 27:43-54.
- Jauncey, K. and Rose, B. (1982): A guide to tilapia feeds and feeding. Institute of Aquaculture, University of Sterling, Scotland.
- Johnstone, R. (1989): maturity control in Atlantic salmon-a review of the current status of research in Scotland. In Symp. On Applications of endocrinology in Aquaculture, Malaga, Spain.
- Johnson, O. W., Dickhoff, W. W. and Utter, F. M. (1986): Comparative growth and development of diploid and triploid coho salmon, *Oncorhynchus kisutch*. *Aquaculture*, 57:329-336.
- Joseph, D. (1981): Partial replacement of concentrated fish food (pellets) with chicken droppings. *Bamidgeh*, 33:24-28.
- Kanazawa, A., Teshima, S. I., Sakamoto, M. and Awal, M. A. (1980): Requirement of tilapia zilli for essential fatty acids. *Bull. Jpn. Soc. Sci. Fish.*, 46:1353-1356.
- Kayano, Y., Yao, S., Yamamoto, S. and Nakagawa, H. (1993): Effect of feeding frequency on the growth and body compositions of young red-spotted grouper, *Epinephelus akaara*. *Aquaculture*, 110:271-278.
- Kepshire, B. M., Tinsley, I. J. and Lowry, R. R. (1983): Effect of temperature on the chemical composition of pink salmon (*Oncorhynchus gorbuscha*) muscle. *Aquaculture*, 32:295-301.
- Kerby, J. H. Everson, J. M., Harrell, R. M., Starling, C. C., Revels, H. and Geiger, J. G. (1995): Growth and survival comparisons between diploid and triploid sunshine bass. *Aquaculture*, 137:355-358.

- Khater, A. A. and Smitherman, R. O. (1988): Cold tolerance and growth of *Oreochromis niloticus*. pp. 215-218. In: In R. S. V. Pullin, T. Bhukaswan, K. Tonguthai and J. L. Maclean (editors). The second international symposium on tilapia in aquaculture. Conference Proceedings 15, 623 pp. International Center for Living Aquatic Resources Management, Manila, Philippines.
- Kiessling, A., Johansson, L. and Storebakken, T. (1989). Effects of reduced feed ration levels of fat content and fatty acid composition in white and red muscle from rainbow trout. *Aquaculture*, 79:169-175.
- Kim, M. K. and Lovell, R. T. (1995): Effect of restricted feeding regimens on compensatory weight gain and body tissue changes in channel catfish, *Ictalurus punctatus* in ponds. *aquaculture*, 135:285-293.
- Kirchgessner, M. and Schwarze, F. F. (1986): Mineral content (major and trace elements) of carp (*Cyprinus carpio* L.) fed with different protein and energy supplies. *Aquaculture*, 54:3-9.
- Kirk, R. G. (1972): A review of recent developments in tilapia culture, with special reference to fish farming in the heated effluents of power stations. *Aquaculture*, 1:45-60.
- Kligerman, A. D. and Bloom, S. E. (1977): Rapid chromosome preparations from solid tissues of fish. *J. Fish Res. Board Can.*, 34:266-269.
- Knud-Hansen, C. F. and Batterson, T. R. (1994): Effect of fertilization frequency on the production of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 123:271-280.
- Kuwaye, T. T., Okimoto, D. K., Shimoda, S. K., Howerton, R. D., Lin, H. R. Pnag, P. K. T. and Grau, E. G. (1993): Effect of 17 α -methyltestosterone on the growth of euryhaline tilapia, *Oreochromis niloticus*, in fresh water and in sea water, *Aquaculture*, 113:137-152.
- Laws, E. D. and Weisburd, R. S. J. (1990): Use of silver carp to control algae biomass in aquaculture ponds. *Progre. Fish-Cultur.*, 52:1-8.
- Li, M. and Lovell, R. T. (1992, a): Growth, feed efficiency and body composition of second and third year Channel catfish fed various concentrations of dietary protein to satiety in production ponds. *Aquaculture*, 103:153-163.
- Li, M. and Lovell, R. T. (1992, b): Comparison of satiate feeding and restricted feeding of channel catfish with various concentrations of dietary protein in production ponds. *Aquaculture*, 103:165-175.
- Li, S., Yang, H. and Lu, W., (1980): Primary study on daily food consumption of silver carp (*Hypophthalmichthys molitrix*), bighead (*Aristichthys nobilis*) and grass carp (*Ctenopharyngodon idellus*). *J. Fish. China*, 4:275-283.

- Liao, H., Pierce, C. L., Wahl, D. H., Rasmussen, J. B. and Leggett, W. C. (1995): Relative weight (Wr) as a field assessment tool: Relationships with growth, Prey biomass, and environmental conditions. *Trans. Amer. Fish. Soc.*, 124:387-400.
- Likongwe, J. S., Stecko, T. D. Stauffer, J. R. and Carline, R. F. (1996): Combination effects of water temperature and salinity on growth and feed utilization of juvenile Nile tilapia, *Oreochromis niloticus*. *Aquaculture*, 146:37-46.
- Lim, C., Ako, H., Brown, C. L. and Hahn, K. (1997): Growth response and fatty acid composition of juvenile *penaeus vannamei* fed different sources of dietary lipid. *Aquaculture*, 151:143-153.
- Lin, H.-R. (1982): Polycultural system of freshwater fish in China. *Can. J. Fish. Aquat. Sci.*, 39:143-150.
- Lin, C. K., Jaijen, K. and Muthuwan, V. (1993): Integration of intensive and semi-intensive aquaculture: concept and example. CRSP Research Reports, 93-54.
- Lin, J.-H., Cui, Y., Hung, S. S. O. and Shiau, S.-Y. (1997): Effect of feeding strategy and carbohydrate source of carbohydrate utilization by white sturgeon (*Acipenser transmontanus*) and hybrid tilapia (*Oreochromis niloticus* × *O. aureus*). *Aquaculture*, 148:201-211.
- Love, R. M. (1970): The chemical biology of fishes. Academic press, London.
- Love, R. M. (1988): The Food fishes: Their intrinsic variation and practical value. Farrand press, London, 276 pp.
- Lovell, R. T. (1980): Feeding tilapia. *Aquaculture magazine*, 7:42-43.
- Lovell, R. T. (1981): Laboratory manual for fish feed analysis and fish nutrition studies. Auburn University, Alabama.
- Lovell, R. T. (1989): Nutrition and feeding of fish. Van Nostrand Reinhold, New York.
- Lovshin, L. L. Tave, D. and Leieutaud, A. O. (1990): Growth and yield of mixed-sex, young-of-the year *Oreochromis niloticus* raised at two densities in earthen ponds in Alabama, USA. *Aquaculture*, 89:21-26.
- Lutes, P. B., Hung, S. S. O. and Conte, F. S. (1990): Survival, growth and body composition of white sturgeon larvae fed purified and commercial diets at 14.7 and 18.4°C. *Prog. Fish-cult.*, 52:192-196.
- Macari, M., Carneiro, D. J., Larson, M. L. and Machado, C. R. (1994): Influence of dietary protein intake and recombinant human somatotropin administration on growth and body composition of juvenile tambacu (a *Piaractus mesopotamicus* × *Colossoma macropomum* cross). *Aquaculture*, 127:363-369.

- Maguire, G. B., Gardner, N. C., Nell, J. A., Kent, G. N. and Kent, A. S. (1995): Studies on triploid oyster in Australia. 2. Growth, condition index gonad area, and glycogen content of triploid and diploid Pacific oyster, *Crassostrea gigas*, from oyster leases in Tasmania, Australia. *Aquaculture*, 137:355-358.
- Maibom, H. (1992): Untersuchung zur induzierten gynogenese beim *Oreochromis niloticus*. M.Sc. Thesis, Institut für Tierzucht und Haustiergenetik and der Georg-August-Universität, Göttingen.
- Mair, G. C. (1993): Chromosome set manipulation in tilapia-techniques, problems and prospects. *Aquaculture*, 111:227-244.
- Mair, G. C., Abucay, J. S., Beardmore, J. A. and Skibinski, D. O. F. (1995): Growth performance trials of genetically male tilapia (GMT) derived from YY-males in *Oreochromis niloticus* L.; On station comparisons with mixed sex and sex reversed male populations. *Aquaculture*, 137:313-322.
- Mair, G. C. and Little, D. C. (1991): Population control in farmed tilapia. *NAGA*, 14:8-13.
- McCarthy, I. D., Carter, C. G., Houlihan, D. F., Johnstone, R. and Mitchell, A. I. (1996): The performance of all-female diploid and triploid Atlantic salmon smolts on transfer together to sea water. *J. Fish Biol.*, 48:545-548.
- McGeachin, R. B. and Stickney, R. R. (1982): Manuring rates for production of Blue tilapia in simulated sewage lagoons receiving laying hen waste. *Prog. Fish-cult.*, 44:25-28.
- McGeachy, S. A., Benfey, T. J. and Friars, G. W. (1995): Freshwater performane of triploid Atlantic salmon (*Salmo salar*) in New Brunswick aquaculture. *Aquaculture*, 137:333-341.
- McGinty, A. S. (1985): Effect of predation by largemouth bass in fish production ponds stocked with *Tilapia nilotica*. *Aquaculture*, 96:119-138.
- Merola, N., Colares de Melo, J. S. and Costa Nascimento, V. M. (1984): Production of the tilapia hybrid *Oreochromus hornorum* (male) $\times$ *O. niloticus* (female) and Common carp in polyculture at different stocking densities. *Annals of the third Brazilian symposium on aquaculture*, pp. 277-285.
- Milstein, A., Alkon, A., Avnimelech, Y., Kachba, M., Hulata, G. and Schroeder, G. (1991): Effects of manuring rate on ecology and fish performance in polyculture ponds. *Aquaculture*, 96:119-138.
- Mires, D. (1982): A study of the problems of mass production of hybrid tilapia fry. pp. 317-329. In R.S.V. Pullin and R. H. Lowe-McConnell (editors). *The biology and culture of tilapias*. International Center of Living Aquatic Resources Management, ICLARM conference proceedings 7, Manila. Philippines.
- Mires, D. (1995): The Tilapias. In C. E. Nash and A. J. Novotny (editors), *production of aquatic animals*. pp. 133-152.

- Moav, R., Wohlfarth, G., Schroeder, G. L., Hulata, G. and Barash, H. (1977): Intensive polyculture of fish in freshwater ponds. 1-Substitution of expensive feeds by liquid cow manure. *Aquaculture*, 10:25-43.
- Moriarty, C. M. and Moriarty, D. J. W. (1973): Quantitative estimation of the daily ingestion of phytoplankton by *Tilapia nilotica* and *Haplochromis nigripinnis* in Lake George, Uganda. *J. Zoo., London*, 171:15-23.
- Morkramer, S., Hörstgen-Schwark, G. and Langholz, H. J., (1985): Comparison of different European rainbow trout populations under intensive production conditions. *Aquaculture*, 44:303-32.
- Muratat, H. and Higashi, T. (1980): Selective utilization of fatty acids as energy in carp. *Bulletin of the Japanese Society of Scientific fisheries*, 46:1333-1338.
- Myers, J. M. (1986): Tetraploid induction in *Oreochromis* spp. *Aquaculture*, 57:281-287.
- Myers, J. M. and Hershberger, W. K. (1991): Early growth and survival of heat-shocked and tetraploid-derived triploid rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 96:97-107.
- NRC (1983): Nutrition requirements of warmwater fishes and shellfishes. Washington, Dc. National Academy of Science
- Nagy, A., (1987): Genetic manipulation performed on warm water fish. In: K. Tiews (ed.). *Proc. World Symp. Selection, Hybridization and Genetic Engineering in Aquaculture*, Bordeaux, 27-30 May 1986. Heenemann, Berlin, pp. 127-145.
- Nell, J. A. Cox, E., Smith, I. R. and Maguire, G. B. (1995): Studies on triploid oysters in Australia. 1. The farming potential of triploid Sydney rock oysters *Saccostrea commercialis* (Iredale and Roughley). *Aquaculture*, 137:355-358.
- Nematipour, G. R., Brown, M. L. and Gatlin, D. M. III (1992): Effect of dietary energy : protein ratio on growth characteristics and body composition of hybrid striped bass, (*Morone chrysops* females $\times$ *M. saxatilis* males). *Aquaculture*, 107:359-368.
- Nilsson, J., (1992): Genetic parameters of growth and sexual maturity in Arctic char (*Salvelinus alpinus*). *Aquaculture*, 106:9-19.
- Noeske-Hallin, T. A., Spieler, R. E., Parker, N. C. and Stuttle, M. A. (1985): Feeding time differentially affects fattening and growth of Channel catfish. *J. Nutr.*, 115:1228-1232..
- O'connell, J. P. and Gatlin, D. M. (1994): Effect of dietary calcium and vitamin D₃ on weight gain and mineral composition of the blue tilapia (*Oreochromis aureus*) in low-calcium water. *Aquaculture*, 124:107-117.

- Ostrowski, A. C. and Garling, D. L. (1988): Influences of anabolic hormone treatment and dietary protein:energy ratio on condition and muscle deposition of rainbow trout. *Prog. Fish-cult.*, 50:133-140.
- Papoutsoglou, S. E., Petropoulos, G. and Barbieri, R. (1992): Polyculture rearing of *Cyprinus carpio* (L.) and *Oreochromis aureus* (St.) using a closed circulated system. *Aquaculture*, 103:311-320.
- Parsons, G. R. (1993): Comparisons of triploid and diploid white crappies. *Trans. Am. Fish. Soc.*, 122:237-243.
- Parsons, G. J. and Dadswell, M. J. (1992): Effect of stocking density on growth, production and survival of the giant scallop, *Placopecten magellanicus*, held in intermediate suspension culture in passamaquoddy Bay, New Brunswick. *Aquaculture*, 103:291-309.
- Payne, A. I. and Collinson, R. I. (1983): A comparison of the biological characteristics of *Sarotherodon niloticus* (L.) with those of *S. aureus* (Steindachner) and other tilapia of the delta and lower Nile. *Aquaculture*, 30:335-351.
- Perera, P. A. B. and De Silva, S. S. (1978): Studies on the biology of young grey mullet, *Mugil cephalus* L. *J. Fish Biol.*, 13:297-304.
- Pillay, T. V. R. (1988): *Aquaculture principles and practices*. Fishing News Book. Pp. 360-376.
- Plavnik, H. Barash, H. and Schroeder, G. L. (1983): Utilization of ducks droppings in fish farming. *Nutrition reports international*, 28:635-641.
- Poston, H. A. and Rumsey, G. L. (1983): Factors affecting dietary requirement and deficiency signs of L-tryptophan in rainbow trout. *J. Nutr.*, 113:2568-2577.
- Power, D. M. (1990): The physiology of growth. *Trout news*, 10:20-23.
- Puckhaber, B. and Hörstgen-Schwark, G. (1996): Growth and gonadal development of triploid tilapia (*Oreochromis niloticus*). p. 377-382. In R. S. V. Pullin, J. Lazard, M. Legendre, B. J. Amon Kothias and D. Pauly (eds.). *The third international Symposium on Tilapia in Aquaculture*. ICLARM Conf. Proc. 41. 575 p.
- Pullin, R. S. V. (1983): Choices of tilapia species for aquaculture. In L. Fishelson and Z. Yaron (Editors), *International Symposium on Tilapia in Aquaculture*, Proceeding, 64-76. Nazareth, Israel.
- Pullin, R. S. V. (1991): Cichlids in aquaculture. P 280-309. In M. H. A. Keenleyside (ed.) *Cichlid fishes: behaviour, ecology and evolution*. Chapman and Hall, London.
- Qin, J. and Culver, D. A. (1992): The survival and growth of larval walleyes, *Stizostedion vitreum*, and trophic dynamic in fertilized ponds. *Aquaculture*, 108:257-276.

- Qin, J. and Culver, D. A. and Ya, N. (1994): Comparative growth of larval walleye and saugeye, and their impact on zooplankton in experimental ponds. *Prog. Fish-cult.*, 56:91-99.
- Qin, J. Madon, S. P. and Culver, D. A. (1995): Effect of larval walleye (*Stizostedion vitreum*) and fertilization on plankton community: implications for larval fish culture. *Aquaculture*, 130:51-65.
- Recoubratsky, A. V., Gomelsky, B. I. And Emelyanova, O. V. and Pankratyeva, E. V. (1992): Triploid common carp produced by heat shock with industrial fish-farm technology. *Aquaculture*, 108:13-19.
- Reich, K. (1975): Multispecies fish culture (polyculture) in Israel. *Bamidgeh*, 27:85-99.
- Reinitz, G. and Hitzel, F., (1980): Formulation of practical diets for rainbow trout based on desired performance and body composition. *Aquaculture*, 19:243-252.
- Rhodes, C. P. and Holdich, D. M. (1984): Length-weight relationship, muscle production and proximate composition of the freshwater cryfish (*Austropotambius pallipes*). *Aquaculture*, 37:107-123.
- Robinson, E., Labomascus, D., Brown, D. B. and Linton, L. T., (1987): Dietary calcium and phosphorus requirement of *O. aureus* reared in calcium-free water. *Aquaculture*, 64:267-276.
- Roem, A. J., Stickney, R. R. and Kohler, C. C. (1991): Dietary pantothenic acid requirement of the blue tilapia. *Prog. Fish-Cult.*, 53:216-219.
- Rothbard, S. (1979): Observations on the reproductive behaviour of tilapia zillii and several sarotherodon spp. Under aquarium conditions. *Bamidgeh*, 31:35-43.
- Saad, C. R. (1979): Use of full-fat roasted soybean in practical catfish diet. M.Sc. Thesis, Auburn University, Alabama.
- Saleh, G., Eleraky, W. and Gropp, J. M. (1995): A short note on the effects of vitamin a hypervitaminosis and hypovitaminosis on health and growth of tilapia nilotica (*Oreochromis niloticus*) *J. Appl. Ichthyol.*, 11:382-385.
- Santiago, C. B. (1985): Amino acid requirements of Nile tilapia. Ph.D. thesis, Auburn University.
- Santiago, C. B. and Reyes, O. S. (1993): Effect of dietary lipid source on reproductive performance and tissue levels of Nile tilapia *Oreochromis niloticus* (Linnaeus) broodstock. *J. Appl. Ichthyol.*, 9:33-40.
- Sarig, S. (1983): A review on tilapia culture in Israel. International symposium on tilapia in aquaculture. Proceeding, pp.116-122, Nazareth, Israel.

- Sarig, S. (1995): The development of polyculture in Israel: a model of intensification. In C. J. Shepherd and N. R. Bromage (Editors). Intensive fish farming. pp. 302-332. Hartonlls Ltd, Bodmin, cornwall, Great Britain.
- Satoh, S., Takeuchi, T. and Watanabe, T. (1984): Effect of starvation and environmental temperature on proximate and fatty acid composition of tilapia nilotica. Bulletin of Japanese Society of Scientific Fisheries, 50:79-84.
- Satoh, S., Takeuchi, T. and Watanabe, T. (1987): Requirement of Tilapia for alphatocopherol. Nippon Suisan Gakkaishi, 53:119-124.
- Schoonbee, H. J. and Prinsloo, J. F. (1988): The use of fish polyculture in integrated agriculture-aquaculture production systems aimed at rural community development. J. Aquat. Prod., 2:99-123.
- Schroeder, G. L. (1978): Autotrophic and heterotrophic production of microorganisms intensively-manured fish ponds, and related fish yields. Aquaculture, 141:303-325.
- Schroeder, G. L. (1979): Microorganisms as the primary diet in fish farming. World symposium on finfish nutrition and fish feed technology, Hamburg (GFR), 20 jun.
- Schroeder, G. L. (1980): Fish farming in manure-loaded ponds. Pages 73-86 in R. S. V. Pullin and Z. H. Shehadeh (Editors). Integrated agriculture aquaculture farming systems, International Center for Living Aquatic Resources, Management, Manila.
- Schroeder, G. L. (1983): The role of natural foods in tilapia growth. A study based on Stable isotope analysis. In L. Fishelson and Z. Yaron (Editors), International symposium on tilapia in aquaculture, Proceeding, PP. 313-322, Nazareth, Israel.
- Schroeder, G. L., Wohlfarth, G., Alkon, A., Halevy, A. and Krueger, H. (1990): The dominance of algal-based food webs in fish ponds receiving chemical fertilizers plus organic manures. Aquaculture, 86:219-229.
- Sevrin-Reyssac, J. and Pletikotic, M. (1990): Cyanobacteria in fish ponds. Aquaculture, 88:1-20.
- Shapiro, J. (1985): Food and intestinal contents of silver carp, *Hypophthalmichthys molitrix* in lake Kinneret between 1982-1984. Bamidgeh, 37:3-18.
- Shearer, K. D. (1994): Factors affecting the proximate composition of cultured fishes with emphasis on salmonids. Aquaculture, 119:63-88.
- Shearer, K. D. and Hardy, R. W. (1987): Phosphorus deficiency in rainbow trout fed a diet containing deboned fillet scrap. Prog. Fish-Cult., 49:192-197.
- Shearer, K. D., Maage, A., Opstvedt, J. and Mundheim, H. (1992): Effect of high ash diets on growth, feed efficiency, and zinc status of juvenile atlantic salmon (*Salmo salar*). Aquaculture, 106:345-355.

- Shelbourn, J. E., Clarke, W. C., McBirde, J. R., Fagerlund, U. H. M. and Donaldson, E. M. (1992): The use of 17 α -methyltestosterone and 3,5,3-triiodo-L-thyronine for sterilizing and accelerating the growth of zero-age coho salmon smolts (*Oncorhynchus kisutch*). *Aquaculture*, 103:85-99.
- Shiau, S.-Y. and Chuang, J.-C. (1995): Utilization of disaccharides by juvenile tilapia *Oreochromis niloticus* $\times$ *O. aureus*. *Aquaculture*, 133:249-256.
- Shiau, S.-Y. and Hsu, T.-S. (1995): L-Ascorbyl-2-sulfate has equal antiscorbutic activity as L-ascorbyl-2-monophosphate for tilapia *Oreochromis niloticus* $\times$ *O. aureus*. *Aquaculture*, 133:147-157.
- Shiau, S.-Y. and Huang, S.-L. (1989): Optimal dietary levels for hybrid tilapia (*Oreochromis niloticus* $\times$ *O. aureus*) reared in seawater. *Aquaculture*, 81:119-127.
- Shiau, S.-Y. and Huang, S.-L. (1990): Influence of varying energy levels with two protein concentrations in diets for hybrid tilapia (*Oreochromis niloticus* $\times$ *O. aureus*) reared in seawater. *Aquaculture*, 91:143-152.
- Shiau, S.-Y. and Peng, C.-Y. (1992): Utilization of different carbohydrates at different dietary protein levels in grass prawn *Penaeus monodon* reared in seawater. *Aquaculture*, 101:241-250.
- Siddiqui, A. and Al-Harbi, A. (1995). Evaluation of three species of tilapia, red tilapia and hybrid tilapia as culture species in Saudi Arabia. *Aquaculture*, 138:147-157.
- Siddiqui, A. and Alnazada, A. R. and Alhinty, H. M. (1993): Growth, survival and production of hybrid tilapia, common carp and sharptooth catfish I mono and polyculture systems during winter in central Saudi Arabia. *Prog. Fish-Cult.*, 55:57-59.
- Siddiqui, A., Howlodder, M. S. and Adam, A. A. (1988): Effects of dietary protein levels on growth, feed conversion and protein utilization in fry and young Nile tilapia, *Oreochromis niloticus*. *Aquaculture*, 70:63-73.
- Small, S. A. and Benfey, T. (1987): Cell size in triploid salmon. *J. Exp. Zool.*, 241:339-342.
- Smith, D. W. (1988): Phytoplankton and catfish culture; a review. *Aquaculture*, 74:167-189.
- Snow, J. R. (1983): Size control of catfish fingerlings by varying the stocking density. International programs. Auburn, Alabama.
- Soliman, A. K. and Wilson, R. P. (1992 a): Water-soluble vitamin requirements of tilapia. 1. Pantothenic acid requirement of blue tilapia, *Oreochromis aureus*. *Aquaculture*, 104:121-126.

- Soliman, A. K. and Wilson, R. P. (1992 b): Water-soluble vitamin requirements of tilapia. 2. Riboflavin requirement of blue tilapia, *Oreochromis aureus*. *Aquaculture*, 104:309-314.
- Spataru, P. (1976): Natural feed of tilapia aurea Steindachner in polyculture, with supplementary feed and intensive manuring. *Bamidgeh*, 28:57-63.
- Spataru, P. (1982): A contribution to the study of the natural food of *Sarotherodon* hybrids grown under conditions of polyculture, supplementary feed and intensive fertilization. *Bamidgeh*, 34:144-157.
- Spataru, P. Wohlfarth, C. W. and Hulata, G. (1983): Studies on the natural food of different fish species in intensively manured polyculture ponds. *Aquaculture*, 35:283-298.
- Steffen, W. (1989): Principles of fish nutrition. New York; Ellis Horwood.
- Stickney, R. R. (1986): Tilapia tolerance of saline waters: A review. *Prog. Fish-Cult.*, 48:161-167.
- Stickney, R. R. and Andrews, J. W. (1971): Combined effects of dietary lipids and environmental temperature on growth, metabolism and, body composition of channel catfish (*Ictalurus punctatus*). *J. Nutr.*, 101:1703-1710.
- Stickney, R. R. and Hardy, R. W. (1989): Lipid requirements of some warmwater species. *Aquaculture*, 79:145-156.
- Stickney, R. R., Hesby, J. H., McGeachin, R. B. and Isbell, W. A. (1979): Growth of tilapia nilotica in ponds with differing histories of organic fertilization. *Aquaculture*, 17:189-194.
- Stickney, R. R. and McGeachin, R. B. (1984): Growth, food conversion and survival of fingerling tilapia aurea fed differing levels of dietary beef tallow. *Prog. Fish-Cult.*, 46:102-105.
- Stillwell, E. J. and Benfey, T. J. (1995): Hemoglobin level, metabolic rate and swimming performance in triploid brook trout (*Salvelinus fontinalis*). *Aquaculture*, 137:355-358.
- Suresh, A. V. and Lin, C. K. (1992): Tilapia culture in saline waters: A review, *Aquaculture*, 106:201-226.
- Suzuki, R., Nakanishi, T. and Oshiro, T. (1985): Survival, growth and sterility of induced triploids in the Cyprinid loach *Misgurnus anguillicaudatus*. *Bulletin of the Japanese Society of Scientific Fisheries*, 51:889-894.
- Szarski, H. (1976): Cell size and nuclear DNA content in vertebrates. *Int. Rev. Cytol.*, 44:93-111.

- Takeuchi, T., Satoh, S. and Watanabe, T. (1983): Dietary lipids suitable for the practical feed of *tilapia nilotica*. Bull. Jpn. Soc. Fish., 49:1361-1365.
- Tang, Y. A. (1970): Evaluation of balance between fishes and available fish foods in multispecies fish culture ponds in Taiwan. Trans. Am. Fish. Soc., 99:708-718.
- Tayamen, M. M. and Shelton, W. L. (1978): Inducement of sex reversal in *Sarotherodon niloticus* (Linnaeus). Aquaculture, 14:349-354.
- Teichert-Coddington, D. R. (1996): Effect of stocking ratio on semi-intensive polyculture of *Colossoma macropomum* and *Oreochromis niloticus* in Honduras, central America. Aquaculture, 143:291-302.
- Teichert-Coddington, D. R. Behrends, L. L. and Smitherman, R. O. (1990): Effects of manuring regime and stocking rate on primary production and yield of tilapia using liquid swine manure. Aquaculture, 88:61-68.
- Teichert-Coddington, D. R. and Green, B. W. (1990): Influence of primary productivity, season and site on tilapia production in organically fertilized ponds in two central American countries. Paper EIFAC/90/Symp. E8, FAO-EIFAC. Symposium on production enhancement in Still Water Pond Culture, Prague, Czechoslovakia.
- Teichert-Coddington, D. R. and Green, B. W. (1993): Influence of day length and incubation interval on water column respiration in tropical fish ponds. Hydrobiologia, 250:159-165.
- Torrans, L. and Hiott, D. (1990): Effect of Brood-stock density on production of Bait- or Forage-Sized Blue tilapias. Prog. Fish-Cult., 52:9-14.
- Torrans, L. ad Lowell. F. (1987): Effects of blue tilapia/channel catfish polyculture on production, food conversion, water quality and channel catfish off-flavor. Arkansas Academy of Science Proceedings, 41:82-86.
- Tung, P.-H. and Shiau, S.-Y. (1991): Effect of meal frequency on growth performance of hybrid tilapia, *Oreochromis niloticus* $\times$ *O. aureus*, fed different carbohydrate diets. Aquaculture, 92:343-350.
- Von der decken, A. (1989): Ration size in the feeding of cod (*Gadus morhua*): effect on skeletal muscle proteins, with special reference to myosin heavy chain. Aquaculture 79:47-52.
- Wange, J.-Q., Flickinger, S. A., Be, K., Liu, Y. and Xu, H. (1989): Daily food consumption and feeding rhythm of silver carp (*Hypophthalmichthys molitrix*) during fry to fingerling period. Aquaculture, 83:73-79.
- Watanabe, T. (1995): Nutrition and growth. In C. J. Shepherd and N. R. Bromage. Intensive fish farming. pp. 154-197. Hartnolls Ltd, Bodmin, Cornwall, Great Britain.

- Watanabe, W. O., Chan, J. R., Smith, S. J., Wicklund, R. I. and Olla, B. L. (1996). Production of Florida red tilapia (*Oreochromis sp.*) in flowthrough seawater pools at three stocking densities. P. 168-174. In R. S. V. Pullin, J. Lazard, M. Legendre, J. B. Amon Kothias and D. Pauly (eds.) the third international Symposium on Tilapia in Aquaculture. ICLARM Conf. Proc. 41, 575 P.
- Watanabe, W. O., Clark, J. H., Dunham, J. B., Wicklund, R. I. and Olla, B. L. (1990). Culture of Florida red tilapia in marine cages: thre effect of stocking density and dietary protein on growth. *Aquaculture*, 90:123-134.
- Weatherley, A. H. and Gill, H. (1987): The biology of fish growth. Academic press, London, pp. 443.
- Weatherup, R. N., McCracken, K. J., Foy, R., Rice, D., Mckendry, J., Mairs, R. J. and Hoey, R. (1997): The effects of dietary fat content on performance and body composition of farmed rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 1515:173-184.
- Wedekind, H. (1995): Dietary influences on product quality in African catfish (*Clarias gariepinus*). *J. Appl. Ichthyol.*, 11:347-353.
- Wellner, E. (1981): On the effect of hogher stocking densities of silver carp on zooplankton production in eutrophic waters. *Binnen Fish. DDR.*, 28:200-202.
- Williams, K., Gebhart, G. E. and Maughan (1987): Enhanced growth of cage cultured channel catfish through polyculture with blue tilapia. *Aquaculture*, 62:207-214.
- Winfree, R. A. and Stickney, R. R. (1981): Effects of dietary protein and energy on growth, feed conversion efficiency, and body composition of tilapia aurea. *J. Nutr.*, 111:1001-1012.
- Winkelman, A. M. and Peterson, R. G. (1994): Genetic parameters (heritabilities, dominance ratios and genetic correlations) for body weight and length of chinook salmon after 9 and 22 months of saltwater rearing. *Aquaculture*, 125:31-36.
- Withler, R. E., Beacham, T. D., Solar, I. I. And Donaldson, E. M. (1995): Freshwater growth, smolting, and marine survival and growth of diploid and triploid coho salmon (*Oncorhynchus kisutch*). *Aquaculture*, 136:91-107.
- Wohlfarth, G. W. (1994): The unexploited potential of tilapia hybrids in aquaculture. *Aquacult. Fish. Manage.*, 25:781-788.
- Wohlfarth, G. W., Hulata, G. (1987): Use of manures in aquaculture. In: D. J. W. Moriarty and R. S. V. Pullin (Editors), *Detritus and microbial Ecology in aquaculture*. ICLARM, Manila, pp.353-367.
- Wohlfarth, G. W., Hulata, G., Rothbard, S., Itzkowich, J. and Halevy, A. (1983): Comparisons between interspecific tilapia hybrids for some production traits. In L.

- Fishelson and Z. Yaron (Editors), International Symposium on Tilapia in Aquaculture, Proceeding, 559-569. Nazareth, Israel.
- Wolters, W. R., Chrisman, C. L. and Libey, G. S. (1982): Effect of triploidy on growth and gonadal development of channel catfish. *Trans. Am. Soc.*, 111:853-863.
- Woo, N. Y. S. and Chung, K. C. (1995): Tolerance of *Pomacanthus imperator* to hypoosmotic salinities: changes in body composition and hepatic enzyme activities. *J. Fish Biol.*, 47:70-81.
- Woynarovich, E. (1975): Elementary guide to fish culture in Nepal. FAO, Rome:133 pp.
- Yamada, R. (1986): Pond production systems: fertilization practices in warmwater. In: J. E. Lannan, R. O. Smitherman and G. Tchobanoglous (Editors), principles and practices of pond aquaculture, Oregon state University press, Corvallis, OR, pp. 85-96.
- Yi, Y., Lin, C. K. and Diana, J. S. (1996): Influence of Nile tilapia (*Oreochromis niloticus*) stocking density in cages on their growth and yield in cages and in ponds containing the cages. *Aquaculture*, 146:205-215.
- Yousif, O. M. (1996): Effects of rearing conditions in closed recirculatory systems on growth, feed utilization, carcass composition and nutrient digestibility of juvenile tilapia (*Oreochromis niloticus* L). Ph.D. thesis, Institute of Animal physiology and Animal Nutrition. University of Göttingen, Germany.
- Zamal, H. and Ollevier, F. (1995): Effect of feeding and lack of food on the growth biochemical and fatty acid composition of juvenile catfish. *J. Fish Biol.*, 46:404-414.
- Zeitler, M. H., Kirchgessner, M. and Schwarz, F. J. (1984): Effect of different proteins and energy supplies on carcass composition of carp (*Cyprinus carpio* L.). *Aquaculture*, 36:37-48.
- Zhu, Y., Yang, Y., Wang, J., Hua, D. and Mathias, J. A. (1990): The effect of manure application rate and frequency upon fish yield in integrated fish farm ponds. *Aquaculture*, 91:233-251.
- Zonneveld, N. and Fadholi, R. (1991): Feed intake and growth of red tilapia at different stocking densities in ponds in Indonesia. *Aquaculture*, 99:83-94.