

REFERENCES

6. REFERENCES

- Abd El-Gawad, A.A.; Gado, M.S.; Abdel-Karim, S.H. and Mahmoud, A.K. (1985).** Effect of protein on certain blood parameters, slaughtering and carcass characters. *Annals of Agric. Sci., Moushtohor.*, 23 : 115-121.
- Abd El-Hakim, N.F.; Attia, F.M.; Amer, A.A. and Younis, T.M. (1982).** Influence of dietary energy source on blood, liver, egg yolk cholesterol, total lipids and liver weight of growing and laying chicks. *Al-Azhar agricultural research bulletin, Faculty of Agriculture, Al-Azhar Univ. Bulletin No.4* .
- Abd El-Hamed, A. M. (1988).** Benefit from agriculture industrialization by-products in rumen animal nutrition. The first national conference on "the role of the scientific research in enhancing the animal wealth" the second forum, developing fodder sources for rumen. Scientific Research Academy. The International Egyptian Centre for Agriculture 25-29 Sep.
- Abd El-Hamed, A.M. and Gabr, A.A. (1991).** Evaluation of water hyacinth as feed for ruminants. *Arch. Anim. Nutr., Berlin*, 41: 745-756.
- Abd El-Khabir, A.M. (1978).** Studies on the feeding quality of water hyacinth with sheep. M.Sc. Thesis, Fac. of Agric., Cairo University.
- Abd El-Malak, N.Y; Abdel-Malak, M.S; El-Gendi, G.M. and Emily, F.N. (1995).** Effect of feeding different levels of herbal feed additive on broiler performance in relation to some metabolic functions. *Egyptian Poultry Sci.*, 15: 111-139.

- Abd Elmoty, A.K.I; El-Bogdady, A.H. and Faltas, A.A. (1986).** Performance and physiological responses of growing LSL chickens fed diets substituted with dried poultry manure. *Egyptian Poultry Sci.*, 6:21-29
- Abd El-Samie, R.E.; Ranaweera, K.N.P. and Nano, W.E. (1983).** The influence of fiber content and physical texture of the diet on the performance of broilers in the tropics. *Bri. Poultry Sci.*, 24: 383-390.
- Abou El-Wafa, S.M. (1994).** Influence of energy preparations and growth promoters on broiler performance. Ph.D. Thesis, Fac. Agric., Cairo University.
- Abou-Raya, A.K.; El-Talty, Y.; Hathout, M.K. and Abdel-Khabir, A.M. (1980).** Utilization of water hyacinth as animal feed: 2-Evaluation of wilted shoots in metabolism trials with sheep with reference to N, Ca, Na and K balances. *Agric. Res. Rev. (Anim. Prod.)* 58 (6): 49-59.
- Abou-Raya, A.K. and Galal, A.Gh. (1971).** Evaluation of poultry feeds in digestion trails with performance to some factors involved. *A.R.E. J. Animal Prod.*, 11: 207-221.
- Adele, M.; Giuseppe, P.; Raffaella, G. and Guglielmo, F. (1992).** Determination of blood constituents, reference values in broilers. *Poultry Sci.*, 71: 337-345.
- Ademosum, A.A. (1973).** Evaluation of brewers dried grains in the diets of growing chickens. *British Poultry Sci.*, 14: 463-468.
- Agroupis, F.M. (1953).** The value of water hyacinth as silage. *Philippine Agriculture J.*, 37 : 50-56.

- Ali, M.A. and Leeson, S. (1994).** Nutritional value and utilization of aquatic weeds in the diet of poultry. *World's Poult. Sci. J.*, 50: 237-251.
- Alicate, M.L.; Leto, G.; Giaccone, P. and Bononno, A. (1992).** Further trials on the use of chick-peas in growing rabbit feeding. *J. Appl. Rabbit. Res.*, 2:1025-1032.
- Asker, N.E.; Hermes, I.H.; Hamza, A.S. and Darwish, A.A. (1988).** Effect of dietary protein on uric acid content in the tissues. Thirteen International Congress for Statistics, Computer Science, Social and Demographic Research.
- Association of Official Analytical Chemists, AOAC (1990).** "Official Methods of Analysis". 15th Ed. Published by the AOAC., Washington, D.C.
- Bagnall, L.O.; Furman, T.Des.; Hentges, J.F., Jr.; Nolan, W.J. and Shirley, R.L. (1974).** Feed and fiber from effluent-grown water hyacinth. Proceedings on waste water use in the production of food and fiber. Oklahoma City, Oklahoma, March 5-7, pp. 116-141.
- Bagnall, L.O., Hentges, J.F.Jr. and Shirley, R.L. (1973).** Aquatic weed utilization-harvesting and processing. *Proc. Weed Sci., Society.* Atlanta, Georgia (C.F. Baldwin et al. 1975).

- Baldwin, J.A. (1973).** Utilization of ensiled water hyacinth in ruminal diets. M.Sc. Thesis., Agric., Florida Univ., Gainesville, U.S.A (C.F. Becker et al., 1987).
- Baldwin, J.A., Hentges, J.F.Jr. and Bagnall, L.O. (1974).** Preservation and cattle acceptability of water hyacinth silage. *Hyacinth Control J.*, 12: 79.
- Baldwin, J.A., Hentges, J.F.Jr., Bagnall, L.O. and Shirley, R.L. (1975).** Comparison of pangola grass and water hyacinth silages as diets for sheep. *J. of Anim. Sci.*, 40 : 968-971.
- Balnove, D. (1971).** The influence of exogenous estrogens and the attainment of sexual maturity on fatty acid metabolism in the immature pullet. *Com. Biochem. Physiol.*, 29:539.
- Batanouny, K.H. and El-Fiky, A.M. (1975).** The water hyacinth (*Eichhornia Crassipes Solms*) in the Nile system. *Egypt-Aquat. Bat.*, 1: 243.
- Bates, J.F. (1973).** Physical characteristics of aquatic plants. "Workshop on Aquatic Weed Management and Utilization. Some Prospects for Aquatic Weed Management in Guyana. (C.F. El-Serafy et al., 1980a).
- Bath, D., Dunlar, J., King, J., Berry, S., Leonoard, R.O. and Obrich, S. (1985).** Composition of by-products and unusual feedstuffs. *Feedstuffs*, 57 (29) : 32-36.

- Becker, R.B. (1946).** Use of dried citrus pulp in chickens diets. Circ. Flo. Univ. Agric. Exp. Stn., No. 423 (cited from El-Moghazy and El Boushy, 1982).
- Becker, K., Mahler, F. and Gall, C. (1987).** Water hyacinth in animal nutrition: The present situation and new research approaches. Animal Research and Develop., 25: 45-65.
- Bell, D.J. and Sturkie, P.D. (1965).** In " Avian physiology " 2nd ed. (P.D. Sturkie, Eds.). Ithaca. N.Y., Cornell University Press, Chapter 2.
- Bell, J.M. and Wilson A.G. (1970).** An evaluation of field peas as a protein and energy source for swine rations. Can. J. Anim. Sci., 50: 15-23.
- Bell, J.M. and Youngs, C.G. (1970).** Studies with mice on the nutritional value of pea protein concentrate. Can. J. Anim. Sci., 50: 219-226.
- Bertschart, A.A. and Kinsella, J.E. (1973).** Extractability and solubility of leaf protein. J. Agric. Fd. Chem., 21:60.
- Best, E.P.H. (1977).** Seasonal changes in mineral and organic components of *Ceratophyllum demersum* and *Elodea canadensis*. Aquatic Botany, 3: 337-348.
- Bhatty, R.S. and Christison, G.I. (1980).** Digestibility of pea proteins by pre-ruminant calves. Can., J. of Anim. Sci., 60: 925-930.

- Biswas, M. and Mandal, L. (1987).** Use of wilted water hyacinth in the ration of growing calves. *Indian J. Anim. Sci.*, 57: 482-485.
- Biswas, M. and Mandal, L. (1988).** Use of fresh water hyacinth in the ration of growing calves. *Indian Vet. J.*, 65: 49-50.
- Block, R.J. and Mitchell, H.H. (1946).** The correlation of amino acid composition of the protein with their nutritive value. *Nutr. Abs. Rev.*, 16: 249.
- Bock, J.H. (1960).** An ecological study of *Eichhornia crassipes* with special emphasis on its reproductive biology. Berkely, Univ. of California, Thesis (C.F. El Serafy et al., 1980a).
- Bock, J.H. (1969).** Productivity of the water hyacinth [*Eichhornia crassipes* (Mart.) Solms]. *Ecology*, 50:460.
- Bonomi, A., Catalano, A.L. and Quarantelli, A. (1985).** Chemical composition and digestibility in vivo of dried pea fodder. *Annali della Facolta di Medicina Veterioria Universite di Parma*, 5: 39-46. (Nut. Abs. and Rev. 57, 12, 1987).
- Bonomi, A., Quarantelli, A., Superchi, P., Sabbioni, A. and Malavasi, P. (1981).** Chemical and nutrient composition of water hyacinth (*Eichhornia crassipes*) and duck weed (*Lemna minor*) grown on partly treated liquid refuse from pig houses. *Annali della Facolta di Parma* 1: 198-216. (C.F. Becker et al., 1987).

Boyd, C.E. (1968a). Fresh-water plants, a potential source of protein.

Economic Botany, 22: 359-368.

Boyd, C.E. (1968b). Evaluation of some aquatic weeds as possible

feedstuffs. Hyacinth Control J., 7: 26.

Boyd, C.E. (1969). The nutritive value of three species of water weeds.

Economic Botany, 23: 123-127.

Boyd, C.E. (1974). Utilization of aquatic plants. In: Aquatic Vegetation

and its Use and Control (Ed. Mitchell, D.S.), United Nations

Education, Scientific and Cultural Organization, Paris.

Boyd, C.E. (1976). Accumulation of dry matter, nitrogen and phosphorus.

Economic Botany, 30: 51-56.

Boyd, C.E. and Blackburn, R.D. (1970). Seasonal changes in the

proximate composition of some common aquatic weeds. Hyacinth

Control J., 8(2): 42-44.

Boyd, C.E. and McGinty, P.S. (1981). Percentage digestible dry-matter

and crude protein in dried aquatic weeds. Economic Botany, 35(3);

296-299.

Boyd, C.E. and Vickers, D.H. (1971). Variation in the elemental content

of *Eichhornia crassipes*. Hydrobiologia, 38: 409-414.

Buriel, P.; Luqo, C.M. and Mata, R.O. (1976). Note on citrus meal in

diets for growing chickens. Agronomia Tropical, 26(3): 216-266.

- Burkill, I.H. (1966).** Eichhornia Crassipes-Utilization. A Dictionary of Economic Products of the Malay Peninsula, 1: 906 "C.F. Soesiawaningini et al., 1977".
- Chatterjee, I. and Hye, M.A. (1938).** Can water hyacinth be used as a cattle feed. Agric. Live-stock, India, 8: 547-553." C.F. Little, 1968".
- Chhibbar, S.S. and Singh, G.D. (1971).** Paddy straw and water hyacinth silage. Indian Farming, 20 (11): 24-28.
- Choy, T.L. and Devaraj, R.M. (1958).** A study of comparative feeding methods on local pigs. J. Malay. Vet. Med. Ass. II, 49" C.F. Soesiawaningini et al., 1977".
- Combs, G.E. (1970).** Aquatic plants for swine feeding. Proceedings of the Aquatic Plant Research., Conference University of Florida, Gainesville, Florida (USA, 60-61). "C.F Baldwin et al., 1975".
- Combs, G.E. (1973).** Processed aquatic plants for animal nutrition. Annual Research., Report of the Institute of Food and Agriculture of Florida, Gainesville, Florida (USA) 81.
- Combs, G.E. and Wallace, H.D. (1973).** Use of dehydrated water hyacinth in swine diets. Annual Research., Report of the Institute of Food and Agriculture of Florida, Gainesville, Florida (USA-No.AL) 73-3.
- Creger, C.R., Farr, F.M., Castro, E. and Couch, J.R. (1963).** The pigmenting value of aquatic flowering plants. Poultry Sci., 42: 1262.

- Damron, B.L. and Wilson, H.R. (1983).** Geese found highly effective in controlling water hyacinths. *Feedstuffs*, 31: 15-17.
- Davies, H.R.J. (1959).** Effect of water hyacinth (*Eichhornia crassipes*) in the Nile Valley. *Nature*, 184: 1085.
- Day, F.W.F. (1918).** The water hyacinth as source of potash. *Agric. Bull. F.M.S.*, 6: 309. In: handbook of utilization of aquatic plants a compilation of the world's publications, 1968. FAO. P1: Cp/20, Rome, Italy.
- Devendra, C. (1976).** The value of the lehmann method. *World Review of Animal Production*, 12(2): 39-42.
- Devendra, C. and Gohl, B.I. (1970).** Performance of broiler fed on diets containing citrus pulp. *Trop. Agric., Trin.*, 47(4): 335. "CF.El-Moghazy and El-Boushy (1982)
- Dessouky, M.S. (1997).** Nutritional studies on rape seeds and sunflower seeds as feedstuffs in broiler rations. Ph.D. Thesis, Fac. of Agric., Al-Azhar Univ.
- Dhar, N.R. (1963).** Organic matter and phosphates in land fertility increase. *Proc. Nat. Acad. Sci., India (B)* 33(1): 16-24. "C.F. Little, 1968".
-

- Dirven, J.G.P. (1965).** The protein content of Surinam roughages. *Qualitas Plantarum et Materiae Vegetabiles*, 12: 172-184. In: handbook of utilization of aquatic plants a compilation of the world's publications, 1968, FAO. P1: CP/20, Rome, Italy.
- Dolberg, F., Saadullah, M. and Haque, M. (1981).** Nutrient elements for livestock in aquatic plants. *Trop. Animal Production*, 6: 322-326.
- “C.F. Wana pat, M., 1983”
- Doumas, B.T. (1971).** Automated analysis boehringer mannheim. *Clin. Chem. Acta.*, 31: 87.
- Driggers, J.C. (1951).** Crop production in dry regions. Florida Agricultural Experiment Station. Bulletin No. 476. “CF.El-Boushy and Vanderpol 1994”
- Duncan, D.B. (1955).** Multiple Range and Multiple F tests. *Biometrics*, 11: 1-42.
- Dutta, R.K., Sahu, B.K., Panda, N.C. and Nayak, B.C. (1984).** Toxic effects of feeding water hyacinth to goats. *Indian. J. Animal. Sci.*, 54(6): 594-597.
- Eggum, B.O. (1970).** The protein quality of cassava leaves. *Br. J. Nutr.*, 24: 761.
- Ekman, P., Enianuelson, S. and Fransson, A. (1949).** Estimation of faecal protein. *K. Lanthbrukse Ann.*, 16: 749.
-

- El-Alaily, H.A. (1974).** Feed for poultry from food processing wastage.
Ph.D. Thesis, Fac. Agric., Ain Shams University.
- El-Bogdady, A.H., Faltas, A.A. and Abd-Elmoty, A.K.I. (1987).**
Effects of starvation and transport on some physiological parameters
of LSL chickens. *Egyptian Poultry Sci.*, 7: 26-36.
- El-Boushy, A.R. and Vanderpol, A.F. (1994).** Poultry feed from waste:
Processing and use .Chapman and Hall, London.(305-320)
- El-Faham, A.I., Ibrahim, S.A. and El-Wardany, I.E. (1995).** Effect of
switching yellow corn with cassava meal on broiler performance,
carcass characteristics and some blood constitutes.*Egyptian Poultry
Sci.*, 15: 261-283.
- El- Gendi, G.M. (1996).** Effect of feeding dietary herbal feed additives on
productive and metabolic responses of broiler chicks. *Egypt. Poult.
Sci*, 16(2): 395- 412
- El-Husseiny, O.M., Ghazalah, A.A., Fayek, H.M. and Abou-El-Wafa,
S.M. (1995).** Enzyme preparations, growth promoters and rumen
contents in broiler diets.*Egyptian Poultry Sci.*, 15: 205-231.
- El-Moghazy, S.A. and El-Boushy, A.R. (1982).** The effect of different
levels of dried citrus pulp in isocaloric isonitrogenous methionine
and lysine supplemented rations: 1-On the performance of broilers.
Ain Shams Univ., Fac. Agric., Research Bulletin 2035.
-

- El-Serafy, A.M.; Allam, S.M.; El-Ashry, M.A.; Kattab, H.M.; Soliman, H.S. and Swidan, F.Z. (1981).** Dry matter intake and nutritive digestibility of water hyacinth hay, haylage and silage by buffalo steers. *Indian J. Anim Sci.*, 51 (7): 698-701.
- Ewing, W.R. (1963).** "Poultry Nutrition". The Ray Ewing Company Publisher, Pasadena, California.
- FAO. (1991).** FAO Production Yearbook, Vol 44, Food and Agriculture Organization of the United Nations, Roma.
- Faris, F. (1972).** Studies on the ecology of *Eichhornia crassipes* (Mart.) Solms in the Sudan. Thesis, Univ., of Khartum, Mimeographed.
- Farrell, D.J., Cummming, R.B. and Hardaker, J.B. (1973).** The effects of dietary energy concentration on growth rate and conversion of energy to weight gain in broiler chickens. *British Poultry Sci.*, 14: 329-340.
- Finlow, R.S. and Mclean, K. (1917).** Water hyacinth and its value as a fertilizer. Bull. 71, Agric. Res. Inst. Puse. In: of utilization of aquatic plants a compilation of the world's publications 1968. FAO. P1: CP/20, Rome, Italy."C.F. Little, 1968".
- Gado, M.S., El-Aggoury, S.A., Radwan, A.A and El-Banayousi, W. L. (1984).** Seasonal variation in some serum constituents for Lohmann broiler chicks of light and heavy hatching weights. *Egyptian Poultry Sci.*, 15:381-390.
-

- Gay, P.A. (1958).** *Eichhornia crassipes* in the Nile of Sudan. *Nature*, 182: 528.
- Gay, P.A. (1960).** Ecological studies of *Eichhornia crassipes* solms in Sudan. *J. Ecol.*, 48: 183-191.
- Gay, P.A. and Berry, L. (1959).** The water hyacinth: a new problem in the Nile. *Geol. J.*, 125: 89.
- Gerard, C. and Troncoso, J. (1980).** Utilization de la jacinth d'eau (*Eichhornia crassipes*) par le lapin de chair. *Rev. Elev. Med. Vet. Pays. Trop.*, 33: 91-96. "C.F. Moreland et al, 1991".
- Gerloff, G.C., Moore, D.D. and Curtis, J.T. (1964).** Mineral content of native plants of Wisconsin. *Agri. Expt. Sta., Univ. Wisc., Madison Research Report 14* "C.F. Boyd and Vickers, 1971"
- Ghazalah, A.A. and El-Shahat, A.A. (1994).** Digestibility and acceptability of some agro-industrial by-products by rabbits. *Egypt. Poult. Sci.*, 14:401-424.
- Ghoneim, A. (1967).** Feed requirements and economical rations. *Anglo Egypt. Lib., Cairo (Textbook in Arabic).*
- Gohl, B. (1981).** Tropical feeds. *FAO Animal Production and Health Series No. 12.* Food and Agriculture organization of the United Nations, Rome, Italy.

- Gopal, B. and Sharma, K. P. (1981).** Water hyacinth (*Eichhornia crassipes*), the most troublesome aquatic weed of the world. Hindosid Publisher (India)
- Gossett, D.R. and Norris, W.E. Jr. (1971).** Relation between nutrient availability and content of nitrogen and phosphorous in tissues of the aquatic macrophyte, water hyacinth. *Hydrobiologia*, 38 (1); 15-28.
- Grandi, A. (1981).** Use of water hyacinth (*Eichhornia crassipes*) in diets for rabbits. *Conigliculture*, 18 (6): 43-48.
- Grandi, A. and Angelis, A. (1983).** Use of agriculture and industrial products for feeding animals. Waste from processing of peas (*Pisum sativum* L.) for feeding fattening rabbits. *Conigliculture*, 20 (3): 45-48.
- Grandi, A. and Marzetti, P.(1981).** Digestibility and nutritive value to the rabbit of two types of water hyacinth. *Conigliculture*, 18 (12): 29-32.
- Grandi, A.; Marzetti, P. and Blasi, F. (1984).** Research on the use of water hyacinth (*Eichhornia crassipes*) flour in the feeding of broilers. *Zoot. Nutr. Anim.*, 10: 229-307. (C.F. Nutr. Abst & Rev., 55: 861).
- Gruhn, k. and Valdivie, D.A. (1982/1983).** Digestibility of the crude nutrients in peas for laying hens. *Tiernahrung and Fulleuing*, 13: 147-150, (C.F. Nutr. Abst. and Rev. 54(5):82,1984.

Gupta, B.N. (1981).Utilization of water hyacinth (*Eichhornia crassipes*) as a source of nutrients for animals. Indian J. Nutr. Dietetics, 18(12): 435-441.

Haller, W.T., Knipling, E.B. and West, S.H. (1970). Phosphorus uptake and distribution in water hyacinth. Proceedings: Soil and Crop Science Society of Florida, 30: 64-68.

Haller, W.T. and Sutton, D.L. (1973). Effect of pH and high phosphorus concentrations on growth of water hyacinth. Hyacinth Control J., 11: 59-61.

Hamdy, S.H.; Khalifah, M.M.; Khalil, A.Z. and Kosba, M.A.. (1978). Effect of corn and bean replacement by dried water hyacinth on the performance of laying hens. Alex. J. Agric. Res., 26 (3): 515-525.

Harms, R.H. (1968). Feed information summaries and nutritive values. Florida Agriculture Experiment Stations-Bulletin No 724.

Hathout, M.K., Abou-Raya, A.K., El-Nouby, H.M., El-Talty, Y.I. and Abdel-Khabir, A.M. (1980).Utilization of water hyacinth as animal feed. Agric. Res. Rev., 58(6): 33-47.

Hbbell, C.H. (1969)."Analysis table for feed ingredients". Feedstuffs, 6:121-128

Hentges, J.F. Jr. (1970).Processed aquatic weeds for cattle nutrition- Proc. Aquatic Plant Res. Conf. Univ. of Florida, Gainesville" C.F. Becker, 1987".

- Hermes, I.H.; Asker, N.E. and Hamza, A.S. (1988).** A comparative study between some protein concentrates used in broiler feeding: 1. Productive performance. Thirteen International Congress for Statistics, Computer Science, Social and Demographic Research.
- Hill, F.W. and Dansky, L.M. (1954).** Studies on the energy requirements of chickens.1. The effect of dietary energy level on growth and feed consumption. Poultry Sci., 33: 112-119.
- Holm, L.G., Weldon, L.W. and Blackburn, R.D. (1969).** The world's worst weeds. Aquatic weeds Sci., 116: 699-709.(C.F. Shoukry, 1982).
- Hora, S.L. (1951).** The water hyacinth problem and pig farming. Science and Culture, 17(6): 231-232.
- Horszczaruk, F. (1976).** Ruminant nutrition in the near east: A practical guide. Near East Animal Production and Health Development Center-Food and Agriculture Organization of the United Nations-United Nations Development Programmer- Handbook No.12(1-10)-Bagdad,
- Hossain, W. (1959).** Investigation on water hyacinth as fodder.Agric. Pakistan, 10(4): 513-518.
- Hussien, A.A. (1994).**The effect of some modifications in managerial and environmental condition on meat production of chickens. M.Sc. Thesis, Fac. Agric., Cairo, University.

- Ibrahim, K.A. (1995).** The value of unusual high protein cereal (gerber) as a replacement for broiler rations ingredients. *Egyptian Poultry Sci.*, 15: 351-380.
- Igbasan, F.A. and Guenter, W. (1994).** Utilization of pea by-products in broiler starter diets. *Poultry Sci.*, 73: 94.
- Igbasan, F.A.; Guenter, W.; Slominski, B.A. and Zhang, W. (1994).** Alternative feed source for poultry diets: an evaluation of feed type pea cultivars from western Canada. *Poultry Sci.*, 73: 9.
- Kabak, A.N. (1993).** Effect of feeding water hyacinth at different levels on animal performance. M.Sc. Thesis, Fac. Agric., Assiut, Univ.
- Kashem, M.A.; Sen, K.; Das, U.; Islam, S.; Khan, N.M.; Huq, A.K.M.A. and Razzaque, A. (1984).** Management of water hyacinth: Utilization of water hyacinth for animal feed. *Proceedings of the International Conference on Water Hyacinth.* (Ed. Thyagarajan, G.), United Nations Environment Programme, Nairobi, pp., 379-387.
- Kazemi, R. and Dagher, A.M. (1971).** Interrelation ship between methionine and vitamin B₆ in chicks diets. *Poultry Sci.*, 50: 1269-1302.
- Kazemi, R. and Kratzer, F.H. (1980).** Source of protein affecting the vitamin B₆ requirement of chicks. *Poultry Sci.*, 59: 95-99.

- Kelley, E.G. (1958).** Plant residues and pomaces. In: Processed Plant Protein Foodstuffs, (Ed. Altschul, A.M) Academic Press Inc., Publishers, New York pp. 859-875. (C.F. El-Boushy and Vanderpol 1994)
- Khalil, A.A.; Khalifah, M.M.; Hamdy, S.M., and Kosba, M.A. (1975).** Dried water hyacinth as a partial substitute for corn in a chick starter diet. Alex., J. Agric., Res., 3 (1): 47-51.
- Khirwar, S.S., Paliwal, V.W. and Pradhan, K. (1980).** Utilization of agro industrial by products by livestock and poultry. 1. Effect of malt sprout and pea waste on broiler chicks. Haryana Agric. Univ. J. of Res., 10 (1): 101-109.
- Kiflewahid, B. (1975).** Nährstoffzusammensetzung und Verdaulichkeit von Wasserhyazin. "C.F. Becker et al., 1987."
- Kiflewahid, B. (1978).** Nutrient composition and digestibility of water hyacinth (*Eichhornia crassipes*, Mart) by cattle. Herbage Abstr., 48: 459.
- Knab, W. (1982).** Untersuchungen über den Futterwert der Wasserhyazinthe (*Eichhornia crassipes* (Mart) Solms) bei Wiederkäuern. Diplomarbeit, Univ., Hohenheim "C.F. Becker et al., 1987".
- Knipling, E.B.; West, S.H. and Haller, W.T. (1970).** Growth characteristics, yield potential and nutritive content of water hyacinth. Soil and Crop Sci., Soc. Of Florida, Proc., 30: 57-63 (Herbage Abstr., 44: 16, 1974).

- Lareo, L. and Bressani, R. (1982).** Possible utilization of the water hyacinth in nutrition and industry. Food and Nutrition Bulletin, 4: 60-64.
- Liang, J.K. and Lovell, R.T. (1971).** Nutritional value of water hyacinth in channel catfish feeds. Hyacinth Control J., 9 (1): 41-44.
- Linn, J.C.; Boodrich, R.D.; Otterly, D.E.; Meiske, T.C. and Staba, E.J. (1975).** Nutritive value of dried or ensiled aquatic plants: Digestibility by sheep. J. Animal Sci., 41: 612.
- Little, E.C.S. (1968).** "Handbook of Utilization of Aquatic Plants." FAO the UN. Rome. Italy.
- Loosli, J.K.; Belmonte, R.P.; Villegas, V. and Cruz, E. (1954).** The digestibility of water hyacinth (*Eichhornia crassipes*) silage by sheep. Philippine Agric. J., 38: 146.
- Lopez, P.L. and Mania, R.P. (1982).** The potential of leaf protein concentrate as supplement in broiler ration. Animal. Production and Health in the Tropics, pp. 223-227.
- Lopez, P.L.; Mania, R.R.; Beldia, M.D.; Trinidad, E.B. and Esmade, J.R. (1981).** The potential of leaf protein concentrate as supplement in broiler ration. Philippine Journal. of Veterinary and Animal Sciences., 7 (1) : 38-39.

- Lyman, C.M. (1956).** Nutritive value of citrus pulp. *J. Agric. Fd. Chem.*, 4: 1008. "C.F. El-Boushy and Vanderpol, 1994"
- Maertens, L. Moermans, R. and DeGrotte, G. (1988).** Predication of the apparent digestible energy (ADE) content of commercial pelleted feeds for rabbits. *J. Appl. Rabbit Res.*, 11: 60-67.
- Mahendranathan, T. (1970).** Water hyacinth has value as a pig feed. *Pig Farmer*, 599-607.
- Mahmud, E.A.B. (1967).** The use of water hyacinth in animal feeding (Pigs) a review. *Handbook of Utilization of Aquatic Plants*, 49-50.
- Mitra, E. and Banerjee, A.O. (1971).** Eradication of Eichhornia plants from fish ponds by bottom soil treatment. *Proceedings of the National Academy of Sciences., (India)* 41, (3), 231-236.
- Moreland, A. F.; Collins, B.R.; Hansen, C.A. and Obrien, R. (1991).** Wastewater grown water hyacinth as an ingredient in rabbit food. *J. Aquat. Plant Manage.*, 29: 32-39.
- Morgan, G.W., Jr. and Glick, B. (1972).** A quantitative study of serum proteins in bursectomized and irradiated chickens. *Poultry Sci.*, 51: 771.
- Mousa, S.M. and Ahmed, S.A. (1992).** Chemical composition of water hyacinth; An introductory study as ruminant foodstuff. *National Symposium on Water Hyacinth, Assiut Univ.* 25-26, Feb.pp, 50-70
-

- Mudgal, V.D. and Singh, N. (1983).** Water hyacinth as animal feed. Indian Farming, 33 (3): 33-36.
- Mukhtar, A.M. S. (1967).** A preliminary study on the chemical composition of water hyacinth in the Sudan. Sudan J. Vet. Sci., and Animal Husbandry, 8 (2): 124-126.
- Muztar, A.J.; Slinger, S.J. and Burton, J.H. (1978).** Chemical composition of aquatic macrophytes III. Mineral composition of fresh water macrophytes and their potential for mineral nutrient removal from lake water. Canadian J. of Poult. Sci., 58: 851-862.
- National Academy of Sciences, NAS (1976).** Making aquatic weeds useful: Some perspectives for developing countries. National Academy of Sciences, Washington, USA, National Academy Press.
- National Research Council, NRC (1971).** Nutrient Requirements of Poultry- 6th Ed. Acad. Sci., Washington, D.C., USA.
- National Research Council, NRC (1977).** Nutrient Requirements of Poultry- 7th Ed. Acad. Sci., Washington, D.C., USA.
- National Research Council, NRC (1984).** Nutrient Requirements of Poultry- 8th Ed. Acad. Sci., Washington, D.C., USA.
- National Research Council, NRC (1994).** Nutrient Requirements of Poultry- 9th Ed. Acad. Sci., Washington, D.C., USA.
- Neumark, H. (1970).** Volcani Institute of Agricultural Research, Israel. In: Handbook Tropical Feeds. FAO Animal Production and Health,

Series, 12. Food and Agriculture Organization of the United Nations, Rome.

Nour, A. M. (1988). Benefit from agriculture industrialization by-products in rumen animal nutrition. The first national conference on "the role of the scientific research in enhancing the animal wealth" the second forum, developing fodder sources for rumen. Scientific Research Academy. The International Egyptian Center for Agriculture 25-29 Sep.

Nour, A.M.; El-Shazly, K.; Abo Akkada, A.R.; Borhami, B.E. and Abaza, M.A. (1980). Evaluation of silage of some by-products from food processing industry. Alex. J. Agric. Res., 28(1): 17-24.

NSRC, Guyana and NAS, U.S.A. (1973). "Workshop on: Aquatic weed management and utilization. Some prospects from aquatic weed management in Guyana". National Academy of Sciences, USA.

Obara, J.; Nyunt, K.M. and Tun, S. (1974). Chemical composition of some green feeds available in Burma for chicken. Nippon Chickusan Gakkai-Ho, 45: 216-217.

Obeid, M. and Tag El-Seed, M. (1973). The water hyacinth (*Eichhornia crassipes* "Mart" Solms) in the Sudanese Nile system. Proc. First Conf. of Pests Control, Univ. of Assiut, Egypt.

Osman, H.E.; El-Hag, G.A. and Osman, M.M. (1975). Studies on the nutritive value of water hyacinth (*Eichhornia crassipes* "Mart" Solms). In: Aquatic weeds in the Sudan with

special reference to water hyacinth. National Council For Research, Khartoum., Sudan: 104-127.

Paliwal, V.W.; Khirwar, S.S. and Pradhan, K. (1980). Utilization of agro-industrial by-products by livestock and poultry.2. Effect of higher levels of pea waste on the performance of broiler chicks. Haryana Agric. Univ., J. Res., 10 (2): 283-288.

Pathak, N.N., Srivastava, S.K. and Ranjhan, S.K. (1980). Note on the nutritive value of water hyacinth (*Eichhorina crassipes*) for pigs. Indian J. Anim. Sci., 50(10): 886-888.

Penfound, W.T. and Earle, T.T. (1948). The biology of water hyacinth. Ecology Monographs, 18: 447-472. "C.F. Little, 1968"

Percheron, D. (1903). La Jacinthe d'Eau. Bull. Union Synd. Agric. D'Egypt, 3^{eme} Ann. No. 30, Alexandria. (C.F. Batanouny and El-Fiky, 1975).

Pescatore, A.J.; Cantor, A.H.; Jackson, K.; Johnson, T.H. and Pfoff, W.K. (1990). Influence of barley-based diets on egg cholesterol content and production of two strains of laying hens. Poultry Sci., 69 supplement 1: 183.

Pfizer, A.D. (1981). Feed formulator. In: Tropical Feeds. FAO Animal Production and Health Series. No. 12. Food and Agriculture Organization of the United Nations, Rome.

- Philipp, O. (1981).** Zur Verwertung der Wasserhyazinthe (*Eichhornia crassipes*; "Mart" Solms). Diss. Univ. Hohenheim. (C.F. Becker et al., 1987).
- Philipp, O.; Koch, W. and Koser, H. (1983).** Utilization and control of water hyacinth in Sudan. Schriftenreihe der GTZ Nr. 122. Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ) GmbH, Eschborn.
- Philip, Y.L. (1996).** Nutritional studies for improving the utilization of water hyacinth by ruminants. M.Sc. Thesis, Fac. Agric., Cairo Univ.
- Pieterse, A.H. (1978).** The water hyacinth (*Eichhornia crassipes*). A review. Abst. of Tropical Agriculture, 4 (2): 9-24.
- Pirie, N. W. (1960).** Water hyacinth: a curse or a crop. In: Handbook of Utilization of Aquatic Plants: A compilation of the World's Publications, 1968. E.C.S. Little ed. (FAO. PL. CP/20).
- Pirie, N.W. (1967).** The present position of research on the use of water hyacinth. In Handbook of Utilization of Aquatic Plants: A compilation of the World's Publications, 1968, E.C.S. Little ed. (FAO. PL. CP/20).
- Pirie, N.W. (1971).** Leaf protein: Its agronomy, preparation, quality and use. In: Pirie, N.W., Ed. IBP Handbook No. 20. Blackwell Scientific Publications. London. pp. 1-4.
-

- Qota, E.M.A. (1994).** Effect of feeding system on poultry productivity under Egyptian farm conditions. M.Sc. Thesis, Fac. Agri., Mansoura Univ.
- Quarantelli, A.; Lucchelli, L.; Anghinetti, A. and Blanco, P. (1987).** Use of blood meal in the feeding of broiler chickens (Experimental contribution). *Amali Della Facolta Di Medicine Veterinari A Univertia Di Parma*.(7):333-353
- Ravindran, V. and Blair, R. (1992).** Feed resources for poultry production in Asia and the Pacific. II. Plant protein sources. *World's Poultry Sci. J.*, 48: 205-231.
- Reddy, K.R. (1984).** Water hyacinth (*Eichhorina crassipes*) biomass production in Florida. *Biomass*, 6 (1/2) 167-181.
- Reddy, K.R.; Hueston, F.M. and Mckim, T. (1985).** Biomass production and nutrient removal potential of water hyacinth cultured in sewage effluent. *Engineering J. Sol. Energ. Engineering*, 107 (2): 128-135.
- Reddy, M.R. and Mohanrao, A.V. (1979).** Studies on supplementation of water hyacinth hay to paddy straw for improved utilization by growing cross-bred calves. *Indian J. Animal Sci.*, 49 (8): 618-621.
-

- Reddy, P.V.S. and Reddy, M.R. (1979).** Utilization of water hyacinth meal (*Eichhorina crassipes*) in the concentrate feeds of cross bred calves. *Indian J. Animal Sci.*, 49(3): 174-179.
- Reduas, B., Wich, G. and Cole, R.K. (1972).** Serum lipid pattern in chickens of the obese strain. *J. Endocrinol.*, 55: 609.
- Reitman, S. and Frankel, S. (1957).** Test combination of glutamic-pyruvic transaminase (GPT) and glutamic-oxaloacetic transaminase (GOT). *Amer. J. Clin. Path.*, 28: 56.
- Reza, A. and Khan, J. (1981).** Water hyacinth as cattle feed. *Indian J. Anim. Sci.*, 51 (7): 702-706.
- Riemer, D.N. and Toth, S.J. (1968).** Survey of chemical composition of plants in New Jersey. *New Jersey Agr. Expt. Sta. Bull.* 820. "C.F. Boyd and Vickers, 1971".
- Ross, E. (1971).** Biological control of pond weeds with white Chinese Geese. *Hawaii Farm Sci.*, 20 (2): 11-12.
- Salem, F. M.S.; El-Shennau, M.M. and Ali, F.R. (1983).** Evaluation of "butaphosphan" as food additive used in improvement of poultry production. *Vet. Med. J.* Vol. 31, No. 2.
- Salveson, R.E. (1971).** Utilization of aquatic plants in steer diets: Voluntary intake and digestibility. M. Sc. Thesis, Univ. of Florida. "C.F. Bagnall et al, 1974".
-

- Santos, E.A. (1982).** Water hyacinth as feed replacement of commercial broiler mash. CLSU (Central Luzon State University) Scintific J, V. 5 (2) and 6 (1) P. 97.
- Sayed, M.A. (1995).** Nutritional studies on shrimp meal in broiler diets. M. Sc. Thesis, Fac. Agric. Cairo Univ.
- Sayed, A.; Soliman, M. and El-Sayed, A.M. (1992).** Seasonal variations in certain nutrient contents of water hyacinth and associated molluscs in Assiut area. National Symp. On Water Hyacinth-Assiut Univ. (25-26 Feb.)
- Schaible. P.J. (1970).** Poultry Feeds and Nutrition. The Avi Publishing Company, Inc ., West Port Connecticut.
- Scott, M.L.; Nesheim, M.C. and Yong, R.J. (1982).** "Nutrition of chicken". Scott, M.L. and Associates, Ithaca, New York, pp. 562.
- Seroux, M. (1984).** Utilization des proteagiheux par le lapion a l'engraissement, Pois lupin fevarol. Proc., World Rabbit Conger., 3: 376-383.
- Sharma, A. (1971).** Eradication and utilization of water hyacinth. A Review. Curr. Sci., 40 (30): 51.
- Shoukry, M.M. (1982).** Optimum utilization of water hyacinth plants in feeding ruminants. Ph. D. Thesis, Fac. Agric. Ain Shams. Univ, Egypt.
-

- Sibbald, I.R.; Slinger, S.J. and Ashton, G.G. (1960).** The weight gain and feed intake of chicks fed a ration diluted with cellulose or kaolin. *J. of Nutrition*, 72: 441.
- Siedel, J. (1983).** Test combination of cholesterol. *Clin. Chem.*, 29: 1075.
- Simeonov, S.; Pashov, D.; Peinikova, Ts. and Dzhurov, A. (1975).** Chronic toxicity of zinc bacitracin in chickens. *Veter., Nauki*, 12 (2): 100-105.
- Smetana, P. (1967).** Report to the Government of Burma. FAO poultry expert., Burma (C.F. Little, 1968).
- Soesiawaningrini, D.P.; Soewardi, B. and Thohari, M. (1977).** Water hyacinth (*Eichhornia crassipes* (Mart.) Solms) in broiler rations. Proceedings of the 6 th Asian-Pacific Weed Science., Society Confernce, Jakarta, Indonesia, 2: 623-627, 1979.
- Soewardi, B. (1974).** On the utilization of water hyacinth (*Eichhornia crassipes*) by Holstein Friesian grade heifers. Proceedings of Southeast-Asian Workshop on Aquatic Weeds Malang . (C.F. Soesiawaningrini, 1977).
- Soliman, H.S.; Mousa, M. and Sayed, A.A. (1992).** Chemical composition of water hyacinth: An introductory study as ruminant feedstuff. National Symposium On Water Hyacinth- Assiut Univ., 25-26 Feb., pp. 55-69.

- Soliman, H.S., Shoukry, M.M., El-Serafy, A. and El-Sokkary, O.** (1987).Effect of grinding and urea or NaOH plus urea treatment on the utilization of water hyacinth hay by sheep.J. Agric., Sci., Mansoura Univ., 12 (4): 742-749.
- Solly, R.K.; Singh, D. and Sharma., V. (1984).**Water hyacinth as feed supplement for small animals.Proceedings of the International Conference on Water Hyacinth (Ed. Thyagarajan,G.) United Nations Environment Programme, Nairobi, pp.398-407.
- Sreemannarayana, O.; Nambiar, K.T.K. and Satchidanandam, V.** (1973). A note on the production of VFA by bovine rumen liquor in artificial rumen in the presence of cashew apple and water hyacinth meals. Indian J. of Animal Sci., 43(6): 545-546.
- Steel, R.G.D. and Torrie, J.E. (1980).** Principles and Procedures of Statistical Approach. 2nd Ed. McGraw Hill Book Co., Inc., New York.
- Stephens, E.L.; Easly, J.F.; Shirley, R.L. and Hentges, J.F.Jr. (1973).** Availability of nutrient, mineral elements and potential toxicants in aquatic plant diets fed steers.Soil and Crop Sci., Soc. of Florida Proc. 30: 32.
- Steward, K.K. (1970).** Nutrient removal potentials of various aquatic plants. Hyacinth Control J., 8: 34-35.
-

- Straskraba, M. (1966).** Das Anteil der höheren Pflanzen an der Produktion der Gewässer. Mith. Int. Ver. Theor. Ang Linnol. 14. Stoffhaushalt der Binnengewässer: Chemie und Mikrobiologie. "C.F. Boyd and Vickers, 1971".
- Sturkie, P.D. (1976).** "Avian Physiology", 3rd Edition.
- Sturkie, P.D. and Newman, H.J. (1951).** Plasma proteins of chickens as influenced by time of laying, ovulation, number of blood samples taken and plasma volume. Poultry Sci., 30: 240-248.
- Surat, R. and Singh, G.S. (1980).** Chemical composition and nutritive value of water hyacinth. Indian J. Dairy Sci., 33 (2): 165-167.
- Susiawaningrini, D.D.; Harijanto, E. and Thohari, M. (1977).** Sebagai Hijauan Makanan Ayam dan Babi. Laporan Akhin Nr. 6/1977. Hasil Penelitian. Masalah and Pengendalian Tumbuhair Pengganggu Air Rawa Pening. No. Doc. Biotrop/TP/77/247. In : Handbook-Utilization and Control of Water Hyacinth in Sudan.
- Sympson, N.D. (1932).** A report on the weed flora of the irrigation channels in Egypt. GOV Press, Cairo
- Tackholm, V. and Drar, M. (1950).** Flora of Egypt, Vol. II, Fuoad I Univ. Press, Cairo, pp. 547. (C.F. Batanouny and El Fiky, 1975)
- Taylor, K.G.; Bates, R.P. and Robbins, R.C. (1971).** Extraction of protein from water hyacinth. Hyacinth Control J., 9(10): 20-22.
-

- Taylor, K.G. and Robbins, R.C. (1968).** The amino acid composition of water hyacinth (*Eichhhornia crassipes*) and its value as a protein supplement. *Hyacinth Control J.*, 7: 24-25.
- Tucker, C.S. and DeBusk, T.A. (1981).** Seasonal growth of (*Eichhhornia crassipes* "Mart." Solms): relationship to protein, fiber and available carbohydrate content. *Aquatic. Bot.*, 11: 137.
- Udedibie, A.B.I., and Igwe, F. O. (1989).** Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens. *Animal Feed Science and Technology*, 24: 111-119.
- Villodolid, D.V. and Bunag, D.M. (1953).** New uses of water hyacinth. In: *Handbook-Utilization and Control of Water Hyacinth in Sudan*.
- Von Loesecke, H.W. (1952).** Citrus fruits industry. *Industrial and Engineering Chemistry*, 44(3): 467-482.
- Wanapat, M. (1983).** Ensilage of rice straw and water hyacinth feed to native cattle and swamp buffaloes. Paper presented at the Second African Workshop "Utilization of low quality roughages with special reference to developing countries". University of Alexandria, Egypt. March, 14-17.
- Weichselbaum, T.E. (1946).** Test combination total protein calorimetric method. *Amer. J. Clin. Path.*, 16: 40.
-

Westlake, D.F. (1963). Comparison of plant productivity. *Biol.Rev.*, 39:385.

Wolverton, B.C.(1975a). Aquatic plants for removal of mevinphos from the aquatic environment. NASA-TM-X-72720 (C.F. Becker et al., 1987).

Wolverton, B.C. (1975b). Water hyacinth for removal of cadmium and nickel from polluted waters. NASA-TM-X-72721 (C.F. Becker et al., 1987).

Wolverton, B.C. (1975c). Water hyacinth for removal of phenols from polluted waters. NASA-TM-X-72722 (C.F. Becker et al., 1987).

Wolverton, B.C. and McDonald, R.C. (1975a). Water hyacinth and alligator weeds for removal of lead and mercury from polluted waters. NASA-TM-X-72723.(C.F. Becker et al., 1987)

Wolverton, B.C. and McDonald, R.C. (1975b).Water hyacinth and alligator weeds for final filtration of sewage. NASA-TM-X-72724.(C.F. Becker et al., 1987)

Wolverton, B.C., and McDonald, R.C. (1975c). Water hyacinth and alligator weeds for removal of silver, cobalt and strontium from polluted waters. NASA-TM-X-72727. (C.F. Becker et al., 1987).

Wolverton, B.C., and McDonald, R.C. (1975d). Water hyacinth for upgrading sewage lagoons to meet advanced waste water treatment standards-Part 1. NASA-TM-X-72729.(C.F. Becker et al ., 1987)

Wolverton, B.C., R.C. McDonald. (1976). Don't waste water weeds.

New Scientist, 71: 318-320.

Yount, J.L. (1964). Aquatic nutrient reduction-potential and possible methods. Rep. 35th Annu. Meeting, Fla. Anti-Mosquito Assoc., PP. 83-85. "C.F. Batanouny and El-Fiky, 1975".

Zeweil, H.S. (1992). Use of a residue from pea (*Pisum sativum*) processing in feeding growing rabbits. Egypt. Poult. Sci., 12: 17-30.

Zeweil, H.S.; Isshak, N.S. and Badawy, N. (1990). Effect of feeding pea by-product to replace yellow corn in the finisher ration on the performance of broiler ducklings. Egypt. Poultry Sci., 10: 69-82.

Zeweil, H.S.; Nour, A.H.; Abou Akkada, A.R. and Isshak, N.S. (1992). The use of water hyacinth (*Eichhorina crassipes* Mart Sloms) in ducklings grower diets. Egypt. Poult. Sci., 12: 287-308.

Zollner, N. and Kirsch. (1962). Test combination of the total lipids. Z-Ges-Exp. Med., 135:545