

REFERENCES

- Abdel-Azim, O.Z. and H.M. Shata (1969).
Pathogenic interaction and physiological studies on four fungi attacking Pelargonium graveolens L.
Proceedings 3 rd. Egypt phytopathology Eng. Cairo: 653-664 .
- Abdel-El-Gallel, A.A. (1977).
Studies on root-rot disease of peanut in A.R.E. and its control.
M.Sc. Thesis, Fac. of Agric. Zagazig Univ., ARE, 1977 .
- Abraham, K.J.; M.R. Menon and M.C. Nair (1976).
Effect of fungicidal treatments on the germination and microflora of sunflower seeds. Agricultural Research Journal of Kerala (1976) 14 (1) 89-90 Coll. Agric., Vellayani, Trivandrum, Kerala, India.
(c.f. Rev. Plant Pathology, 56: 815).
- Acimovic, M. (1962).
Macrophomina phaseoli as a causal agent of sunflower wilt in Vojodina. Zast. Bilja (Plant Prot., Beograd), 1962 (69-70): 125-138, 1962.
(c.f. Rev. Appl. Mycol., 43: 811, 1964).
- _____ (1964).
The occurrence of M. phaseoli on some Agricultural crops and morphological and ecological properties of the parasite. Savr. Poljopr., 12 (1): 55 - 66, 1964 .
(c.f. Rev. Appl. Mycol., 45: 1323, 1966).
- Alabouvette, C. and P. Bremeersch (1975).
Two new diseases of sunflower crops in France. Comptes Rendus des Seances de l' Academie d'Agriculture de France (1975) 61 (11) 626 - 636 Cent. Rech. Agron., Dijon, France.
(c.f. Rev. Plant Pathology, 55: 3243, 1976).

- Allam, A.I.; J.B. Sinclair and P.E. Schilling (1969).
Laboratory and greenhouse evaluations of four
systemic fungicides.
Phytopathology, 59: 1659 - 1662 .
- Anon., (1976).
Agricultural and Economic Statistics Ministry
of Agriculture, ARE .
- A.O.A.C. (1950).
Association of Official Agricultural Chemists,
Second edition.
- Ashour, W.A. and I.F. Gamal El-Din (1962).
Effect of certain environmental factors
on Pre-emergence and Post-emergence damping
off of some ornamentals.
Ann of Agric. Sc., Fac. of Agric., Ain Shams
Univ., Cairo V. 7 No. 1 June 1962.
-
- (1964).
Corm disease of gladiolus in UAR (Egypt).
Ann. of Agric. Sc., Fac. of Agric. Ain Shams
Univ., Cairo, V. 9 No. 2 Dec. 1964 .
- Ashour, W.A.; I.S. Elewa; K.G.M. Ahmed and A.I. El-Fiki
(1978).
Enzyme activity of Macrophomina phaseoli
(Maubl.) Ashby, Fusarium oxysporum Schlecht
and Rhizoctonia solani Kuhn in relation to
damping-off of water melon.
Ann. of Agric. Sc., Fac. of Agric., Ain Shams
Univ., Cairo, V. 9 No. 1 June 1977 .
-
- (1978).
Effect of fungicides on the fungal causing
damping-off of water melon plant in A.R.E.
I: Fungal growth and enzyme activity in Vitro.
Ann. of Agric. Sc., Fac. of Agric., Ain Shams
Univ., Cairo V. 9 No. 1 June, 1977 .
- Békési, P.; J.Voros and O.H. Calvert (1970).
Macrophomina phaseoli in Hungary damaging sunflower
Pl. Dis. Reptr., 54: 286 - 287 .

- Bouhot, D. (1967).
Observations on some diseases of cultivated plant in Senegal. Agron. Trop., 22 (9): 888 - 890. (I.N.R.A.)
(c.f. Rev. Appl. Mycol., 47: 83, 1968).
- Brandes, E.W. (1919).
Bamana wilt.
Phytopathology, 9: 339 - 389 .
- Chan, I.D. (1966).
Saprophytic behaviour; inoculum potential and interaction of cotton root infecting fungi. Ph D. Thesis, Fac. of Agric., Cairo Univ., 174 pp., 1966 .
- Chan, Y.H. and W.E. Saakston (1967).
Translocatable toxic substance produced on sunflower by S. bataticola Taub. Thirty-third Meeting of Candian Phytopath. Ological Society, University of Ottawa, Ontario, May 31, June 2, 1967- Abs. in Proc. Can. Phytopath. Soc., 34: 15 - 26, 1967 .
-
- (1973).
Non specificity of the necrosis-inducing toxin of S. bataticola. Candian Journal of Botany (1973) 51 (3) 690 - 692.
- Crossan, D.F. (1965).
Field and greenhouse experiments for control of Rhizoctonia solani root rot of snapbean. Phytopathology, 55: 503 .
- Dastur, R.H.; R.D. Asana; K. Sawhney; S.M. Sikka and R.S. Vesudeva (1960).
Cotton in Indina, A monograph, Vol. 2. Indian Central Cotton committee, Botany, p. 169 .
- Eisa (Nawal) A., and F.M. Barakat (1976).
Effect of some contact and systemic fungicides on the control of damping-off and Powdery mildew of Cucurbits.
Egypt phytopathology Soc., 1-4 No. 1976-Proc. No. 91.

El-Zarka, A.M. (1976).

The status of sunflower as a promising soil crop in Egypt with ramphasis on disease spectrum.

VII: International sunflower conference
Krasnodar, USSR June 27 - July 3, 1976 .

Enrico, P. and J. Krag (1961).

Influence of date, depth of sowing, and seed dressing on emergence and post-emergence damping-off of groundnuts.

Phytopathologia Mediterranea No. 3 Nov. 1961.

Fahmy, T. (1927).

The Fusarium Disease (Wilt) of cotton and its control.

Phytopathology, 17: 749 - 767 .

————— (1928).

Fusarium Wilt disease of cotton and its control.
Egypt Min. Agric. Tech. and Sci. Bull. 74,
106 pp.

Fakir, G.A.; M.H. Rao and M.J. Thirumalachar (1976).

Seed transmission of M. Phaseolina in sunflower.

Pl. Dis. Repr., 60: 736 - 737 .

Fikry, A. (1932).

Investigation on the wilt disease of Egyptian cotton caused by various species of Fusarium.
Egypt. Min. Agric. Bull. No. 119 .

Ghaffar, A. and D.C. Erwin (1969).

Effect of soil water strass on root-rot of cotton caused by Macrophomina phaseoli.
Phytopathology, 59: 795 - 797 .

Gulidova, L.A. (1970).

Effect of mineral fertilizers on the resistance of winter wheat and barley to root-rot Khisniya sel'Khoz., 8 (4): 46 - 48 .

(c.f. Rev. Plant Pathology, 49: 562 ,1970).

- Ilyas, M.B.; M.A. Ellis and J.B. Sinclair (1976).
Effect of soil fungicides on M. phaseolina,
sclerotia viability in soil and soybean
stem pieces.
Phytopathology, 66: 355 - 359 .
- Joshi, L.K. and S.C. Vyas (1975).
Screening of systemic and non systemic fung-
icides against F. oxysporum causing foot
and root-rot wheat. Indian Phytopathology
(1974), Publ. (1975). 27 (4) 650 - 651.
(c.f. Rev. Plant Pathology, 55: 804).
- Khristova, E. and A. Iordanka (1962).
Some new diseases of plants Rast. Zasht.,
10 (5): 27 - 29, 1962 .
(c.f. Rev. Appl. Mycol., 43: 419 , 1964).
- Kolte, S.J. (1974).
Influence of fungicidal seed treatment on
emergence of sunflowers under field conditions
Pesticides 1974, 8 (9), 40 - 41 (Eng.).
(c.f. Chemical abstracts 82 (15) 23, 1975).
- Korshunova, A.F.; A.E. Tchumacov and R.I. Shekotchikhina
(1966).
Root-rot diseases of wheat and their control.
Koloc, Leningrad. (In Russian) .
(c.f. Ghobrial, E. (1976). Sources of resistance
to root-rot diseases of barley. Agric. Res. Rev.,
2: 9 - 16, 1976).
- Kraeva, G.A. (1960).
Root(rot diseases on wheat in virgin lands and
their control. Ph.D. Thesis, Leningrad (In
Russian).
(c.f. Ghobrial, E. (1976). Sources of resistance
to root-rot diseases of barley. Agric. Res.
Rev. 2: 9-16, 1976 .
- Littrell, R.H. (1966).
Effect of nitrogen nutrition on susceptibility
of chrysanthemum to an apparently new Bitype
of F. oxysporum.
Pl. Dis. Reptr., 50: 882 - 885. 1966 .

- Luciano, A. and M. Davereux (1967).
Sunflower production in Argentina. publ. tec.
Estac. Exp. Pergamino 37, Vii + 53 pp.
(c.f. Rev. Appl. Mycol., 47: 1222, 1968).
- Maklad, F.M. (1978).
Studies on some fungal sunflower diseases and
their control.
M.Sc. Thesis, Fac. of Agric., El Azhar Univ.,
Cairo 109 pp., 1978 .
- Marengo, L.V. (1942).
Fusariosis of the sunflower (Helianthus annuus L.).
Rev. Fac. Agron., B. Aires., X, 1, pp. 130-147,
1942.
(c.f. Reg. Appl. Mycol., 22: 361, 1943).
- Marras, F. (1963).
Ground nut; sunflower and cowpea, new hosts
for M. phaseoli in Sardinia. Note fitopat.
Sard. 3, 10 pp., 2 pl., 1963 .
(c.f. Rev. Appl. Mycol., 43: 421, 1964).
- Mathur, S.B. (1968).
Production of toxins and pectolytic enzymes
by two isolates of S. bataticola Taub. and
their role. in pathogenesis. Phytopathology
Z., 62 (4): 327 - 333 .
(c.f. Rev. Appl. Mycol., 48: 376, 1969).
- Middleton, K.J. (1971).
Sunflower diseases in South Queensland.
Qd. Agric. J. 97 (11): 597 - 600 .
(c.f. Rev. Plant Pathology, 51: 2749, 1972).
- Mostafa, M.A.; M.S. Naim and S.K. Moawad (1957).
Studies on the interaction between F. oxysporum
schlecht and M. phaseoli (Maubl.) Ashby in
parasitizing "Karnak" and "Giza 30" cotton
varieties and in cultures 11: Pathogenicity
of Fusarium and Macrophomina, singly and in
association, under different nitrogenous man-
urial treatments.
The Third Arb. Sci. Cong., 10-20 Sept., 1957,
Beirut: 57 - 67 .

- Orellana, R.G. (1971).
Fusarium Wilt of sunflower (Helianthus annuus):
first report.
Pl. Dis. Reprtr. 55 (12): 1124 - 1125 .
- Pastorino, A. (1965).
A polyphagous fungus S. rolfsii. Bol. Fac.
Agron. Univ., Montevideo 750: 1 - 6, illus,
1964. (Biol. Abstr., 46 (19), 86591, 1965).
(c.f. Rev. Appl. Mycol., 45: 775, 1966).
- Sackston, W.E. (1957).
Diseases of sunflowers in Uruguay.
Pl. Dis. Reprtr., 41: 885 - 889, 1957 .
- _____ (1958).
Black root-rot of sunflowers in Uruguay caused
by S. bataticola Taub. Abs. in Proc. Canad.
Phytopath. Soc., 26 (1958), p. 14, 1959.
(c.f. Rev. Appl. Mycol., 38: 370, 1959).
- _____ (1978).
Sunflower disease mapping in Europe and Adjacent
Mediterranean countries.
Helia. Information Bulletin of the F.A.O.
Research Network on sunflower Nr. 1-December
1978. p. 21 - 28 .
- Sahran, G.S. and V.K. Gupta (1974).
Control of collar rot of soybean with fungicides.
Indian Journal of Mycology and Plant Pathology
(1973. Publ. 1974) 3 (2) 222 - 224 .
(c.f. Rev. Plant Pathology , 54: 573, 1975).
- Sacker, P. (1946).
The effect of planting on the emergence and
surivval of blue lupine.
Phytopathology, 36: 479 - 480.
- Shabana , R. and S. Abou-Khadra (1976).
The possibility of sunflower seed production,
the first crop for producing oil yield, in
Egypt for sailing.
The Egyptian second congress for year 2000,
Scientific meeting of outside Egyptian Collage.
December, 1976. (In arabic).

- Sharvelle, E.G. (1961).
The nature and uses of modern fungicides
Burgess Publishing Company 1961, printed in the
united states of America pp.308.
- Simmonds, J.H. (1956).
Science Branch. Plant Pathology Section.
Rep. Dept. Agric. Od., 1955 - 1956, pp. 65-66,
1956 .
(c.f. Rev. Appl. Mycol. 36: 307 -, 1957).
- Sivaprakasam, K.; K., Pillayarsmy and S.Ayyamperumal.
(1975).
Studies on sunflower root-rot disease. II.
Effect of seed treatment. Madras Agric.
J. (1975). 62 (2) 74-75 Reg. Agric. Res.Sta.,
Kovilpatti, Tamil Nadu, India.
(c.f. Rev. Plant Pathology, 55: 5306, 1976).
- Sivaprakasam, K.;K. Pillayarswamy and S. Rajaram (1975).
Effect of NPK fertilizers on root-rot disease
incidence in sunflower. Madras Agric. J. (1975).
62 (5) 308 - 309 Cotton and Millet Exp. Sta.,
Kovilpatti, Tamil Nadu, India.
(c.f. Rev. Plant Pathology, 56: 816, 1977).
- Smith, A.L. (1940).
A regional study of the relationship of potash
treatments to the development of cotton with
under widely varying conditions of soil and
environment.
Phytopathology, 30: 707 .
- Thrap, W.H. and C.H., Wadleigh (1939 a).
The effects of nitrogen sources, nitrogen level,
and relative acidity on Fusarium wilt of cotton.
Phytopathology, 29: 756 .
(1939b).
Effect of nitrogen, phosphorus, and ptassium
nutrition on the Fusarium wilt of cotton.
Phytopathology, 29: 756 - 757 .

- Tikhonov, O.I.; V.K. Nedelko and T.A. Perestova (1976).
Infection period and spreading pattern of
S. bataticola in sunflower.
- VII. International sunflower conference abstracts
of papers Krasnodar June 27-July 3, 1976 .
- Vasudeva, R.S. (1930).
Studies on the physiology of parasitism X (11)
on the effect of one organism in reducing the
parasitism activity of another.
Ann. Bot., 44: 557 - 564 .
- Wolf., T.F. and F.A. Wolf (1948).
A toxic metabolic product of F. oxysporum var.
nicotiana in relation to a wilting of tobacco
plants.
Phytopathology, 38: 292 - 298 .
- Wallen, V.R. and W.Bell (1956).
Treatment of vegetable seed for improved
emergence.
Pl. Dis. Repr., 40: 129 - 132 .
- Wellman, F.L. (1943).
Comparative toxix effect of extracts from mild
and virulent isolates of tomato-wilt.
Phytopathology, 33: 1004 - 1017 .
- Woltz, S.S. and A.W.Engelhard(1973).
Fusarium wilt of chrysanthemum : I Effect of
nitrogen source and live on disease develop-
ment.
Phytopathology, 63: 155 - 157 .