

REFERENCES

LITERATURE CITED

- Abd El-Fattah, A.; M.H. Hilal and M. Saber (1984). Effect of sulphur application on pH, EC and availability of phosphorus in different Egyptian soils. Proc. Second National Conf. on the Problems of Land Degradation, Cairo, Egypt (Nov. 19-22).
- Ahuja, L.D.; C.M. Verma; S.K. Sharma and T.R. Lamba (1978). Range management studies on the contribution of ground storey (grass) in afforested areas in arid regions. Annals of Arid Zone, 17 (3): 304-310.
- Alawi, B.J.; J.L. Strohlein; E.A. Hanlon and F. Turner (1980). Quality of irrigation water and effects of sulphuric acid and gypsum on soil properties and Sudan grass yields. Soil Sci. 129:315-340.
- Association of Official Agriculture Chemists (1970). Official methods of analysis. A.O.A.C. Washington, 11th ed., D.C. pp. 832.
- Aulakh, M.S. and N.S. Pasricha (1977). Interaction effect of sulphur and phosphorus on growth and nutrient content of mung bean (*Phaseolus aureus* L.). Plant and Soil, 47: 341-350.
- Bakhshish, S.; H.N. Mathur and P. Joshie (1980). Effect of tree shade on grassland production in the moist subtropical region of northern India. Indian J. Forestry 3 (4): 345-348.
- Barrow, N.J. (1977). Phosphorus uptake and utilization by tree seedlings. Aust. J. Bot., 25: 571-584.

- Church, D.C. (1980). Digestive physiology and nutrition of ruminants. Vol. 3, Proccical Nutrition; 2nd Edt.
- Draz, M.Y.; M.M. Wassif and S.E. El-Maghraby (1993). Improvement of Siwa dune sand for the growth of acacia plants under saline water regime. Fourth Congress, 24-25 Nov. Egypt. Soil Sci. Soci., Cairo, Egypt.
- Dubbs, A.L. (1971). Competition between grass and legume species in dryland. Agron. J. 63: 359-362.
- Dubois, M.K.; J.K. Gilles; P.A. Hamilton and F. Smith (1950). A colormetric method for the determination of sugar. Nature; 168-169.
- El-Bagouri, I.H.; M.A. El-Kadi and S.A. Sabet (1978). Nitrogen, phosphorus and copper requirements of barley and the conditions of highly calcareous soils. Desert Inst. Bull., A.R.E., 28 (2): 437-447.
- El-Gala, A.M.; M.A. Mostafa and S.E. El-Maghraby (1989). Influence of sulphur and saline irrigation water on growth and elemental status of barley plant grown on calcareous soils. Egypt. J. Soil Sci. Special Issue 31: 443-455.
- El-Hakeem, M.S. (1990). Effect of forage mixtures at different mixing ratios on yield and quality. Egypt. J. Appl. Sci., 5 (7): 804-817.
- El-Maghraby, S.E.A. (1988). Effect of sulphur as a soil amendment on ionic balance in calcareous soil under saline water regime. M. Sc. Thesis Soil Dept., Ain Shams Univ.

- El-Naphate, M.; G. El-Zamzame and A. El-Hamrone (1986). Range plants in arid zones of North Africa. Green Belt Project for North Africa; pp. 260.
- El-Rashidi, M.M.; F. Labib and M.Y. Tayel (1975). The effect of super phosphate fertilization on barley plants grown on highly calcareous soil. Egypt. J. Soil Sci., Special Issue, pp. 327-337.
- Farid, M.F.A.; M.F. Wardeh; L. Cerol; L. Haris and H. Loid (1979). Arab and Middle East Tables of Feed Composition. Arab Center for the Studies of Arid Zones and Dry Lands (ACSAD) and Utah Agric. Exp. Stat. pp. 554.
- Frie, E.; K. Payer and E. Schutz (1964). Determination of phosphorus by ascorbic acid. Schw. Landwert Sch. Forschung. Heft, 3: 318-328.
- Gabra, M.A.; M.A. Nor El-Din and E.Z. Youssef (1984). The effect of N.P.K. on the yield of berseem/barley mixture. Proceeding ENCIP Symposium; 6-7 Nov. Agric. Res. Center, Egypt.
- Garcia, L.; M.E. Desa and L. Carloni (1977). The effect of sulphur on the solubility and forms of phosphorus in soil. Agrochemica, 21: 163-169.
- Gill, A.S.; B.O. Patil and C.L. Yadav (1988). Mixed cropping of barley and senji for green fodder in partially irrigated lands in dryland. Agric. Sci. Digest India, 8 (1): 12-14.
- Goss, D.W. and B.A. Stewart (1979). Efficiency of phosphorus utilization by Alfalfa from manure and superphosphate. Soil Sci. Amer. J., 43: 523-528.

- Gupta, V.K. and D.S. Mehla (1980). Influence of sulphur on the yield and concentration of copper, manganese, iron and molybdenum in berseem (*Trifolium alexandrinum*) grown on two different soils. *Plant and Soil*, 56: 229-234.
- Gutteridge, R.C. (1991). Agronomic evaluation of tree and shrub species in South-East Queensland. *Fer. Abst.*, 52 (3): 1366.
- Hassan, M.Th. (1994). Legume and grain production within rainfed agroforestry system in the Mediterranean Coast of Egypt. First International Symposium on Silviculture of Protection Forestry in Arid Regions and the Agroforestry Potential, ARC; NARP and USAID, Alexandria.
- Hassan, N. and R.A. Olson (1966). Influence of applied sulphur on availability of soil nutrient for corn (*Zea mays* L.) nutrition. *Soil Sci. Soc. Am. Proc.* 30: 286-290.
- Helal, M.H. and A. Abd-El-Fattah (1987). Effect of CaCO_3 and clay content of alkaline soils on their response to added sulphur. *Sulphur in Agriculture*, 11: 15-19.
- Hingston, F.J.; N. Malajczuk and T.S. Grove (1982). Acetylene reduction (N_2 -fixation) by jarrah forest legumes following fire and phosphate application. *J. Appl. Ecol.* 19 (2): 631-645.
- Ibrahim, M.K. (1981). Shrubs for fodder production. *Advances in food producing systems for arid and semi-arid lands*, National Agricultural Research Station, Kenya 601-642.

- Ivanic, J. (1977). Effect of various application rates of sulphur on uptake of nutrients and yield of spring barley. *Soils and Fertilizers*, 43 (4): 3366.
- Jasper, D.A.; L.K. Abbott and A.D. Robson (1989). Acacias respond to additions of phosphorus and to inoculation with VA mycorrhizal fungi in soils stockpiled during mineral sand mining. *Plant and Soil*, 115 (1): 99-108.
- Jones, U.S. (1982). *Fertilizers and soil fertility*. Second Edition, Aeslon Publishing Company, Reston, Virginia
Aprentice-Hall Company.
- Kandeel, S.A.; A.E. Kherallah and B.E. Bourthami (1994). Biomass utilization of juvenile *Acacia cyanophylla* Lindl. grown in Egypt. First International Symposium on Silviculture of Protection Forestry in Arid Regions and the Agroforestry Potential, ARC, NARP and USAID, Alexandria.
- Kashired, A. (1972). Effect of sulfur and residual sulphur on pH and availability of phosphorus in calcareous soil and effect of applied nitrogen on yield and chemical composition of corn. *Zeitschrift für Pflanzenernährung und Bodenkunde*, 133: 93-98 (c.f. *Soil and Fert.* 37; 405, 1974).
- and J. Bazargani (1972). Effect of sulfur on pH and availability of phosphorus in calcareous soils. Influence of sulphur and nitrogen on yield and chemical composition. *Zeitschrift für Pflanzenernährung und Bodenkunde*, 131: 6-13 (c.f. *Soil and Fert.* 35, 4391).

- Khater, A.M. (1981). A study of sulphur and petroleum products as efficient materials affecting the availability of certain nutrients in soils. M. Sc. Thesis, Fac. of Agric., Ain Shans Univ.
- Le Clerg, E.L.; W.H. Leonard and A.G. Clark (1966). Field plot technique. Burgtoss Publishing Co. Minneapolis Minnesota, U.S.A.
- Lunnan, T. (1987). Mixtures of barley and legumes for green feed. Nordisk-Jordbrugsfors Kning Bibliographic Citation, Nordisk-Jordbrugsforskning, 69 (2): 293; Norwegian Language Translated in English.
- Maasdorp, B.V. and R.C. Gutteridge (1986). Effect of fertilizer and weed control on the emergence and early growth of five leguminous fodder shrubs. Tropical Grasslands. 20 (3): 127-134.
- Mahmood, H.A.M.; P.K. Ponnuswamy and M. Viswanathan (1980). Nutrition studies on black wattle (*Acacia mearnsii*). 1. Effect of different N.P.K. fertilizer combinations up to the 6th year. Indian Forestry, 106: 397-402.
- Maltby, J.E. and A.A. Webb (1983). Studies on solidic soils under *Acia harpophylla*. Queensland J. Agric. of Anim. Sci. 40 (1): 53-63.
- Malavolta, E.; G.C. Vitti; C.A. Rosolem; H.J. Klemann; N.K. Fageria; P.T.G. Guimaraes and M.L. Malavolta (1987). Sulphur in Brazilian Agriculture, 11: 2-5.
- Mehrotra, O.N. and S.H.A. Ali (1970). Studies on mixed cropping of barley and grass. Indian J. Agron. 15 (1): 277-280.

- Mehta, U.R. and H.B. Singh (1979). Response of greengram to sulphur on calcareous soils. Indian J. of Agric. Sci. 49 (9): 703-706.
- Messina, M.G. and I.L. Barton (1985). Early growth and survival of *Acacia melanexylon* effect of weed control and fertilizer. Newzealand J. of Forestry Sci., 15 (1): 111-116.
- Milton, S.J. (1983). *Acacia tortilis* sub sp. heteracantha productivity in the tugela dry valley Bushveld: preliminary results. Proceeding xth AETFAT congress. SCIR Conference Centre, 19-23 January. Pretoria, Republic of South Africa.
- Moosa, G.D. (1986). Yield dynamics and chemical composition of canopy components in alfalfa-grass associations. Dissertation Abst.; Sci. and Engin. 46; (12): 4074 B.
- Orphanos, P.I. (1987). Direct and residual effect of fertilizer phosphorus on barley in cyprus. International Center for Agricultural Research in Dry Areas, (ICARDA), Syria. 120: 44-49.
- Overstreet, R.; J.C. Martin and M.L. King (1951). Gypsum, sulfur and sulfuric acid for reclaiming on alkaline soil of the Fresno series. Hilgardia. 21: 113-127.
- Paper, I.L. and R.H. Miller (1978). Comparison of the oxidation of thiosulfate and elemental sulfur by two heterorophic bacteria and thiobacillus. Soil Sci. 126 (1): 9-14.
- Peach, K. and M.V. Tracey (1956). Modern methods of plant analysis. 1. Springer-Verlag Berlin.

- Prasad, D.; K.M. Singh; R.N. Katiyar and R.N. Singh (1987). Impact of intercropping on the plant growth, pest incidence and crop yield of pea, *Pisum sativum* Linn. Indian. J. of Enomology 49 (2): 153-172.
- Pressland, A.J. (1976). Possible effects of removal of mulga (*Acacia onura*) on rangeland stability in south western Queensland. Australia Rangeland J. 1 (1): 24-30.
- Radwan, M.S. and A.K. El-Fakhry (1975). The value of vetches for forage production in Northern Iraq. Mesopolamia J. Agric. 10 (1-2): 35-40.
- Radwan, M.S. and A.K. El-Fakhry (1976). Forage crops and pastures. Agronomy Dept., Fac. of Agric. and Forestry, El-Mosel Univ. pp. 651.
- Ramos, J.M.; L.F. Garcia-Del-Moral; J.L. Molina-Cano; P. Salamanca; F. Roca-De-Togores; L.F.G. Moral and F.R. Togares (1989). Effects of an early application of sulphur or ethephon as foliar spray on the growth and yield of spring barley in a Mediterranean environment. J. Agro. of Crop Sci. 163 (2): 129-137.
- Rawal, O.R. and G.L. Yadava (1988). Response of barley to N, P, K and Zn under dryland conditions in Chittogarrh district. Annals Agric. Res. 9 (2): 263-265.
- Rehm, G.W. (1987). Application of phosphorus and sulfur on irrigated alfalfa. Agron. J. 79 (6): 973-979.
- Reuter, D.J.; A.M. Alston and T.G. Heard (1973). Correction of manganese deficiency in barley crops on calcareous soils. III. Application of elemental sulphur. Aust. J. Exper. Agric. and Anim. Husb., 13 (63): 446-451.

Ryan, J. and J.L. Stroehlein (1979). Sulfuric acid treatment of calcareous soils. Effects on phosphorus solubility, inorganic phosphorus forms and plant growth. Soil Sci. Soc. Am. J. 43: 731-735.

Sabet, S.A.; M.A. Abdel-Salam; A. El-Leboudi and A.A. Robishy (1970). Biological evaluation for phosphorus availability in certain calcareous soils of the U.A.R. Desert Inst. Bull. 20: 135-150.

Salem, M.O.; M.S. El-Hakeem and H.M. Abd El-Aziz (1985). Seed collection project for the north western zone of Egypt. Final Report. Desert Research Center.

----- ; A.A. El-Sall; M.M. Wassif and M. Hilal (1989). Effect of sulfur, nitrogen and organic manure application on the growth characters of wheat plant under calcareous soil and saline irrigation water conditions. Desert Inst. Bull., A.R.E., 39 (1): 123-156.

----- and M.S. Barsoum (1993 a). Response of some legume forage plants to bacterial inoculation and phosphorus fertilization under Al-Arish region conditions. Egypt. J. Appl. Sci., 8 (2): 593-602.

----- and M.S. Barsoum (1993 b). Effect of bacterial inoculation and phosphorus fertilization on some forage legumes under dry farming conditions at Rafah region, North Sinai. Egypt. J. Appl. Sci., 8 (6): 61-75.

Saper, R.J. and I.H. El-Bagouri (1964). The effect of soil carbonate level on availability of added and native phosphorus in some calcareous Manitoba soils. Can. J. Soil Sci. 44: 327-344.

- Schonau, A.P.G. (1984). Fertilization of fast-growing broadleaved species. South African Forest Res. Inst. 17: 253-268.
- Seif, S.A.; M.S. El-Hakeem and S.H. Hendawy (1990). Growth behaviour of two acacia species as affected by nitrogen and phosphorus application. Egypt. J. Appl. Sci., 5 (8): 96-106.
- Sheha, M.Y. (1984). Adaptation of plants in Kuwait. Kuwait Foundation for Sci. Development, First Edition.
- Shekhawat, J.S.; N.L. Sen and L.L. Somani (1988). Evaluation of agro-forestry systems under semi-arid conditions of Rajasthan. Indian Forester, 114 (2): 98-101.
- Skerman, P.L. (1977). Tropical range legumes. FAO, Plant Production and Protection Publication No. 2, p. 524.
- Steel, R.G.D. and J.H. Torrie (1980). Principles and procedures of statistics. 2nd, McGraw-Hill Co. N.Y., U.S.A.
- Tiedman, J.A. and D.E. Johnson (1992). *Acacia cyanophylla* for forage and fuelwood in North Africa. Agroforestry Systems, 17: 169-180.
- Thalen, D.C.P. (1979). Ecology and utilization of desert shrub range lands in Iraq. Dr. W. Junk bv. Publishers.
- Thomas, M.B. (1981). Nutrition of five species of container-grown acacia, boronia, choisya and eucalyptus. Scientia Horticulturae, 14: 55-68.

- Trevino, J.R.; R. Caballera and J. Gill (1980). Studies on the utilization of algarroba (*Vicia monantha* Rehz) as a forage plant, growth and yield analysis. Pastos 10 (2): 138-143.
- Trevino, T.; R. Caballero and J. Gil (1981). The utilization of tare (*Vicia monantha* Pehiz) as a forage plant, chemical composition, digestibility and energy value at different stages of maturity. Pastos, 11 (2): 361-370.
- Venkataramanan, C.; B. Haldorai; P. Samraj; S.K. Nalatwadmath and C. Henry (1983). Return of nutrient by the leaf-litter of bluegum (*Eucalyptus globulus*) and black wattle (*Acacia mearnsii*) plantations of Nilgiris in Tami Nadu. Indian Forester 109 (6): 370-378.
- Wassif, M.M. (1994). The relationship between soil amendments application and utilization of saline irrigation water in arid regions. Symposium on Desert Studies on the Kingdom of Saudi Arabia, 2-4 October, Center for Desert Studies; King Saud University.
- ; A.A. El-Shall and I.H. El-Bagouri (1986). Alfalfa response to phosphate fertilizers and its residual effect under calcareous soil conditions. Desert Inst. Bull., A.R.E., 36 (1): 1-18.
- ; H. Bassiouny and M.A. Abdel Salam (1980). Phosphorus utilization under saline water irrigation using P33 labelled fertilizer. Isotope Rad. Res. 12 (2): 127-136.
- Willey, R.W. (1979). Intercropping, its importance and research needs. Part I. Competition and yield advantages. ICRISAT; Hyderabad 500016, A.P., India (c.f. Feld Crop Abst. 32 (1): 1-10).

Yadav, J.S.P. (1980). Salt affected soils and their afforestation. Indian Forester, 106 (4): 259-272.

Younis, A.A.; M.A. Harfoush and K.N. Ghobrial (1986). Effect of interseeding some winter leguminous forage crops with barley on forage yield and quality. Proceediing Second Conf. Agron., Alex. Egypt. 1: 763-771.

Zaght, M.F.; S.S. Tag Eldin and Elsh. Osama (1993). Guide for the improtant plants of the desert garden. Center of Research Studies Station, King Saud University, Kingdom of Saudi Arabia, Riyadh.