

*R*EFERENCES

REFERENCES

- Abd El-Kader, M.A. (1977).
Studies on some diseases of lentil.
M. Sc. Thesis. Fac. Agric., Assiut Univ.
- Abd El-Moity, T. H. and M.N. Shatla (1978).
Biological control of white rot of onion
(Sclerotium cepivorum Berk.) by Trichoderma
harzianum the third International Plant Path-
ology conference p. 197.
- Abd El-Moity, T. H. and M.N. Shatla (1981).
Biological control of white rot disease of
onion (Sclerotium cepicorum) by T. harzianum
Phytopathology Z. 100 : 29 - 35.
- Abd El-Rehim, M.A. (1962).
Studies on the organisms causing root-rot and
wilt of horse beans, Vicia fabae var. ecuine
in U.A.R Ph. D Thesis, Fac. Agric., Alex. Univ.
- Abdou, Y.A., M. M. Ragab and Olfat M. Mousa (1970).
Microbial association in Rhizoctonia root-rot
of beans. U.A.R J. Phytopathol. 2, 59 - 62.
- Agrawal, S.C.; N.M. Kare and S.P. Agrawal. (1977).
Biological control of Sclerotium rolfsii
causing collar rot of lentil.
Indian Pathopathol. 30 (2) : 176 - 179.
- Ali, M. D. H. (1967).
Studies on root-rot disease of broad bean and
its control in U.A.R Ph. D. Thesis, Fac. Agric.,
Ain Shams Univ.

- Aly, A. Z. (1974).
Studies on soil fungi causing root-rot to lupine.
M. Sc. Thesis, Fac. Agric., Zagazig Univ.
- Alimeida, R. T. and C.M.U. Landim. (1981).
Preliminary studies on the biological control
of Sclerotium rolfsii sacc., causal agent of
sclerotial wilt of cowpea (*Vigna unguiculata*
(L.) Walp.)
CNPQ, Fortaleza, Ceara, Brazil.
Fitossanidade 1981. 5 (1) : 15 - 20.
(c.f. the C A B abstracts database).
- Armstrong, G. M., and J.K. Armstrong (1963).
Lupine and other legume hosts of the Fusarium
wilt from U.S cotton.
Netheral. J. Plant pathol. (Tijdschr. Plziekt.)
69 (1) : 3-75 . (R.A.M., 42 : 551, 1963).
- Ashour, W.A., A.R. Sirry and H.S. Salem (1964).
Studies on French bean root-rot caused by Scl.
rolfsii sac. in U.A.R (Egypt).
Ann. Agric. Sci., Fac. Agric., Ain Shams Univ.
Cairo, Vol. 9 No. 1 , June (1964)
- Badr, A. B. (1975).
Studies on some important damping off and root
rot fungi of lentil.
M. Sc. Thesis, Fac. Agric. Al-Azhar Univ.
- Baker, K. F. and R. J. Cook (1974).
" Biological Control of Plant Pathogens "
W.H. Freeman and CO., San Francisco, 433. P.

- Baker, R (1968).
Mechanisms of biological control of soil-borne pathogens.
Annu. Rev. Phytopathol. 6, 263.
- Barentt, H.L. and B.B. Hunter (1972).
Illustrated Genera of Imperfect Fungi.
Burgess publishing company. Minneapolis,
Minnesota.-- pp
- Broadbent, P. and K.F. Baker (1969).
Bacteria and Actinomycetes antagonistic to root pathogens in Australian soil.
Phytopathology, 59 : 1019 - 1021.
- Broadbent P. and V. Waterworth (1971).
Bacteria and Actinomycetes antagonistic to fungal root pathogens in Australian soils.
Aust. J. Biol. Science, 24 (5).935-944.
- Broadbent, P., K.F. Baker, N., Franks, and J., Holland (1977).
Effect of Bacillus spp. on increased growth of seedlings steamed and in non-treated soil.
Phytopathology, 67, 1027.
- Carrera, C. M. (1940).
The genus Fusarium in the Argentine Republic.
Study and Systemic classification (second contribution)
Rev., Argentin, Agron. 7 : 277 - 286
(R. A. M., 20 : 235. 1941).

- Carrera, C.M. and M. Noll (1941).

The importance of some species of Fusarium in the root-rot and wilt of lupines (Lupinus albus and Lens esculenta in Uruguay. An. Soc. Cient. Argent. 131 (4) : 152 - 184, (5) : 185 - 211 (c.f.R.A.M;23 : 230, 1944).

- Clements, F. E and G. L. Chear (1957).

The Genera of Fungi.

Hofner Publishing Co. New York. 496 P.

- Chidambaram; P. S. B. Mathur and P. Neergaard (1973).

Identification of seed-borne Drechslera species. Friesia, 10 : 165 - 207.

- Damicone, J. P. and W. J. Manning (1980).

Biological control of Fusarium crown rot of Asparagus. Greenhouse, and field results compared (Abstract in Beltsville symposia in Agricultural Researchs. V. Biological control in crop production, May 18 - 21, 1980. P. 3.

- Domsch, K.H., W. Games and T. Anderson (1980).

Compendium of Soil Fungi

Academic press. A subsidiary of Harcourt Brace

Hovanovich Publishers, London, New York,

Toronto, Sydney and San Francisco, (1) : 859 PP.

- Dumitras, L., T. Sesan (1980)

Features of antagonism of Trichoderma virde
Pers. ex Fr. to Rhizoctonia solani Kuhn.
Res. Inst. Pl. Prot., Bucharest, Romania.
Studu si cercetari de Biologie. Biologie
Vegetala 1980. 32 (1) : 87 - 92.
(c.f R. Pl. Pathol. 1973)

- Elad, Y., I. Chet and J. Katan (1980).

Trichoderma harzianum a biological agent effective against Sclerotium rolfsii and Rhizoctonia solani. Phytopathology, 70 : 119 - 121.

- Elad, Y., I. Chet and Y. Henis (1980).

Biological control of Rhizoctonia solani in
strawberry fields by Trichoderma harzianum.
Plant and soil 60, 245 - 254.

- El-Helaly, A. F., I. A. Ibrahim; M. W. El-Shawah, H.M. El-Arosi, M.K. Abo El-Dahab, S.H. Michail, M. A., Abd El-Rehim, E.H. Wasfy, a,d M.A. El-Goorani (1966).

General survey of plant diseases and pathogenic
organisms in U.A.R.
(Egypt) Until 1965. Alex. J. Agric. Rev. Bull.
No. 15 (1966).

- El-Shanwani, M. (1973).

Studies on root-rot diseases in lentil.
M. Sc. Thesis, Fac. Agric. Al-Azhar Univ.

- Fedorinchik, N.S., T. A. Terunina, M.G. Tyutinnikov and K.I. Kudryavtva (1975).
Trichoderma -4- a new biological preparation for plant disease control.
VIII Intern. Cong. Plant prot., Vol. III :
67 - 72, Moscow 1975.
- Georgey, N.I. (1977).
Studies on Sclerotium cepivorum Berk.
causing white rot of onion and its control.
Ph. D. thesis, Fac. Agric. Ain Shams Univ.
- Gilman, J.C. (1957).
A Manual of Soil Fungi (Second Edition)
State College Press, Ames, Iowa, U.S.A
- Guy, S. O and R. Baker (1977).
Inoculum potential in relation to biological control of Fusarium wilt of peas.
Phytopathology 67 : 72.
- Hadar, Y., I. Chet and Y. Henis (1979).
Biological control of Rhizoctonia solani damping - off with wheat bran culture of Trichoderma harzianum. Phytopathology, 69, 64 - 68.
- Hansen, H.N. (1926).
A simple method of obtaining single spore
Science, 64 : 384 , 1959.

- Henis, Y., J.A. Lewis, and G. C. Papavizas (1984).
Interactions between Sclerotium rolfsii and Trichoderma spp. relationship between antagonism and disease control.
Soil Biol. Biochem. 16 (4). 1984. 391 - 396
(c. f the CAB abstracts database)
- Huang, H. C (1979).
Biological control of Sclerotinia wilt in sunflower.
Canada Agriculture (1979) 24 (3) 12 - 14
Canda Res. Sta., Morden, Man., Canda.
(R.P.P. 59 : 2295, 1979)
- Johenson, L.E., E.A.Curl, J.H.Bond and H.A. Fribourg (1960).
Methods for Studying Soil Microflora. Plant disease relationships.
Second printing, 1960.
Burgess Publishing Company, Minn. U.S.A
- Katznelson, H. and S.E. Cole (1965).
Production of gibberellin-like substances by bacteria and actinomycetes. Gen. J. Microbial 11, 733.
- Lang, K.J. (1975).
Experiments with fungi causing damping-off. 1- Interrelation between Trichoderma viride Pers. ex. fr and races of pathogenic fungi in the genera Pythium, Fusarium and Rhizoctonia. Eurs for Pathol. 5 (4) : 225 - 240.

- Neweigy. N.A., N.A. Eisa, and L.A.El-Shewy (1982).
Biological control of damping-off in broad bean cultivars Giza 2 and rebaya 40.
Ain Shams Univ. Fac. Agric. Res. Bull.
No. 1778; 1 - 28.
- Noll, W. (1939).
Studies on foot and wilt diseases in leguminosae
Zeitschr. Pflanzenkr., 49., 385-430.
(c.f., Biol. Abs., 15 : 3015, 1941).
- Old, K.M. (1965).
Fungistatic effects of soil bacteria on root rotting fungi with particular response to Helminthosporium sativum.
Phytopathology, 55 : 501 - 505.
- Qasem, S.A and C.M., Christensen (1958).
Influence of moisture content, temperature, and time on the deterioration of stored corn by fungi. Phytopathology, 48 : 544 - 549.
- Ram Nath, S.M.Mathur and P. Neergaard (1970).
Seed borne fungi of mung bean (Phaseolus aureus Roxb.) from India and their significance.
Proc. Int. Seed test Assoe., 35 : 225 - 241.
- Sahab, A.F. (1976).
Rhizosphere and physiological studies on the effect of some pesticides on some plant pathogens.
Ph. D. Thesis, Fac. Agric, Ain Shams Univ.

- Saltenis, B. (1967).
Fusariosis of leguminous crops in lithuania.
1 - Specific composition and distribution of
the causal agents of the disease.
T. S. R. Mokslu Akad. Darb. Ser. C. 2 (43):87-98.
(c.f.R.A.M., 47:419- 1968).
- Schultz, H. (1939).
Investigations on the role of Pythium species
as agents of lupine root-rot.
Phytopathology. Z., 12 (4) : 405 - 420.
(R.A.M., 19 : 282, 1940).
- Shekunova, E.G (1970).
Disease of fodder seed lupine in the Chernigov
region.
Tree dyvses. Inst. Zashch. Rast. 29 (1). 44 - 51.
(R.P.P; 50 : 542, 1971).
- Sirry, A. R., W.A. Ashour and M.D.H. Ali (1970).
Studies on root-rot disease of broad bean.
Res. Bull. 393, Fac. Agric., Ain Shams Univ, 21pp .
- Sivans, A. and I Chet. (1984).
Biological control of Fusarium spp.
(c.f. proc. 6 th Cong. . Un. Phytopath.
Mediterr. Cairo Egypt (1984).
- Tschen, J.S.M. and W.L. Kuo (1985).
Antibiotic inhibition and control of Rhizoctonia
solani by Bacillus subtilis.
Dep. of Botany, Nath. chung-Hsing Univ.,
Taichung, Tiwan, Roc.
Plant Prot. Bull. 27 (2) 1985. 95 - 103.

- Vesely, D. (1979).

Use of Pythium olgandrum to protect emerging
sugar beet (In soil-borne plant pathogens)

Schippers, B. and W. Gams (Eds.) (1979).

Academic Press, London New York, San Francisco,
593 - 595.

- Weimer, J.L. (1944).

Some root-rots and a foot-rot of lupines in the
southeastern part of the United States.

J. Agric, Res., 68 (12) : 441-457.

- Weindling. R. (1932).

Trichoderma lignorum as a parasite of other
soil fungi.

Phytopathology, 22 , 837.

- Wen-Shi-Wu (1976).

Biological control of seed and soil borne fungi
associate with wheat and oats.

Bot. Bull. Academia Science, 17 : 161 - 168.

- Wilson, V. E. and J. Bransberg (1965).

Fungi isolated from diseased lentil seedlings.

Pl. Dis. Reprtr., 49 (8) 660 - 662.

- Windaham, M.T., Y.Elad and R. Baker (1986).

A mechanism for increased plant growth induced
by Trichoderma spp.

Phytopathology, 76 : 518 - 521.

- Wood, R.K.S. (1951).

The control of lettuce by the use of antagonistic organisms. II- the control of *R. solani* Kuhn.
Ann. Appl. Biol., 38 : 217 - 230.

- Yamamata, W., N.oyasu and K.Takigawa (1955).

Studies on the wilt disease of broad bean
J. Sci. Rep. Hyogo. Univ. Agric. 2.1,
PP. 52 - 62 (1955)
(R.A.M., 37 : 139).

- Yehia, A. H., S. A. El-Hassan and A.H.El-Bahali (1982).

Biological seed treatment to control Fusarium
root-rot of broad bean. Egypt. J,
Phytopathol., 14, 1-2 59-66.

- Yu, T. F. and C. T. Fang (1984 a)

Fusarium diseases of broad bean.
II. Further studies on broad bean wilt caused
by Fusarium oxysporum.
Phytopathology, 38 : 331.

- Yu, T.F. and C.T. Fang (1984 b)

Fusarium diseases of broad bean.
III- Root-rot and wilt of broad bean caused
by two new forms of Fusarium.
Phytopathology, 38 : 587.

- Zgorkiewiez, A. (1963)

Results of investigation on vascular wilt of
lupine in Poland.

Biul. Inst. Ochr. Roal. Poznan, 24:389-404

(c.f. R.A.M, 43:314, 1964)