

REFERENCES

- Abdalla, A.M.(2002).**Effect of bio- and mineral phosphorus fertilizer on the growth, productivity and nutritional value of faba bean. Egyptian J. Hort ., 29(2): 187-203.
- Abd El-Aziz, El-Set. A.-A. and N.A. Anton (1999).** Influence of applying methods and micronutrient on growth, yield and quality of faba bean. Zagazig J.Agric.Res., 26 (5):1245-1257.
- Abd-El-Gawad,A.A.; D.M. El-Hariri ; A.M.A. Abo-Shetaia and,A.A. Bahr (1993).** Yield and yield components responses of chickpea (*Cicer arietinum*, L.) to phosphorus fertilization and micronutrients. African J. Agric. Sci., 18(1):61-71.
- Abd El-Lateef, E. M.; M. M. Selim and T. G. Behairy (1998).** Response of some oil crops to bio-fertilization with phosphate dissolving bacteria associated with different levels of phosphate fertilization. Bull. NRC. Egypt, 23(2): 193 – 202.
- Abdel-Wahab, A.F.; M.A. El-Deeb; A.A. Ragab and M.M. El-Sawi (1999).** Effect of soil bio-conditioner and NPK on nodulation, growth and yields of faba bean at the newly reclaimed soils. J. Agric. Sci. Mansoura Univ., 24(11): 6295-6301.

- Abo-Shetaia, A. M. and Mokhtar, Soheir. A. M. (2001).**Yield and yield components responses of chickpea (*Cicer arietinum*, L.) to phosphorus fertilization and micronutrients. Arab Univ. J. Agric. Sci., 9(1): 235-243.
- Agrawal, M. M. and C. P. Sharma (2005).**Effect of sulphur and molybdenum with *Rhizobium* and P-solubilizer (PSB) on yield and nutrient uptake in chickpea.Farm Sci., 14(2): 20-22.
- Ahmed, M. K. A.; M. H.Afifi and M. F. Mohamed (2003).** Effect of biofertilizers, chemical and organic fertilizers on growth, yield and quality of some leguminous crops. Egyptian J. Agron., (25): 45-52.
- Ajay, G.; V.K .Sharma; G.D. Sharma and P.Chopra(2006).** Effect of biofertilizer and phosphorus levels on yield attributes, yield and quality of urd bean (*Vigna mungo*).India.J. Agro.,51(2): 142-144 .
- A.O.A.C. (1970) .** Association of Official Agricultural Chemists. Official Methods of Analysis. Washington 11th ED. D.C. PP.832.
- Arf, O.; M.E.S.Buzetti; G.N.Guerreiro and J.A.Gutierrez (1994).** Inoculation, nitrogen fertilizer application and micronutrient application to increase yield of beans (*Phaseolus vulgaris*, L.) in the selviria region,Mato .Cultura,Agro.,3(1):151-166.
- Azad, A.S; J.S. Manchanda; A.S .Gill and S.S .Bains(1993).** Effect of zinc application on grain yield, yield components and nutrient content of Lentil.Lens Newsletter, 20(2): 30-33.

- Bahr, Amany. A. (1997).** Response of chickpea crop to some fertilization treatments. Ph.D.Thesis, Fac. Agric., Suez Canal Univ.,Egypt.
- Bhuiyan, M. A. H; D. Khanam; M.R. Khatun and M.S. Hassan (1998).** Effect of molybdenum, boron and Rhizobium on nodulation growth and yield of chickpea. Bull. of the institute of tropical Agric. Kyushu Uni. (2):1-7.
- Braga, N.R. and C. Vieira (1998).** Effect of inoculation with Bradyrhizobium sp., nitrogen and micronutrients on chickpea yield. Bragantia, 57(2) : 349-353.
- Chandra, R. (1995).** Response and economics of rhizobium, phosphorus and Zinc application in gram in Western U.P. plains. Legume Res., 8 (2) : 103-108.
- El-Awag, T.L.; A.M. Hanna and I.M. El-Naggar (1993).** Influence of bio and mineral phosphate fertilization on soybean production and some characters under different levels of soil moisture. Egypt. J. Appl. Sci., 8(11): 575-594.
- El-Awag, T. L.(1998).** Effect of soil moisture levels, Rhizobium inoculation and nitrogen application on broad bean yield. J. Agric. Sci. Mansoura Univ., 23(10): 4695-4704.
- El-Gizawy, N.K.B. and S.A.S. Mehasen(2004).** Yield and seed quality response of chickpea to inoculation with phosphorus fertilizer and spraying with iron. The 4 th Scientific Confer. of Agric. Sci. Assuit, December, 2004.

- El-Hadi, E.A. and E.A. El-Sheikh (1999).** Effect of Rhizobium inoculation and nitrogen fertilization on yield and protein content of six chickpea (*Cicer arietinum*, L.) cultivars in marginal soils under irrigation. Nutrient Cycling in Agroeco Systems.; 54(1):57-63. (C.F. Sammaries of Monograph, Record 1779 of 210- CAB Abstr. 1998-2000).
- El-Kalla, S.E.; A.K. Mostafa; A.A. Leilah and R. A. Awad (1999).** Mineral and bio-phosphatic fertilization for intercropped faba bean and onion. Egypt. J. Agric. Res., 77(1): 253-271.
- El-Karamany, M. F. and Amany. A. Bahr (1999).** Effect of mineral fertilization, organic manuring and biofertilization on yield and yield components of chick pea (*Cicer arietinum*, L.) cultivars in sandy soil . Egypt. J. Appl. Sci., 14(11): 1999.
- Enania, A.R. and A.K. Vyas (1994).** Effect of phosphorus and zinc application on growth, biomass and nutrient uptake by chickpea in calcareous soil. Annals of Agric. Res., 15(4): 397-399.
- Gangwar, K.S. and N.P. Singh (1986).** Effect of zinc fertilization on lentil yields and yields attributes . Lens Newsletter, 13(1):17-20.
- Gangwar, K. S. and N.P. Singh (1994).** Studies on zinc nutrition on lentil in relation to dry matter accumulation, yield and N,P uptake . Indian J. pulses Res., 7(1): 33-35.

- Gautam,V.S.; O.P. Girothia and S. Olkar (1995).**Performance of Bengal gram genotypes under integrated fertility management practices. crop. Res. Hisar. ; 10(3):247-252. (C.F. Sammaries of monograph, Record 146 of 210- CAB Abstr.1996-1998).
- Gupta, O.(1998).**Effect of micronutrients on growth of some root pathogens of chickpea.International Chickpea and Pigeonpea Newsletter.,(5): 14-15.
- Gupta, S.C. and S.L. Namdeo (1996).**Effect of rhizobium inoculation on symbiotic traits, grain yield and quality of chickpea genotypes under rainfed conditions.Crop Res.Hisar, 12(2):127-132.
- Hafiz, S.I.(1999).**Productivity of some chickpea varieties as influenced by plant density under sandy soil conditions. Annals of Agric. Sc., Moshtohor, 37(3):1605-1617.
- Hafiz, S. I. (2004).** Response of chickpea crop to biofertilization and foliar spraying with zinc under different levels of N and P fertilization in newly reclaimed sandy soils. Annals of Agric. Sci. Moshtohor., 42(3): 933-948.
- Hamed, M.F. (2003).** Faba bean as affected by zinc,phosphorus fertilizer and phosphoren. Annals of Agric. Sci. Moshtohor., 41 (3):1047-1056.
- Hamissa, A.M.; F.M. Hammouda and R.E. Knany (2000).** Response of nodulated faba bean crop to phosphate solubilizing bacteria under phosphorus fertilization and copper foliar spray application. J. Agric. Sci.Mansoura Univ.,25(5):2995-3007.

- Hegazy, M.H.; D.N. Abdi and S.A.Genaidy (1993).** Effect of some micro-nutrients and methods of application and rhizobium inoculation on faba bean. *Egyption J. Agric. Res.*, 71(1): 21-33.
- Hegazy, M.A.; M. Monib; H. Amer and E. Shoher (1992).** Inoculation of faba bean with bionitrogen under Egyptian condition. First inter. African Conf. on Biofert., Cairo ,Egypt, 23-28March.
- Herbert, D.; P. Phlipps and R. Strange (1971).** Determination of total carbohydrates. *Methods in Microbial*, 58:209-344.
- Hernandez, L.G. and G.D. Hill (1983).** Effect of plant population and inoculation on yield and yield components of chickpea (*Cicer arietinum*, L.). *Proceedings Agronomy Society of New Zealand*, (13):75-79 .(C.A. Bahr 1997).
- Hossain, M.M. and M. Suman (2005).** Effect of biofertilizer and nitrogenous fertilizer on growth, yield and N-uptake of lentil (*Lens culinaris Medik.*). *International J. Sustainable Agric.technology.*, 1(6): 76-81.
- Hussein, A.H.A.; N.M. Abou-Zeid and M.E. Hassan (1991).** Effect of N, P fertilizers, rhizobium inoculation and seed fungicides on yield, yield components, nodulation and seed contents of faba bean. *Egypt. J. Agric. Res.*, 69(3): 695-708.

- Hussein, A. H.A.; S.A. Saleh and M.A. El-Deeb (1999).** Effect of rhizobium inoculation, on growth, nodulation and yield of faba bean cultivated in the newly reclaimed soils of middle Egypt. Bull. Fac. Agric., Cairo, Univ., 5-9 September.
- Islam,M.A.; U.K .Saha; M.S .Islam and R.R .Saha (1995).** Fertilizing chickpea for high yield with special emphasis on boron and molybdenum in grey floodplain soils of Bangladesh. Annals of Bangladesh Agric. Res.Institute, 5(1):43-49.
- Jain, P.C.; P.S. Kushwaha; U.S. Dhakad; H. Khan and S.K. Trivedi (1999).** Response of chickpea (*Cicer arietinum,L.*)to phosphorus and biofertilizer. Legume Res., 22(4):241-244.
- Jain, L. K. and P. Singh (2003).** Bio-fertilizers and phosphorus nutrition Growth and nutrient uptake of chickpea (*Cicer arietinum, L.*) as influenced Crop Research Hisar. India. J.article., 25(3):410-413.
- Jain, L. K.; P. Singh and J. K. Balyan(2006).**Productivity and profitability of chickpea (*Cicer arietinum, L.*) cultivation as influenced by biofertilizers and phosphorus fertilization.India. J. of Dryland Agric. Res. and Development, 21(1): 82-84.
- Jain, A.K.; S. Kumar and J.D.S. Panwar (2007).** Studies on Rhizobium with phosphorus and micronutrients on mung bean (*Vigna radiata L. Wilczek*) in relation to yield and yield attributes. Adva. in Plant Sci., 20(2): 373-376.

- Johnson, S.E.; J.G .Lauren; R.M. Welch and J.M. Duxbury (2005).** A comparison of the effects of micronutrient seed priming and soil fertilization on the mineral nutrition of chickpea (*Cicer arietinum*, L), lentil (*Lens culinaris*, L), rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Nepal. *Experimental Agric.*, 41(4): 427-448.
- Kabesh, M. O.; M. S. M. Saber and T.G. Behaiey (1987).** Utilization of biofertilizers in field crop production.1- Effect of the increasing rates of phosphate fertilizers with and without biofertilization on growth and yield of soybean plants in pot experiments.*Egypt.J.Agron.*,12 (1-2):123-130.
- Kale, N.Y.; P.L. Patil and B.C.Patil (1982).** A study on effect of Rhizobium and Azotobacter inoculation on nodulation, N₂ fixation and yield of gram (*Cicer arietinum*, L.). *Indian J. Microbiology*,22(3): 203-205.
- Kalyan,S.; I. Raok; J.S. Janwal; J.S. Bohra and k. Singh (1983).** Effect of phosphorus and bio-fertilizers on growth yield and quality chickpea (*C.arietinum*,L.) transaction of Ind. Soci. of Desert. technology and univ. center of Desert studies., 8(1):33-38.
- Kedar, P.; S. Kumar; P. Ram and J. P. S. Rathi (2005).** Effect of FYM and biofertilizer in conjunction with inorganic fertilizer on growth, yield and profit of chickpea (*Cicer arietinum*, L.).*Plant Archives*, 5 (2): 609-612.

- Khanal, N.; K.D. Joshi; D. Harris and S.P. Chand (2005).** Effect of micronutrient loading, soil application, and foliar sprays of organic extracts on grain legumes and vegetable crops under marginal farmers' conditions in Nepal. Micronutrients in South and South East Asia Proceedings of an International.
- Khune, N.N.; S.O. Kolte; A. J. Chitriv and P.D .Wanghar (1989).** Biofertilizers in Vidarbha-present and prospects. PKV Res. J., 13 (2) : 168 -169.
- Knany,R.E.; A.M.Masoud and A.M.Konsouh (2004).** Impact of phosphorus fertilization and some micronutrients spraying on seed yield and quality of new cowpea cultivars.2nd Inter.Conf. Hort. Sci., Fac. Agric., Kafr El-Sheik, Tanta Univ., Egypt (28) : 1035-1047.
- Kosgey,J.R.; B.A.Mckenzie and G.D.Hill (1993).** Rhizobium and nitrogen effects on chickpea sown on two dates in canterbury. Proceedings – ann.Conference.Agron-Soci. New-Zealand.,(23):87-92.
- Krishnareddy, S.V. and I.P.S.Ahlawat (1996).** Growth and yield response of lentil cultivars to phosphorus,zinc and biofertilizers.J. Agron. and Crop Sci.,177 (1): 49-59.
- Mahmoud, S.M.(1995).** Effect of inoculation of faba bean with four genetically marked strains of rhizobium leguminosarum on nodulation, nodule occupancy and yield. Assiut J. Agric. Sci., 26(4).

- Mahmoud, S.M. and L.A. El-Far (2000).** Effect of inoculation with Bradyrhizobium (Hypogaeae) and nitrogen fertilization on the productivity of peanut (*Arachis hypogaea* L.) grown on sandy calcareous soil. Assiut J. Agric. Sci., 31(3): 57-72.
- Maurya, B.B.; C.I. Sanoria and P.C. Ram (1987).** Combined culture treatment enhances nodulation, yield and quality of chickpea. Seeds and Farms, 13 (8):25-27.
- Meena , K. N. ; R. G. Pareek and R. S. Jat (2001).** Effect of phosphorus and biofertilizers on yield and quality of chickpea (*Cicer arietinum*, L.). An. Agric. Res., 22(3): 388-390.
- Meena, L.R.; R.K. Singh and R.C. Gautam (2003).** Yield and nutrient uptake of chickpea (*Cicer arietinum*, L.) as influenced by moisture conservation practices, phosphorus levels and bacterial inoculation. Legume, Res., 26 (2):125-127.
- Mehasen, S.A.S. and M.A.El-Ghozoli (2003).** Response of soybean plants to foliar application with iron and molybdenum and soil fertilization with rock phosphoate and phosphoate dissolving bacteria. Monufiya J. Agric. Res., 28(1):87-102.
- Menaria, B.L.; P.Singh and R.K. Nagar (2003).** Effect of nutrients and microbial inoculants on growth and yield of soybean (*Glycine max* (L.) Men-ill. J. Soil and Crops. 13(1): 14-17.

- Misra, S.K.; R.M. Upadhyay and V.N .Tiwari (2002).** Effects of salt and zinc on nodulation, leghaemoglobin and nitrogen content of rabi legumes. India. J. Pulses Res ., 15(2): 145-148 .
- Monib, M.; A. Higazy; M. E. Hassan and A.A. Ragab (1994).** Possibility of magnifying broad bean productivity in fertile Nile Valley soils. Annals Agric Sci., Ain-Shams Univ., Cairo 39(1): 53-63.
- Nassar, K.E.; S.A .Radwan and A.A .Rahmou (2002).** Effect of seed coating with some micronutrients on Faba bean (*Vicia faba*, L.). I - Effect on photosynthetic pigments micronutrient contents and plant growth characters. Egyptian J. Soil Sci., 42(2): 347-362.
- Negm, M.A.; R.G. Kerlous and Y.B. Besada (1998).** Different sources of nitrogen and rhizobium inoculation effect on peas growing on a calcareous soil. Egypt, J. Soil Sci. 38(1-4): 69-79
- Neweigy, N. A.; T. A. Tewfike; M. A.El-Sibaie and M. A. El-Sayed (2000).**Effect of biological fertilization by *Rhizobium* and phosphate dissolving bacteria on broad bean grown in sandy soil.Annals of Agric. Sci. Moshtohor.,38(4): 2125-2139.
- Pathak, S.; K. N.Namdeo; V. K. Chakrawarti and R. K.Tiwari (2003).** Effect of biofertilizers, diammonium phosphate and zinc sulphate on growth and yield of chickpea (*Cicer arietinum*, L.).Crop Res. Hisar., 26(1): 42-46.

- Pawar, K.B.; N.J.Bendre; R.B. Deshmukh and R.R. Perane (1997).**Field response of chickpea seed inoculation of Rhizobium strains to nodulation and grain yield.J. of Maharashtra. Agric.unvi.publ.1998.,22(3):370-371.
- Peksen,E. and A. Gulumser (1996).**Effects of seed inoculation with three different rhizobium strains on seed yield and protein content of chickpea cv. ILC 482.ondoku zmayis, Unive. ziraat. fakuletsi. Dergis., 11 (2):69-77. (C.F. Sammaries of Monograph,Record 123 of 210- CAB Abstr.1996-1998/07).
- Pramanik, K. and R. K. Singh (2003).**Effect of levels and mode of phosphorus application with and without biofertilizers on yield and nutrient uptake by chickpea (*Cicer arietinum*,L.). Annals of Agri. Res., 24 (4): 768-775.
- Ram,G.; B.S.Chandrakar; M.K.Misra and R.K.Katre(1984).** Effect of seed treatment with Rhizobium and dithane M-45 on plant height,nodulation and yield of chickpea. Indian J. Agric.Sci.,54(3):214-216.
- Ram, V., P. K. Yadav; A. P. S. Dohare and M. D. Lodhi (2007).** Effect of biofertilizers and phosphorus levels on growth and yield of chickpea (*Cicer arietinum*, L.)in rainfed area under bundelkhand conditions. Plant Archives.,7 (1):237-239.
- Rao, K.L.N.; P.J.R. Reddy; B. K. Mahalakshmi and C. L.N. Rao (2005).** Effect of plant growth regulators and micronutrients on flower abortion, pod setting and yield of chickpea. Annals of plant physiology.19 (1): 14-16.

- Reddy, N.R.N. and I.P.S. Ahlawat (1998).** Response of chickpea (*Cicer arietinum*, L.) genotypes to irrigation and fertilizers under late. sown conditions. Ind. J . Agron., 43(1):95-101.
- Rizk, T.Y.; A.A.A. Haleem and I.K.A. Mohamed (2006).** Effect of organic, bio and mineral fertilization treatments on yield, yield components and chemical contents of seeds of faba bean "*Vicia faba*, L". Egyptian J. Agric.-Res., 84(1): 153-163.
- Rodelas, B. J.; C.Gonzalez- Lopez; V.Salmeron and M.B. Martinez-toledo (1998).** Response of faba bean to combined inoculation with azotobacter and rhizobium leguminosarum bv. Viceae, Al. Soil Ecol., 12. 51-59.
- Roy, S.K.; S.M.L. Rahaman and A.B.M. Salahuddin (1995).** Effect of Rhizobium inoculation and nitrogen on nodulation growth and seed yield of gram(*Cicer arietinum*, L.). Ind.J.Agric.Sci., 65(12):853-857.
- Saleh, S.A.; M.A. El-Deeb and A.A. Ragab (2000).** Response of faba bean (*Vicia faba*, L.) to rhizobium inoculation was affected by nitrogen and phosphorus fertilization. Bull. Fac. Agric., Cairo Univ., (51): 17-30.
- Santos, O.S. and V. Estefanel (1986)** .Effects of micro-nutrients and sulphur on soybean seeds .Revista Centor De-ciencias Rurais .16:1,5-17.
- Sapatnekar, H. G. ; P. H. Rasal and P. L. Patil (2002).** Role of biofertilizers and P-sources in chickpea cultivation.J. of Maharashtra Agricultural Universities., 27(2): 225-226.

- Sattar, M.A.; A.K. Podder; M.C.Chanda; M. Rahman; A.K. Podder; H.c. Van; Z.N.T. Begum, T. Heulin and A. Hartmann (1996).** Rhizobial biofertilizers:the most promising BNF technology for increased grain legume production in Bangladesh.Bio.,nitr.,fix., sociated with rice production,Based on selected papers presented in the inter, symp., Dhaka, Bangladesh.
- Sawires, E. S. (2001).**Effect of phosphorus fertilization and micronutrients on yield and yield components of chickpea (*Cicer arietinum*, L.). Arab Universities J. Agric. Sci., 46(1): 155-164.
- Sendecor, G.W. and W.G. Cochran (1980).** Statistical methods. 6 th Iowa State Univ. press, Ames, Iowa, U.S.A., 369-375.
- Sharma, A.K.; S. Haribendra; S. Sushil; S. Roham; K.N. Namdeo; H. Singh; S. Singh and R. Singh (1989).** Response of gram (*Cicer arietinum*, L.) to rhizobial and N fertilizeation.Ind J. Agron., 34(3):381-383.
- Sharma, R.A. and B.B. Parmar (1997).** Influence of biofertilizers and indigenous sources of nutrients on nutrient uptake and productivity of rainfed soybean-gram cropping sequence .Crop, Res.,Hisar, 13(1):13-18.
- Singh, R. K.; D. N. Shukla and De Nirmal (2006).** Effect of biofertilizers, fertility level and weed management on weed growth and yield of late sown chickpea (*Cicer arietinum*,L.).India. J. Agric. Sci., 76(9): 561-563.

- Smith, G.; W.Wiebold ; T.L.Niblack; P.Scharf and D.Blevins (2000).** Yield components of soybean plants infected with soybean cyst nematode and sprayed with foliar applications of boron and magnesium.J plant Nutrition.23(6):827-834.
- Srivastava, P.C.; U. Khan and L.M. Pant (2006).**Effect of N and Zn interaction on nodulation, yields and nutrient concentrations of French bean inoculated with *Rhizobium leguminosarum* bv. *phaseoli* (Strain-9R). Crop Res. Hisar., 31(1): 43-51.
- Subba-Rao, N.S.; K.V.Tilak and C.S. Singh (1986).** Dual inoculation with *Rhizobium* sp.*Glomus Fasciculatum* enhances nodulation, yield and nitrogen fixation in chickpea (*Cicer arietinum*, L.). Plant and soil ,95 (3):351-359.
- Sunder, S.; B.L .Pareek and S.K .Sharma (2003).**Effect of phosphorus and zinc on dry matter, uptake of nutrients and quality of clusterbean [*Cyamopsis tetragonoloba* (L) Tabu.]. An. Agri. Res., 24(1): 195-196.
- Tellawi,A.; N.Haddad and B.Hattar (1986).** Effect of several *Rhizobium* strains on nodulation, nitrogen uptake and yield of chickpea (*Cicer arietinum*, L) .Zeitschrift-fur pflanzenernahrung und Bodenkund, 149 (3):314-322.
- Tippannavar, C.M.; S.A. Desai; K. Krishnamurthy and S.B. Patil (1992).** Performance of biofertilizers in northern dry zone of Karnataka. J. Maharashtra,Agric.,Univ., 17(3): 494-495.

- Tomar, R. K. S. and R.S. Raghumanshi (1995).**Effect of nutrients on yield of chickpea under irrigated condition. Bhartiya. Krishi. Anus and Han. Patrika., 10(1-2):7-10.
- Tomar, R.K.S.; K.N. Namdeo and J.S. Raghu (1996).**Efficacy of phosphate solubilizing bacteria biofertilizer with phosphorus on growth and yield of gram (*Cicer arietinum*, L.) Ind.J. Agron., 41(3):412-415.
- Ulukan, H.; M. Guler and M.S .Adak (2002).**Effect of zinc fertilization on yield and quality of faba bean (*Vicia faba*, L.). India.J. Agric.Sci., 72(12): 742-744.
- Yadav,R.P. and U.k.Shrivastava (1997).** Response of chickpea (*Cicer arietinum*,L.) to phosphorus and biofertilizer legume.Res.,20(2):137-140.
- Yakout, G.M.; M.H. Greish and V. Romheld (2001).** Response of faba bean crop to phosphatic, foliar and bio-fertilization under new reclaimed sandy soil conditions. (C.F.Record.2 of CAB Abstract, 200/08 – 2002/07).
- Yanni, Y.G.(1992).** Performance of chickpea,lentil and lupin nodulated with indigenous or inoculated rhizobio micropartners under in nitrogen, boron, cobalt and molybdenum fertilization schedules.World. J. Microbiology and Bio., 8 (6) : 607- 613.
- Zaghloul,R.A.; M.A. El-Ghozoli and S.A.S. Mehasen (2002).** Effect of dual inoculation (VA-mycorrhizae and Rhizobium) and zinc foliar application on growth and yield of mungbean.An. Agric.Sci.Cairo., 47(2): 501-525.

Zeidan, M.S.; M.O. Kabesh and M.S.M. Saber (2001).

Utilization of biofertilizers in field crop production. 14-
Effect of organic manuring and biofertilization on yield
and composition of two faba bean varieties cultivated in
a newly reclaimed soil. Egypt. J. Agron., (23): 47-57.