

6. REFERENCES

- Abbas, A. M. (1978): Growing and fattening of early weaning lambs. M.Sci. Thesis, Fac. Of Agric., Zagazig Univ., Egypt.
- Abdel-Reheem, I. A. (1994): Improvement of some Ossimi productive and reproductive traits through grossing with Chios breed sheep. M.Sc. Thesis, Fac., of Agric., El-Minia University.
- Aboul-Naga, A. M. and Afifi, E. A. (1980): Crossing sub-tropical Egyptian sheep with Hampshire to improve their lamb production. J. Agric. Sci. (Camb.), 95:365-370.
- Aboul-Naga, A. M. and El-Shobokshy, A. S. (1974): Fattening performance and carcass quality of Sufflok crossbred lambs compared with Ossimi. Agric. Res. Rev., 52:63.
- Aboul-Naga, A. M., El-Tawil, E., Galal, E. S. E., Labban, F. and Khishin, S. (1972): The effect of crossing Merino with Ossimi and Barki sheep on some productive traits. J. Agric. Sci. (Camb.), 78:275-280.
- Aboul-Naga, A. M., Labban, F. M., Ghonem, K. E. and Issawy, H. F. (1968): Effect of crossing Merino with Ossimi sheep on body measurements and development. J. Anim. Prod., U.A.R., 8(1-2):31-44.
- Aboul-Naga, A. M., Markotic, B. and Abdel-Salam, N. (1973): Early fattening of local sheep on different feeding systems. Alex. J. Agric. Res., 21(2):193-198.
- Akcapinar, H., Tekin, M. E., Kadak, R., Akmaz, A. and Müftüoğlu, S. (1992): Growth, fattening and carcass characteristics in Turkish Merino, German Blackheaded Mutton × Merino, Hampshire Down × Merino and Lincoln Longwool × Merino (F1) lambs. A. U. Veteriner Fakültesi Zootekni Anabilim Dalı. Ankara, Turkey. A.B.A.2(2):18-23.
-

- Akgündüz, V., AK, I., Koyuncu, M., Filya, I., Deligözoglu, F. and Tuncel, E. (1994): Fattening performance and carcass characteristics of F₁ carcass of meat type breeds and Kivircik sheep. *Koyunculuk Arastivma Enstitusu Müdürlü gü, Bandirma, Turki. A.B.A. 34(3-4) 48-64.*
- Aksoy, A. R. (1995): Fattening performance and carcass traits of Red Karaman and Tushin ram lambs at different slaughter weights. *Univ. Veteriner Fakültesi Zootekni Anabilim Dali. Kars, Turkey. A. B. A. 42(1):15-23.*
- Anderson, M. K., Field, R. A., Riley, M. L., McCormick, R. J., Snowden, G. D. and Bailey, D. G. (1991): Effect of age, castration and season on difficulty of pelt removal in lambs. *J. Anim. Sci., 69;3284-3291.*
- A.O.A.C. (1990): Official Methods of Analysis, 15th Ed. Association of Official Analytical Chemists. Inc. USA.
- Arik, Z., Alarslan, O. F., Kor, A., Ertugrul, M. and Polatsu, S. (1996): Slaughter and carcass traits of Malya and White Karaman lambs fattened on complete diets based on barley and different roughages.. *Ankara Univ., Veteriner Fakültesi, Zootekni Anabilim Dali, Ankara, Turkey. A. B. A., 36(2):68-87*
- Arnold, G. W. (1975): Herbage intake and grazing behaviour in ewes of four breeds at different physiologicat states. (*Aust. J. Agric. Res., 26:1017*).
- Arpacik, R., Aydogan and Özcelik, M. (1993): Comparison of live weight gain and feed consumption of Ile-de-France × Turkish Merino and Ile-de-France × White Karaman F₁ ram lambs. *Veteriner. Fakültesi, Zootekni Anabilim Dali, Ankara, Turkey. A.B.A., 17(3):187-192.*

- Arora, C. L. and Acharya, R. M. (1972): Factors affecting growth, carcass yield and wool yield in Indian breeds of sheep and their crosses. *Ind. Veter. J.*, 49(6):578-584.
- Arthur, P. F. and Ahunu, B. K. (1988): Features of growth and development in Ghanaian crossbred sheep. *Ind. J. Anim. Sci.*, 58(9):1098-1101.
- Asker, A. A., El-Kholsy, I. G. and Juma, K. H. (1964): Effect of docking on growth and development of Awassi lambs. *J. Anim. Prod. U. A. R.* 4:11.
- Asker, A. A., Ragab, M. T. and Bastawisy, A. E. (1954): Effect of crossing Egyptian sheep on growth and development of lambs. *Empire J, Exper. Agric.*, 22(88):256-260.
- Azamel, A. M. (1978): A study on fattening lambs on concentration and Alfa alfa. M. Sc. Thesis, Fac. of Agric., Zagazig Univ. Egypt.
- Azarov, P. V. (1985): Food conversion for body weight and wool in purebred and crossbred young sheep. *Razvedenie orets I Koz. Sherstovedenie*, 39-45 (Ru, 7 ref.).
- Babadaeva, L. M. (1974): Meat production of crossbred meat wool sheep. *A. B. A.*, 43 (10):4551.
- Badreldin, A. L. (1951): Growth and carcass percentage in Ossimi and Rahmani sheep. *Fac. of Agric. Cairo Univ., Bull. No. 3.*
- Bali, G. S., Das, S. A., Gopinthan, V. K. and Sharma, T. R. (1976): Preliminary studies on some characteristics of Bannur mutton, *J. of Food Sci. and Technol.*, 13(2):97-98.
-

- Barrton, R. A. (1967): A review of meat research at Annual Mossey University. Sheep farming Annual, Mossey Univ. 1967:127-139.
- Basuthakur, A. K. and Singh, O. N. (1969): A study on body weight and conformation of Coimbatore and Corriedal sheep. Ind. Vet. J., 46(12):1065-1068.
- Bicer, O., Güney, O. and Pekel, E. (1995): Effect of slaughter weight on carcass characteristics of Awassi male lambs. J. of Applied Anim. Res., Dep. of Anim. Sci., Univ. of Mustafa Kamal, Fac. of Agric., Antakya, Turkey. 8(1):85-90
- Bondarenko, G. M. (1971): Food conversion and daily gain in young crossbred and fine-wooled sheep. Outsevootive., Mesk., 16(8):34 (A. B. A., 39:4741).
- Bonsembiante, M., Andrighetto, I. and Cozzi, G. (1988): Effect of Finn sheep crossbreeding on Lamon sheep performance. Post-mortem traits. J. of Agric. Sci., in Finland, 60:515-517.
- Botkin, M. P., Field Ray, A. and Johnson, LeRoy, C. (1988): Science, production and management. Prentice Hall, New Jersey.
- Bouton, P. E., Harris, P. V. and Shorthose, W. R. (1971): Effect of ultimate pH, upon the water-holding capacity and tenderness of mutton. J. Food Sci., 36(3):435-439.
- Boylan, W. J., Berger, Y. M. and Allen, C. E. (1976): Carcass merit of Finn sheep crossbred lambs. J. Anim. Sci., 42:1413.
-

- Burghart, M. and Schmidt, L. (1966): Investigations on gross dissection of the carcass of Merino slaughter lambs. Bayer landw. Jb., 43:857-864. (A.B.A. 35(3):2567).
- Burns, R. H. (1961): Crossbreeding of sheep. Wool Short Course. Feb. 1 (1961).
- Capistrak, A., Malik, J. and Apolen, D. (1988): Carcass quality of three-breed crossbreds based on the Improved Valachian sheep. Czechoslovakia. A.B.A. 33(7):577-587.
- Carena, A. and Costa, F. (1971): Fattening and slaughter trial with improved Lucanian lambs and their crosses with Berrichon and precoce at different ages (42-91 and 140 days). A.B.A. 41(9):3964.
- Carpenter, Z. L. (1966): What is consumer-preferred lamb. J. Anim. Sci., 25:1232.
- Cattaneo, P. Bianchi, M. A., Beretta, G. and Cantoni, C. (1979): Keeping characteristics of lamb and its nutritive value. Industrie Alimentari, (C.F.F.S.) T.A., 12, 05 735, (1980):18(9):641-643.
- Cebetov, E. F. (1965): Change with age in the carcass components and meat production of weathers. Vestn. Sel. Hoz. Nauk., 10:116-120. (Abst. A, B. A. 33:3416).
- Charyulu, E. K. and Munirathnam, D. (1984 a): A comparative study on some pre-and post-slaughter parameters of Nellore, Mandya and their crosses with Dorest and Sufflok rams under feed lots. Ind. J. Anim. Sci., 54(2):175-179.
- Charyulu, E. K. and Munirathnam, D. (1984 b): Factors affecting birth, weaning and six months body weight in Mandya, Nellore and their crosses with Dorset and Sufflok rams. Ind. J. Anim. Sci., 54(7):709-710.
-

- Chatterjee, A. K., Balgia, B. R., Panda, B. and Khabade, V. S. (1969): Carcass evaluation of Bannur and non-Bannur lambs. Ind.Fd. Packer, 23:1-6 (A. B. A., 40(3):3168).
- Chawadhary, S. S. (1978): Evaluation of carcass traits in Noli, Lohi, Magra and Hissardole $\times$ Magra breeds of sheep. Thesis Abst., Haryana Agric. Univ. 4:185-186. (Abstr. A.B.A. 48 No. 1206).
- Chipeav, O. M. (1963): Lambs born in winter can be economically slaughtered at 8-9 months of age. (Abst. A. B. A., 32, 1076).
- Ciolca, N., Ursescu, A. Petcu, D., Avran, A. and Cusa, S. (1972): Evaluation of growth performance and carcass quality of lambs of local breeds. A.B.A. 41 (3):1116.
- Coetzee, C. G. (1972): Intensive mutton production from wool-bearing sheep determination of the optimum slaughter mass of early weaned sheep. A.B.A., 41 (11):4831.
- Cumlivski, S. (1959): Growth and body development of crossbreds from Valachian ewes and Wurttemberg rams. Sborn. Osl. Akad. Zmed., Ved. Zivoc. Vyr. 4(3):331-340. (A. B. A., 27:1376).
- Dag, B. and Ertugrul, M. (1993): Fattening performance and carcass traits of Karayaka and Border Leicester mult Karayaka (F_1) crossbred male lambs. S.U. Ziraat Fakültesi, Zootekni Bölümü, Konya, Turkey. A.B.A. 33(3-4):42-57.
- Dally, R. A. (1969): A note on the fat content of lean meat. L J. of the Association of Public Analysts, 7(1):33.

- Darwish, M. Y. H. (1962): Meat production. First edition, Cairo, in Arabic.
- Darwish, M. Y. H., El-Samman, S. and Abou-Hussein, E. R. M. (1973): Meat production from Rahmani lambs. Egypt. J. Anim. Prod., 13(1):35-48.
- Darwish, M. Y. H., El-Samman, S. and Abou-Hussein, E. R. M. (1975): Meat production from Rahmani lambs. A. B. A., 43(1):203.
- Das Kornel and Surindra Kumar (1987): Body weight confirmation of imported and farm-bred Australian Corriedale sheep in India. Ind. J. Anim. Sci., 57(2):153-154.
- Dattilo, M., Congiu, F. Rossi, G. and Brandano, P. (1975): Comparative study of physico-chemical characteristics of meat and fat of different age and sexes, Attidella Societa Italiana dell Scienze Veterinarie, (c.f. F. S. T. A.) Vol. 7 (1975). 27: 409-414.
- Dinescu, S. (1972): Results obtained from crossing Lincoln and South down rams with Turcana ewes. A. B. A., 41(3):1120.
- Druzenjkov, G. I. and Oleinik, T. I. (1970): Tien Shen semifine woolled sheep as improvers of carcass characters in crossbred. A. B. A., 39(1):482.
- Duncan, D. B. (1955): Multiple range and multiple F tests. Biometrics, 11:1-42.
- Efner, T. (1972):The slaughter yiely of wethers killed at different age. A. B. A., 40(4):4552.

- El-Fiky, F. A. A. (1984): The effect of early weaning of lambs on mutton production. M. Sci. Thesis, Fac. Agric. Al-Azhar Univ. Cairo.
- El-Hamadani, O. T. (1974): Effect of crossing on feed efficiency in sheep. M.Sci. Thesis, Fac. of Agric., Ain Shams Univ., Cairo.
- El-Kimary, A. F. (1975): Crossbreeding of sheep under Farm Condition Ph.D. Thesis. Fac. of Agric., Alex. Univ.
- El-Kouni, M. H., Karam, H. A., Galal, E. S. E. and Afifi, E. A. (1974): Crossbreeding and the effect of certain environmental factors on body weight of Barki and German Merino sheep. J. Agric. Sci. (Camb.), 82:349-352.
- El-Mahdy, M. R. (1985): Studies on carcass evaluation in lambs. M. Sc. Thesis, Fac. of Agric. Zagazig Univ., Benha Branch, Moshtohor, Egypt.
- El-Qobati, A. A. (1990): Studies on some productive traits in sheep. M.Sc. Thesis of Agric. Zagazig Univ., Benha Branch, Moshtohor, Egypt.
- El-Said, A. I. (1983): Studies on some production traits in sheep. M. Sc. Thesis, Zagazig Univ., Benha Branch, Fac. of Agric., Moshtohor, Egypt.
- El-Sayed, A. I., El-Mahdy, M. R. and Soliman, A. S. (1995): Body weights and measurements of Ossimi and Rahmani $\times$ Ossimi Crossbred lambs as affected by some factors. Annals of Agric. Sci., Moshtohor, 33(4):1251-1263.
- El-Shaer, H. M., Salem, O. A. and Khamis, H. S. (1992): Fattening sheep on ration of broiler litter ensiled with green Bersem (*Trifolium alexandrinum* L.). 2- Animal performance and carcass traits. Egypt. J. Anim. Prod., 29(2):263-272.

- El-Shahat, A. A., El-Basiony, A. Z., Abdel-Rahman, G. A. and Aly, H. M. (1986): Effect of interaction between breed and plane of nutrition on sheep carcass tissues. *Egypt. J. Anim. Prod.*, 26(2):129-135.
- El-Shahat, A. A. and El-Ashry, M. A. (1987): Specific gravity measurements of lambs carcass joints. *Beitrage Zur Tropischen Landwirtschaft und Veterinarmedizin.*, 25(4):439-442.(A.B.A., 1988, 56:4417).
- Eltya, J. and Juma, K. H. (1970): Weight and body dimensions in Awassi sheep. *Iraqi J. Agric. Sci.*, 5:3-8 (A.B.A., 40:1350).
- Epstein, H. (1961): The development and body composition of docked and undocked fat-tailed Awassi lambs. *Empire J. Exper. Agric.*, 29(114):110-118.
- Epstein, H. and Herz, A. (1968): The body composition of 3-month-old Rambouillet lambs of different live weights. *J. Agric., Sci., (Cam.)*, 70(1):29-32.
- Ermekov, M. A. and Shrobogotov, Ju. A. (1965): The meat characters of Semi-fine woolled lambs reared in the semi desert zone of South Kazakhstan. (Abst. A.B.A.), 33:3418.
- Fahmy, M. H., Galal, E. S. E., Ghonem, Y. S. and Khishin, S. S. (1969) :Crossbreeding of sheep under semi-arid conditions. *J. Anim. Prod.*, 11(3):351-360.
- Falvey, L. and Hengmichai, P. (1979): Carcass studies of small ruminants in the northern highlands. *Thai. J. Agric. Sci.*, 12(4):301-308.

- Farhan, S. M. A., Al-Khalisi, I. J. and Hameed, M. A. (1969): Effect of docking on growth and fattening of Awassi lambs. *Iraqi, J. Agric. Sci.*, 4(2):57-65.
- Farid, A., Makarechian, M. and Sefidbakht, N. (1977): Crossbreeding of Iranian-fat-tailed sheep. Lamb performance of Karakul, Mehraban and Naeini breeds. *J. Anim. Sci.*, 44:542-547.
- Faten, F. Mohamed (1986): Studies on lamb production from some crossbreeding system. M. Sc. Thesis, Fac., of Agric. Zagazig Univ., Egypt.
- Faten, F. Mohamed (1992): Effect of ration and breed on the distribution of fat in sheep. Ph. D. Thesis, Fac., of Agric. Ain Shams Univ., Egypt.
- Fernandes, A. P., Thorat, B. P. and Deshmukh, A. P. (1988): Effect of crossbreeding on growth performance, carcass traits and meat characteristics in Deccani (native) sheep. 3rd World Congress on sheep and Beef Cattle Breeding. A.B.A. 1:483-486.
- Fogarty, N. M., Hall, D. G. and Atkinson, W. R. (1992): Management of highly fecund ewe types and their lambs for 8-monthly lambing. 2- Effect of weaning age and sex on lamb growth and carcass traits. *Aust. J. of Exp. Agric. (Aust.)*, 32(8):1031-1036.
- Gaili, E. S. E. and Ali, A. E. (1985): Meat from Sudon Desert sheep and Goats: Part 1 carcass yield, offals and distribution of carcass tissues. *Meat Sci.*, 13:217-227.
- Gaili, E. S. E. (1993): A comparison of body components of indigenous Saudi sheep breeds related to relative size and growth rate. *Anim. Prod.*, 57:113-118.

- Galal, E. S. E. and Ghanem, Y. S. (1970): The interaction between breeds of sheep and pattern on production and management. U.A.R., J. Anim. Prod., 10(1):7-12.
- Galal, E. S. E., Seoudy, A. M., Younis, A. A. and Khishin, E. S. (1971): Feed lot performance and carcass characteristics of yearling Barki sheep and their crosses with Merino. Alex. J. of Agric. Res. 19:15-24.
- Galal, E. S. E., Ghonem, Y. S., Fahmy, M. H., Seoudy, A. M. and Farid, N. A. (1975): Carcass traits and feedlot performance of Barki, Merino and Awassi breeds of sheep and some of their crosses. Egypt. J. Anim. Prod., 15:33.
- Galmez, J. and Santisteban, E. (1971): Effect of body weight at slaughter on carcass weights yield and composition in French precoce lambs. A.B.A., 40(3):3173.
- Ghanekar. V. M., Soman, B. V. and Bhatawadekar, S. D. (1972): A note on some vital and meat characteristics in Bannur sheep. Ind. J. Anim. Sci., 42(8):584-586.
- Ghoniem, K. E., Aboul-Naga, A. M. and Labban, E. (1968): Effect of crossing Merino with Ossimi sheep on growth and body weight. Egypt. J. Anim. Prod., 8:45-56.
- Ghoneim, K. E. (1973): Studies on commercial Karadi flocks of sheep in Northern. 11. Body weights and dimensions. Publication No. 24, Hamman Al-Alii, Mosul Univ. (52) PP.
- Ghoneim, K. E., Taha, A., Taka, M. R., Abdallah, R. K. and Kazzal, N. T. (1973): Some economic characters of Awassi sheep in North of Iraq. Publication No. 22 United Nations. FAO. Special Fund Project (Iraq 69/515).

- Gohler, H. (1970): The fattening of lambs with special reference to the estimation of carcass value. A.B.A., 39(2):1885.
- Goot, H., Hasdai, A., Bor, A., Zenou, A. and Gootwine, E. (1988): Body and carcass composition of fat-tailed Awassi and Booroola-Awassi F₁ ram lambs. 3rd World Congress on sheep and Beef Cattle Breeding. 1:491-493.
- Grumbach, S. and Nürnberg, K. (1996): Fattening procedures for lambs. Germany A. B. A. 8:70-72.
- Guney, O. (1990): Commercial crossbreeding between Ile-de-France, Rambouillet, Chios and local fat-tail Awassi for market lamb production. Small Ruminant Res., 3(6):449-456.
- Hamada, M. K. O. and Badawi, A. M. (1955): A study of birth weight in the Egyptian Rahmani sheep. Alex. J. Agric. Res., 3(2):137-146.
- Hamada, M. K. O. (1976): Variations in correlated weights of Rahmani lambs. World Revi. Anim. Prod., XII (1):51-60.
- Hammond, J. (1932): "Growth and the development of mutton qualities in the sheep". 2nd ed. Oliver and Boyd, London.
- Hammond, J. J. R., Bowman, J. C. and Robinson, T. J. (1983): Farm Animals. Fifth Edition. Edward Arnold.
- Hassan, H. A. (1984): Studies on some of the Egyptian sheep and their crosses under-Minia environmental conditions. Ph.D. Thesis in Anim. Prod. Fac. of Agric. El-Minia Univ.
-

- Hassan, H. A. (1988): Effects of crossbreeding and certain environmental factors on twinning percentage in sheep. *Minia J. Agric. Res. and Dev.* 10(3):1441-1452.
- Hassan, H. A. (1993): The effect of crossing Chios rams with Ossimi and Saidi ewes on growth performance and viability of lambs. *Egyptian J. Anim. Prod.*, 30(1):39-53.
- Hassan, H. A. and El-Feel, F. M. R. (1988): The effect of crossing among Ossimi, Barki and Saidi sheep and some other factors on body weight and daily gain of lambs. *Minia J. Agric. Res. and Dev.*, 10(4): 1607-1628.
- Hassan, H. A. and El-Feel, F. M. R. (1991): The effect of breed, level of feeding, age and slaughter weight on performance and carcass traits of lambs. *Egypt. J. Anim. Prod.*, 28(2):225-268.
- Hassona, E. M. A. (1980): Some studies on sheep performance fed different levels of feeding. M. Sc. Thesis, Fac. of Agric. Zagazig Univ. Egypt.
- Havitonov, K. A. (1969): Evaluation of Romney Marsh and Romney Marsh × Tsigai rams on the fattening performance and food conversion of their progeny. *Inst. Zhivot.*, 15:81-82. (A.B.A., 39:3324).
- Hofmeyr, H. S. (1975): Efficiency of feed utilization in farm animals: Study of the that influence the total efficiency of energy utilization in different sheep breeds. *Agric. Res. Pretoria, South Africa, Dep. of Agric. Tech. Serv.* (1975):63-64.
- Huidolra, F. and Jurado, J. J. (1989): Meat production of crossbred Manchega sheep. *Sanidad, Anim.*, 4, 1:35-44.

- Husaini, S. A., Deatherage, F. F. and Kunkle, L. E. (1950): Studies and meat, II- Observation on relation of biochemical factors to changes in tenderness. *Food technol.*, 4:366-369.
- Ibrahim, M. K. El-Sayed, A. I., El-Mahdy, M. R. and Hamdieh, H. Khalil (1994 a): Carcass traits and non-carcass components weight of Ossimi and Crossbred lambs in relation to pre-slaughter weight. *Annals of Agric. Sci., Moshtohor*, 32(3):1461-1471.
- Ibrahim, M. K. El-Sayed, A. I., El-Mahdy, M. R. and Soliman, A. S. (1994b): Studies on some productive traits in sheep. 1- Evaluation of carcass characteristics of lambs. *Annals of Agric. Sci., Moshtohor*, 32(3):1473-1488.
- Ibrahim, M. K., Hamdieh, H. Khalil and El-Qubati, A. A. (1991 a): Studies on some productive traits in sheep. 1- some factors affecting body measurements of sheep. *Annals of Agric. Sci., Moshtohor*, 29(1):167-185.
- Ibrahim, M. K., Hamdieh, H. Khalil and El-Qubati, A. A. (1991 b): Studies on some productive traits in sheep. II- Prediction of carcass traits from body weight and measurements. *Annals of Agric. Sci., Moshtohor*, 29(1):187-198.
- Imanalinov, S. T. (1973): Tajik and Saraja rams-successful improves of Kazakh fat-rumped sheep. *A.B.A.*, 42(2):587.
- Imam, S. A. (1962): Genetic improvement in body weight and growth rate in a flock of Ossimi sheep. *M. Sci. Thesis, Fac, of Agric. Cairo Univ.*
- Jamison, H. K., Carter, R. C., Gains, J. A. and Kincaid, C. M. (1961): The effect of breed of sire on body size of lambs at birth. *J. Anim. Sci.*, 20:154-158.

- Jensen, N. E. (1987): Performance testing of ram lambs. Individprover med lam Beretning fra statens Husdyrbvugs forsog. Statens Husdyrbvugsforsog, Foulum, Tjele, Denmark. No. 632, 38 PP (Da, en)
- Jensen, N. E. (1989):[Individual testing of ram lambs in (1989.3.)]. Individprover of vaedderlam. 111-Tidsskrift for Fareavl, 54 (10):7-9 (Da).
- Jensen, N. E. (1990): (Performance testing of ram lambs in 1990, 2:) Individprover of vaedderlam. II- Tidsskrift for Fareavl, Statens Husdyrbvugsforsog, Foulum, Tjele, Denmark. 55 (9):21-23 (Da)
- Jensen, N. E. (1992): Performance test of ram lambs in 1992. A.B.A. No. 713, 41 pp.
- Jensen, N. E. (1993): Performance testing of ram lambs in 1993. Denmark. A.B.A. 58(9). 4-8 (Da).
- Kadak, R., Akcapinar, H., Tekin, M. E., Akmaz, A. and Müftüoglu, S. (1993): Fattening and carcass characters of F₁ German Blackheaded Mutton×Akkaraman, Hampshire Down × Akkaraman, German Blackheaded Mutton×Awassi and Hampshire Down × Awassi lambs. Hayvancilik Merkez Arastirma Institisti Müdürlüğü, konya, Turkey. A.B.A. 3(1):1-7.
- Khalil, I. (1966): Factors affecting lamb production. M.Sc. Thesis. Fac. of Agric., Cairo Univ., Egypt.
- Kirton, A. H., Hight, G. K. and Duganzich, D. M. (1967): A comparison of the carcass quality of Romney with Border Leicester ×Romney lambs and South-down×Romney with South down×(Border Leicester×Romney) lambs. N.Z.J. Agric. Res., 10:33-42.

- Kirton, A. H., Carter, A. H. Ciarke, J. N., Sinclair, D. P. and Jury, K. E. (1968): Breeds of sire for export lamb production. Sheep farming Annual Mossey Univ., 39-48.
- Kleemann, Do., Dolling, CHS. And Ponzoni, R. W. (1990): Carcass and non carcass characteristics of Suffolk-sired lambs from South Australian Merino, Poll Dorest × Merion and Border Leicester × Merino ewes. Small Ruminat Res., 283.
- Koch, R. M., Swiger, L. A., Chambers, D. and Gregory, K. E. (1963): Efficiency of feed use in beef cattle. J. Anim. Sci., 22:486.
- Koncar, L. (1957): Comparisons of the bony development in the first five months of F₁ crossbreds of varieties of Pramenkes and Merinos. Veterinaria, 6:549 (A.B.A., 26:1401).
- Koretskii, P. S., Batchaeu, S. U. SH., Karadogeu, A. K. and Gochiayer, KH. N. (1984): Some results of the use of Australian rams on Soviet Mutton-wool ewes. Razredenie Ovesti Koz. Sherstoredenie, 8-12 (Ru, 6 ref.).
- Krilova, N. N. and Liskovskaia, U. N. (1961): Physical and chemical methods of analysis of animal products, Food ind. Pub., Moscow.
- Laban, F. M. and Radwan, A. (1963): The effect of crossing Egyptian sheep with Merino on live weight of lambs. Proceedings of the 2nd Anim, Prod. Conference, Cairo, vol., 1:47-59.
- Labecka, S. (1973): Meat characters of Romney Marsh sheep kept under production conditions in West Pomerania. A. B. A., 43(8):3421.

- Labusca, I., Plamadeala, C., Labusca, S., Grigori, I. and Pepa, I. (1976): Fattening performance of different type of sheep at the Movileni intercooperative center. *Zootehnie-Medicina Veterinaria* (1976):61-65 (A.B.A., 44:1182).
- Lane, S. F. and Kemp, R. A. (1990): The effects of shearing and Ralgro implants on feedlot performance of lambs during the summer. *Canadian J. of Anim. Sci.* 70(3):743-750.
- Latham, S. D., Moody, W. G. and Kemp, J. D. (1966): Techniques for estimating lamb carcass composition. *J. Anim. Sci.*, 25:492-496.
- Lawrie, R. A. (1985): *Meat Science*, Fourth Edition. Pergamon Press. New York.
- Layla, B. Bahgat (1991): Studies on crossbred sheep. Ph.D. thesis, Fac., of Agric. Zagazig Univ., Egypt.
- Lewczuk, A. (1975): Comparative studies on growth and carcass quality of purebreds Merino lambs and Lincoln $\times$ polish Merino crossbreds. Pt., 2. Growth and carcass quality of crossbred and purebred Merino wethers aged 180 or 240 days. *Roczniki Nauk Rolniczych*, B. 96(4):53-63. (A.B.A., 44(5) No.:2153).
- Lohse, B. and Kallweit, E. (1968): Characters for the estimation of carcass composition in Blackheaded Mutton fat lambs. *A.B.A.*, 37(1):462.
- Mabrouk, M. M. S., El-Oksh, H. A., Afifi, E. A. and Kadry, A. E. H. (1983): Lambs growth and carcass merit of Barki, Ossimi and Rahmani and their cross with Ile-de-France. *Al-Azhar Agric. Res. Bull.* No.95:1-25.

- Makarechian, M., Farid, A., Sefidbakht, N. and Mostafavi, M. S. (1975): The influence of breed and weaning age on feedlot performance of Iranian fet-tailed sheep. *Iranian. J. of Agric. Res.*, 2:21-29 (A.B.A., 43:1108).
- Marai, I. F. M., Daader, A. H. and Bahagat, L. B. (1998): A trial for evaluation of the Egyptian program for improvement of sheep. *Dep. of Anim. prod., Fac. of Agric., Zagazig Univ., Zagazig, Egypt.*
- Marzouk, K. M. (1986): Studies on productive and reproductive traits of Ossimi, Barki and Saidi sheep and their crosses under El-Minia environmental conditions. *M. Sci. Thesis Fac. of Agric. El-Minia Univ.*
- Meat Research Institute (1971 a): *Photographic Judging Standards lambs; London Agric. Res. Councile.*
- Meat Research Institute (1971 b): *Lamb Judging Score Card; London Agric. Res. Councile.*
- Minev, p., Katsarov, Ya., Dobrev, D. and Boshnakov, I. (1971): Body conformation in sheep. *Zhivot. Mouk.* 0(6):73-80 (A.B.A., 40:4535).
- Ministry of Agric., Egypt, (1968): *Animal and Poultry Nutrition. Bull. No. 3.*
- Mirzabekov, S. SH. And Dulsemin, D. (1988): *Carcass quality of lambs.. A.B.A. No. 3:30-31 (Ru).*
- Mokhtar, M. M. (1974): Effect of length of fattening period on gain and carcass traits in sheep. *M. Sci. Thesis, Fac. Of Agric. Ain Shams Univ., Egypty.*
- Moskalenko, A. N. (1970): Meat production of young crossbred in relation to type of crossbreeding. *A. B. A., 39(4):4684.*

- Mousa, T. M. (1991): Effect of crossing Ossimi, Awassi and Chios sheep on some productive traits. Ph.D. thesis in Anim. Prod. Fac. of Agric. Assiut Univ.
- Mowafy, M. A. (1968): Comprative studies on the meat quality of the local and imported sheep. M.Sc. Thesis, Fac. of Agric., Ain-Shams Univ.
- Mutaev, M. and Gulin, P. A. (1989): Intensive finishing of lambs in the non-chernozem region. *Zootekhnika*, No. 7, 48-50 (Ru).
- Netyaga, M. I., Verbina, V. D. and Makhmetov, E. Kh. (1992): Production traits of Australian-North Kazakh Merinos. *A.B.A.* 3:13-15 (Ru).
- Neville, W. F., Pope, A. L. and Chapman, A. B. (1955): The effect of breed of two size on dimensions of lambs at birth. *J. Anim., Sci.*, (A.B.A.), 14:1187.
- Nianogo, A. J. and Lankoande, L. (1991): Effect of early weaning on growth of West African Dwarf lambs. *Univ. de Ouagadougou, Upper volta. A. B. A.*, 39(3):327-328.
- Oliveira, N. M. DE, Osorio, J. C. and Monteiro, E. M. (1996): Meat production of lambs of five breed types. 1. Growth and development. EMBRAPA, Centro de pesquisa de pecuaria dos Campos Sulbrasilereiros, Bage, Rs, Brazil. *A. B. A.*, 26(3):467-470
- Oltenacu Branford, E. A. and Boylan, W. J. (1981): Productivity of purebred and crossbred Finnsheep. II. Lamb weights and production indices of ewes. *J. Anim. Sci.*, 52:998-1006.
- Osman, A. H. and El-Shafai, S. A. (1967): Carcass characteristics of Sudan Desert sheep. *Sudan J. Vet. Sci., Anim. Husb.*, 8:115-119.

- Osorio, J. C., Oliveira, N. M. DE., Nunes, A. and Pouey, J. (1996): Meat production in lambs of five breed types. 3. Losses and morphology. Brazil, A. B. A., 26(3):477-481.
- Ospanov, B. K., Tsol, L. F. and Ten, V. M. (1978): Meat production of lambs from sheep inseminated at 7.5 to 9 months of age. Vestnik Sel'sk-Khozyaistvennoi Nauki Kazahstana, 21(9):17-18. (A. B.A., 43:705).
- Palian, B. and Bayaric, D. (1959): Differences in the development during Suckling of twin and single Pramenka lambs at Kupris. Veterinaria (Sarajevo) 8:55-68. (A.B.A., 27:424).
- Pilar, R. DE. C., Pires, C. C., Restle, J., Silveira, S. DAS, E., Goncalves, J. M. and Fernandes, F. (1994): Growth and carcass composition of confined sheep of different breed types, slaughtered at 12 months of age. Brazil. A.B.A. 24(3):607-612.
- Plemyannikov, A. G. and Ilyaledinov, E. V. (1975): Breed differences in the utilization of nutrients during fattening of crossbred lambs. Ovtsevodstvo (9)., 17-18. (A.B.A., 43:705).
- Prasad, R. D. D., Charyulu, E. K., Sreenivasulu, D., Rao, T. M. and Munirathnam, D. (1991): Carcass characteristics of Mandya, Nellore and their halfbreds with Dorset under feedlots. Andhra Pradesh Agric. Univ., Palamaner, Andhra Pradesh 517-408, Ind. A.B.A., 61(2):226-227.
- Raichev, S. and Stankov, I. (1990): Finishing and carcass traits of semifine-wooled commercial crossbred lambs. Stara Zagora, Bulgaria. A.B.A., 27(8):26-31.

- Ramano, M. J. L., Martinez, R. L. and Rodriguez, V. A. (1990): Carcass measurements and Omental fat in sheep bred for hides and wool in different environment. Memoria, III., Congreso Nacional de production Ovina, 100-102.
- Raziq, F., Ahmed, I., and Shah, S. B. A. (1979): Comparative nutritive value of beef, mutton, and chicken meats and their extracts, Pakistan, J. of Sci. Res., 31, 1/2 , 22-25.
- Rodin, V. P. (1968): Meat quality of Suffolk×Tsigai crossbreds. A.B.A., 36(3):2639.
- Rouse, G. H., Topel, D. G., Veter, R. L., Rust, R. E. and Wickersham, T. W. (1970): Carcass composition of lambs at different stages of development. J. Anim., Sci., 31:846-855.
- Safa, K. Al-Amin (1976): Evaluation of carcass quality of sheep. M. Sci. Thesis, Fac. of Agric., Cairo Univ. Cairo Egypt.
- Safari, E. (1985): Comparison of feed-lot performance and carcass characteristics in rams, cryptorchids and wethers. Dep. Anim. Sci., College of Agric., Shivaz, Iran. World Rev. of Anim. Prod. 21(3) 3, 19-24 (En, 19 ref.).
- S. A. S. (1989): SAS User[^] F, [^] FS Guide:Basics SAS Ins, Cary. NS.
- Schreier, V. (1990): Results of a reciprocal cross-breeding trial involving German Mutton Merino and German Blackheaded Mutton sheep with special reference to fattening performance and carcass yield, Tierzucht., 44:4, 161-162.

- Seidzanov, E. M. (1964): Change with age and conditions of management in the meat of Kazakh Merino sheep. Abst. (A.B.A.), 33, 1297.
- Semenov, S. I., Sergeeva, L. G. and Kotvitskii, V. A. (1988): Carcass quality of lambs of different origin. Zootekhnika, A.B.A. No., 10, 31-32 (Ru).
- Sharaby, M. A. and Suleiman, I. O. (1988): Slaughter weights, carcass weight and dressing percentages of five breeds of sheep slaughtered in Saudia Arabia. Aust. J. Exp. Agric., 28(5):567-570.
- Sharafeldin, M. A. and K. E. Ghoneim (1966): Sheep production. 2nd ed. Cairo, in Arabic.
- Singh, V. K., Mathur, P. B. and Bohra, D. J. (1973): A note on carcass traits of coimbatore male lambs weaned at two ages. Ind. J. Anim. Sci., 43(8):783-785.
- Singh, B. B. and Singh, M. R. (1974): Relative importance of body measurements for body weight and wool yield in Bikaneri lambs. Ind. J. Anim. Health, 12(1):37-40.
- Singh, B., Singh, B. and Sangwan, M. L. (1988): Body conformation of sheep at weaning and six months of age. Ind. J. Anim. Prod. And management, 4(1):27-30.
- Soliman, H. S. (1971): Allowance of starch and protein for fattening lambs. M.Sc. Thesis, Fac. of Agric., Ain Shams Univ., Cairo.
- Srivastava, V. K. and Roy, A. (1969): Influence of season of birth and age on the carcass quality of Magra lambs. Ind. J. Anim. Sci., 39:294-306.

- Suliman, A. A. (1994): Improvement of some Ossimi productive and reproductive traits through crossing with Chios breed of sheep. M. Sci. Thesis, Fac. of Agric. El-Minia Univ. Egypt.
- Swidan, F. El-Shoboksy, A. S., Aboul-Naga, A. M. and Abbas, A. M. (1979): Fattening performance and carcass traits of early weaned Rahmani lambs. Egypt. J. Anim. Prod., 19, No.2, PP. 169-177.
- Tapiljskii, I. A. and Urickii, V. A. (1966): The meat production of crossbred sheep in Uzbekistan. A. B. A., 35(1):410.
- Thakur, A. K. B., Mahajan, J. M. and Singh, O. M. (1967): Study on body weight and conformation of Gadi and Romney March sheep. Ind. Vet. J., 44:589-596. (A.B.A., 36:1968).
- Todorovski, V., Risterski, K. and Popovski, K. (1973): Body measurements and weight in Awassi sheep reared in Yugoslavia. Veteriannviski Glasnik (1973), 27(6):427-434. (A.B.A., 41:5285).
- Uljjanov, A. N. (1968): Meat production of crossbred lambs in Kuban. A.B.A., 36(3):2647.
- Venkatesan, E. S. and Ranganathan (1972): The influence of age and sex on the carcass yield in Nellore and local Madras sheep. A.B.A., 41(9):3973.
- Vold, E. (1972): Investigations on meat and fat quality in two Norwegian sheep breeds. A. B. A., 41(1):168.
- Younis, A. A., El-Shobokshy, A. S., Hafz, M. and Danasoury, M. S. (1973): Effect of different levels of energy intake through alteration of roughage to concentrate ratio in rations for fattening lambs. Alex. J. Agric. Res., 21:229.

- Younis, A. A., Seoudy, A. M., Galal, E. S. E., Ghanam, Y. S. and Khishin, S. S. (1975): Effect of plane of nutrition on feed lot performance and carcass traits of desert sheep. *Trop. Agric.*, 52:233.
- Yusupov, S. Yu. And Basarov, S. (1991): Constitution in the selection of Karakul sheep. *Ovtsevodstvo*. No., 4, 21-23 (Ru).
- Zamarozov, A. V. (1963): The meat production of Romanov and crossbred lambs. *Abst. (A.B.A.)*, 33:374.
- Zamoryshev, A. V. and kuznetsova K. I. (1971): Meat production in Romanov sheep. *A. B. A.*, 41(8):3474.
- Zelenskii, G. G. (1971): Meat production of weathers obtained by commercial crossing with meat/short-wooled rams. *A. B. A.*, 40(4):4585.
- Zupp, W. and Grumbach, S. (1994): Testing of different sheep breeds for meat production. Commercial crossbreds have the best performance. Institut für Tierzucht, Landesforschung-sanstalt Mecklenburg-Vorpommern, Dummerstorf, Germany. *A. B. A. No. 9*, 69-70 (De)
- Zygoyiannis, D., Stamataris, K., Kouimtzis, S. and Doney, J. M. (1990): Carcass composition in lambs of Greek dairy breeds of sheep. *Anim. Prod.*, 50:261-269.