

- National Research Council, N. R. C. (1984).
Nutrient requirements of poultry, 8th ed. Nat.
Acad. of Sci., Washington, D.C.
- Omar, N. E. (1987).
Effect of different dietary protein and energy
levels in a three-stage diet system on general
performance of broiler chicks.
Ph. D. Thesis, Fac. Agric., Cairo Univ.
- Payne, I. F. (1930).
Malformed leg bones in young chickens. Science,
71: 664.
- Perlman, R. L., Telser, A. and Dormann, A. (1964).
Studies on the role of manganese in bone
formation.
J. Biol. chem., 237: 3623.
- Plavnik, I., Bornstein, S. and Hurwitz, S. (1981).
Evaluation of methanol-grown bacteria and
hydrocarbon-grown yeast as sources of protein for
poultry: Studies with young chicks.
Br. Poultry Sci., 22: 123.
- Rehnberg, G. L., Hein, J. F., Carter, S. D., Linko, R. S.,
and Laskey, J. W. (1981).
Chronic ingestion of manganese oxide by young
rats: tissue accumulation, distribution, and
depletion.
J. Toxicol. Environ. Health, 7: 263 - 272.
- Renner, R., and Hill, F. W. (1959).
Metabolizable energy value of fats and fatty acids
for chickens.
Proc. Cornell Nutrition conference, page 95-100.
- Robinson, H. C., Telser, A. and Dormann, A. (1966).
Studies on the role of manganese in bone
formation.
Proc. Natl. Acad. Sci. U.S., 56: 1859.

Sadzhad, K. H. (1983).

Productivity and metabolism of substances in hens in relation to dietary manganese level. Nutr. Abst. & Rev., 55 (4) : 2055 (Series B, 2055) , 1985.

Salmon, W. D., Jr. and Daughaday, W.H. (1957).

A hormonally controlled serum factor which stimulated sulfate incorporation by catilage in vitro.

J. Lab. Clin. Med., 49: 825 -836.

Saly, J., Fried, K., Jantosovic, J., Kusev, J. and Benhatchi, M. (1985).

The effect of manganese on egg shell quality.

Folia Veterinaria, 29 : 91 - 99

(Nutr. Abst. & Rev. , 55 (4) : 777, 1985).

Schaible, P. J., Bandemer, S. L. and Davidson, J. A. (1938).

The manganese content of feedstuffs and its relation to poultry nutrition.

Michigan state college Agric. Exp. Sta. Bull.No. 159.

Seth, P. C. C. and Clandinin, D. R. (1973).

Effect of including rapseed meal in the ration of broiler-type chickens on the incidence of perosis and the ineffectiveness of supplemental manganese.

Poultry Sci., 52 : 1158 - 1160.

Settle, E. A., Mraz, F. R., Douglas, C. R. and Bletner, J. K. (1969).

Effect of diet and manganese level on growth, perosis and Mn uptake in chicks.

J.Nutrition, 97: 141 - 146.

Shannon, D. W. F., and McNab, J. M. (1972).

The effect of different dietary levels of n-paraffin-grown yeast on the growth and food intake of broiler chicks. Br. Poultry Sci., 13 : 267.

- Silbert, J. E. (1964).
Studies on the role of manganese in bone formation.
J. Biol. Chem., 239: 1310.
- Southern, L.L., and Baker, D.H. (1983 a).
Eimeria acervulina infection in chicks fed deficient or excess levels of manganese.
J. Nutr. , 113 : 172 - 177.
- Southern, L. L., and Baker , D. H. (1983 b).
Excess manganese ingestion in the chick.
Poultry Sci., 62: 642 - 646.
- Steel, R. G.D., and Torrie, J.H. (1980).
"Principles and Procedures of Statistics;" A Biometrical Approach. 2nd ed, Mc Graw-Hill Book Co., New York , N. Y.
- Stephens, J. F., Bletner, J. K. and Goff, O. E. (1961).
Supplementation and processing methods of feather meal used in broiler diets.
Tenn, Farm Home Sci., Prog. Rep., 39:2.
- Sturkie, P. D. (1954).
" Avian Physiology".
Constock Publishing Associates, Cornell Univ. Press., Ithaca, New York, Page 224.
- Succi, G., Pialorsi, S., Fiore, Di., L. and Cardini, G. (1980).
The use of methanol - grown yeast LI-70 in feeds for broilers.
Poultry Sci., 59: 1471.
- Suso, F. A., and Edwards, Jr. , H.M. (1968).
A study of techniques for measuring manganese absorption and biological half-life in the chicken.
Poultry Sci., 47: 991 -999.

- Telser, A. Robinson, H. C. and Doremann, A. (1965).
Studies on the role of manganese in bone
formation.
Proc. Nat. Acad. Sci. U. S, 54: 912.
- Terpstra, K. and De Hart, N. (1974).
The estimation of urinary nitrogen and faecal
nitrogen in poultry excreta.
Z.Tierphysiol., Tierernahrg. U.
Futtermittelkde, 32:306-320.
- Thomas, K. W. and Lowther, D. A. (1976).
Manganese levels and the morphology of the
epiphyseal plate in broilers with slipped tendons.
Poultry Sci., 55 : 1962- 1968,
- Titus, H. W. (1932).
Perosis, or deforming leg weakness, in the
chicken.
Poultry Sci., 11: 117-125.
- Turk, D. E. (1974).
Intestinal parasitism and nutrient absorption.
Fed. Proc., 33: 106-111.
- Turk, D. E. (1978).
The effects of coccidiosis on intestinal function
and gut microflora. Pages 227-267 in Avian
coccidiosis. P.L. Long, K. N. Boorman, and B. M.
Freeman, eds. Br. Poultry Sci. Ltd., Edinburgh,
Scotland.
- Turk, D.E., Gunji, D. S. and Molitoris, P. (1982).
Coccidial infections and manganese absorption.
Poultry Sci., 61: 2430-2434.
- Underwood, E. J. (1977).
" Trace elements in human and animal nutrition."
4th ed. Academic Press, New York, N. Y.