




LITERATURE CITED

LITERATURE CITED

- Aggour, R. A., L.A. Badr, M. korayem and M. M. Mansour (1998).** Genetic studies on root phenols content as a defense mechanism against root knot nematode in tomato. Egypt. J. Plant Breed. 2, 111–133.
- Ahuja, S. and M.C. Muchopadhyaya (1985).** Variability in the occurrence of root-knot nematode, *Meloidogyne incognita* infection in various vegetable crops. Bulletin of Entomology. 1985, 26: 1, 17-24.
- Bajaj, K.L. and R. Mahajan (1977).** Phenolic compound in tomato susceptible and resistant to *Meloidogyne incognita* (Kofoid & White) Chitwood. Nemat. Mediterr. 5:329-333 .
- Bajaj K.L. , Y.K. Arona and R. Mahajan (1983).** Biochemical differences in tomato cultivars resistant and susceptible to *Meloidogyne incognita*. Helm. Abstr. 52:1541.
- Bajaj, K.L., P. Singh and K. Mahajan (1986).** Changes induced by *Meloidogyne incognita* in superoxide dismutase and peroxidase activity in tomato roots : Helm. Abstr. 55:96 .

Bhatt, G.M. (1971). Heterotic performance and combining ability in a diallel cross among spring wheats (*T. aestivum* L.). *Austr.J. of Agric. Res.* 22:359-369.

Bingefors, S. (1982). Nature of inherited nematode resistance in plants . pp. 188-219 in K.F.Harris and K. Maramorsch. eds. *Pathogeus, Vectors and plant desease*. Orlando. FL : Academic press.

Boerma, H.R. and R.S.Hussey (1992). Breeding plants for resistance to nematode . *J. Nematol.* 24(2) : 242-252.

Canto-Saenz, M. (1983). The nature of resistance to *Meloidogyne incognita* (Kofoid & While) Chitwood, pp.160-165. In *proc. Third Res. And Plann. Conf. on Root Knot Nematode, Meloidogyne spp.*, March22-26,1982,ed. C.C. Carter, International Meloidogyne project, Lima, Peru. 233 pp.

Castagnone, S.P., M. Bongioranni and A. Dalmasso (1992). Differential expression of root knot nematode resistance genes in tomato and pepper : evidence with *Meloidogyne incognita* virulent and avirulent near-isogenic line ages. *Annals of Applied Biology* 120 (3) : 487-492.

Cho. H.J., S.C. Han. and D.R. Choi. (1986). Screening groundnut, sweet pepper, cucumber and tomato varieties for resistance to root-knot nematode,

Meloidogyne hapla. Research Reports, Rural Development Administration, Korea Republic, Plant Environment, Mycology and Farm Product Utilization. 1986, 28 : 1, 94-97.

Dalmasso, A, M.C. Cardin, E. Pochard and M.C. Daunay (1985). Pathogenicity of *Meloidogyne* nematodes and the genetic resistance in some solanaceous vegetables. *Compte-Rendus-des-Seances-de-l'Academie-d'Agriculture-de-France*. 1985, 71: 7, 771-779; Numero Special : Creation de varietes resistantes aux nematodes des cultures: interet, possibilites et limites.

Das, P., P.N. Phukang and D.K. Sarmah (1988). Screening of some vegetables varieties against root knot nematode *Meloidogyne SPP*. Research, Assam, agric. Univ., 9 (1-2) ; 56 -57.

Di-Vito, M. (1986). Population densities of *Meloidogyne incognita* and growth of susceptible and resistant pepper plants. *Nemat. Mediterr.* 1986, 14: 2, 217-221.

Di-Vito, M., Saccardo and N. Sasanelli (1989). Reaction of lines of *Capsicum SPP.* to root knot nematodes (*Meloidogyne SPP.*). Proceeding of EUCARPIA VIIth meeting on genetics and breeding on capsicum and eggplant, Krogujevac, Yugoslavia. pp. 171-175.

- Di-Vito, M., G. Zaccheo and J.M. Poulos (1996).** Reaction of Asian germplasm of pepper (*Capsicum annuum* L.) to Italian populations of root knot nematodes (*Meloidogyne* spp.). *Capsicum and Eggplant Newsletter*. 15 ; 63-66.
- Di-Vito, M., F. Saccardo and G. Zaccheo (1991).** Response of lines of *Capsicum* spp. to Italian population of four species of *Meloidogyne*. *Nematologia Mediterranea*. 19 (1) : 43 – 46; 8 ref
- Di-Vito, M.P, F. Saccardo, A. Errico and G. Zacchea (1992).** Genetic resistance to root knot nematodes (*Meloidogyne* spp.) in *Capsicum chacoense*, *C. Chinese* and *C. frutescence*. *Capsicum– Newsletter*, Special Issue : 205–209 ; EUCARPIA VIIIth meeting on genetics and breeding on capsicum and eggplant, Rome, Italy, 7-10 September 1992 ; 6 ref.
- Di-Vito, M., F. Saccardo and G. Zaccheo (1993).** Response of new lines of pepper to *Meloidogyne incognita*, *M. javanica*, *M. arenaria* and *M. hapla*. *Afro-Asian J. Nematol.* 3: 2, 135-138.
- Di-Vito, M.D., F. Saccardo, A. Errico, G. Zaccheo, and F. Catlano (1993).** Genetics of resistance to root knot nematodes (*Meloidogyne* spp.) in *Capsicum chacoense*, *C. Chinese* and *C. frutescence*. *Genetics and Breeding* . 47 (1) : 23-26.

- Dukes, P.D.S. and L. Fery (1997).** Charleston Hot, a southern root knot nematode resistance, Yellow fruited Cayenne pepper with a compact plant Habit. HortScience. 32 (5): 947-948.
- El-Sherif, M.A., M.H. Belal and T. Badra (1980).** Activity of polyphenol oxidase in tomato healthy and infected by *Rotylenchus reniformis* and *Meloidogyne incognita*. Ans. Schadl. Pflanz. Umwelt. 53:10-13.
- Fassuliotis, G. (1987).** Genetic basis of plant resistance to nematodes. pp. 364-371 in J.A. Veech and D.W. Dickon, eds. Vistas on nematology. Society of nematologists.
- Fery, L. and D. Dukes (1996).** The inheritance of resistance to southern root knot nematode in "Caroline Hot" Cayenne pepper. J.Amer. Soc. Hort .Sci . 121(6) : 1024-1027.
- Fery, R.L. and J.A. Thies (1997).** Evaluation of *Capsicum chinense* Jacq. cultigens for resistance to southern root knot nematode. HortScience, 32(5) : 923-926.
- Fery, R.L. and J.A. Thies (1998 a).** PA -353, PA -398 and PA-426: Southern root knot nematode resistance *Capsicum chinense* Jacq. Germplasm lines. HortScience. 33 (4) : 760-761.

- Fery, R.L and J.A. Thies (1998 b).** Genetic analysis of resistance to the southern root knot nematode in *Capsicum chinense* Jacq. J.Amer. Soc. Hort. Sci. 123(6) : 1008-1011.
- Fery, R.L., D.S. Philip and A. Judy (1998).** Carolina Wonder and Charleston Bell : southern root knot nematode resistant Bell pepper : HortScience. 33(5) : 900-902.
- Ganguly, S. and D.R. Dasgupta (1982).** Characterization of polyphenol oxidase from susceptible and resistant varieties of tomato inoculated with the root knot nematode, *Meloidogyne incognita* . Ind. J. Nematol. 12(2) :320-390.
- Gardner, C. O. (1963).** Estimates of genetic parameters in crosses fertilizing plant and their implication in plant breeding P. 225-252. In : W.D.Hanson and H.F. Robinson (eds.) statistical genetics and plant breeding. Nat. A cad. Of Sci. Nat. Res. Counc. Pub. pp 982.
- Giebel, J. (1974).** Biochemical mechanisms of plant resistance to nematodes. J. Nematol. 6:175-184.
- Gomez, K.A. and A.A. Gomez (1984).** Statistical procedures for agricultural research and. John Wiely and Sons. New York.

- Griffing, B. (1956).** Concept of general and specific combining ability in relation to diallel crossing systems. *Austral. J. Biol. Sci* 9:463-93.
- Hassan, A.A. (1993).** Breeding of vegetable crops. In Arabic. El-Dar El-Arabia Lil Nashr Wa Tawzee. Cairo, Egypt. pp. 799.
- Hassan, H.M. (1990).** Studies on the nature of resistance to root-knot nematode in tomato. Ph. D. Thesis , Fac. Agric., Alex. Univ., Egypt.
- Hassan, N. and S. K. Saxena (1977).** Effect of inoculating tomato with root-knot nematode, *Meloidogyne incognita* on the phenolic content and polyphenol oxidase activity. Procedures of Symp. On : physiology of the host-pathogen interaction . Nov. 25-26, 1977, Madras. India.
- Hayman, B.I. (1954).** Theory and analysis of Diallel crosses. *Genetics* 39 : 789-809.
- Hendy, H, A. Dalmasso and M.C. Cardin (1985).** Differences in resistant *Capsicum annuum* attacked by different *Meloidogyne* species. *Nematologica*. 31: 1, 72-78.
- Houssny, H.H. and B.A. Oteifa (1956).** Preliminary field tests for evaluating some tomato varieties for resistance to

root-knot nematodes (*Meloidogyne* species.) PL. Dis. Rept . 40 : 974-976.

Jinks, J.L. (1954). The analysis of continuous variation in a diallel cross of *Nicotiana rustica* varieties . Genetics 39: 767-788.

Khan, B, A.A. Khan and M.R. Khan (2002). Response of some cultivars of pepper, eggplant and tomato to races of root-knot nematodes, *Meloidogyne incognita* and *M. javanica*. Ind. J. Nematol. 32: 2, 147-152.

Khelu, A.Z., V.G. Zaets and A.A. Shesteporov (1989). Tests on resistance to root-knot nematodes in tomato, cucumber and pepper varieties grown under cover in central Lebanon. Byulleten'- Vsesoyuznogo- Instituta- Gel' mintologii-im.-K.-I.-Skryabina. 1989, No. 50, 85-89.

Lamberti, F. (1979) .Economic importance of *Meloidogyne* spp. In suberropical and Mediterranean climates, P341 – 357. In : F. Lamberti and C.F. Taylor (eds) . Root-knot nematodes (*Meloidogyne* species) : Systematics, biology, and control. Academic, New-York .

Marwoto, B . (1989) . Response of various pepper cultivars to root knot nematodes (*Meloidogyne* spp.): Buletin Penelitian Hortikultura. 17: 4, 15-19. AGRICOLA Data Base INTERNET.

- Masood, A. and S.I. Hussain (1979).** Phenolic and ortho-dihydroxy phenolic changes and their role in the resistance and susceptibility of three tomato varieties to *Meloidogyne incognita*. Helm. Abstr. 48:1119.
- Mather, K. and J.L.Jinks (1971).** Biometrical genetics . Cornell University Press, Ithaco,NY.
- McClure, M.A.; T.H. Kruk and I. Misaghi (1973).** A method for obtaining quantities of clean *Meloidogyne* eggs. J. Nematol. 5:230.
- Mekete, T, W. Mandefro and N. Greco (2003).** Relationship between initial population densities of *Meloidogyne javanica* and damage to pepper and tomato in Ethiopia. Nematologia Mediterranea.31: 2, 169-171.
- Narayana, Y.D. (1979).** Studies on the nature of resistance in tomato (Var. NTDR-1) to root-knot nematodes. Helm. Abstr 48:1578.
- Narayana, Y.D. and D.D.R. Reddy (1980).** The role of nitrogen, ammonia acid and phenols in resistance of tomato to root-knot nematodes Nemat. Mediterr. 8:51-57.
- Ogbuji, R.O. and M.O. Okafor (1984).** Comparative resistance of nine pepper (*Capsicum annuum* L.) cultivars to three root-knot nematode (*Meloidogyne*) species and their related use in traditional cropping

systems. Beitrage-zur-Tropischen-Landwirtschaft-und-
eterinarmedizin. 22: 2, 167-170. AGRICOLA Data
Base INTERNET.

Okophyi, N.S. (1983). Polyphenol oxidase isoenzymes in roots of
tomatoes resistance to *Meloidogyne incognita*. Helm.
Abstr. 52:669.

Okophyi, N.S. and A.V. Sadykin (1978). The resistance of
tomatoes to the gall nematodes. Pl. Breed. Abstr. 48
(7) : 7128.

Okophyi, N.S., N.N. Zaginailo and A.V. Sadykin (1987). A
method of selection nematode resistant tomato
varieties and lines. Pl. Breed. Abstr. 57:3324.

Oostenbrink, M. (1966). Major characteristics of the relation
between nematodes and plants. Meded. Landbouw-
hogesch. Wageningen, 46 pp.

Peixoto, J.R., W.R. Moluf, and V.P. Campos (1995). Evaluation
of pepper genotypes for resistance to *Meloidogyne*
incognita (race 2) and *Meloidogyne Javanica* . Horti
Bxasileira . 13(2) 154-158.

Peter, K.V., R.W. Goth and R.E. Webb (1984): Indian hot
peppers as new sources of resistance to bacterial wilt,
Phytophthora root rot, and root knot nematode. :
HortScience. 1984, 19: 2, 277-278.

- Potenza, C.L., S.H. Thomas, E.A. Higgins and C. Sengupta (1996).** Early root response to *Meloidogyne incognita* in resistant and susceptible Alfalf cultivars. J. Nematol. 28 (1) : 175-181.
- Ramana, K.V. and C. Mohandas (1986).** Reaction of black pepper germplasm to root-knot nematode, *Meloidogyne incognita*. Ind .J. Nematol. 16: 1, 138-139.
- Riggs, R.D. and N.N. Winstead (1959).** Studies on resistance in tomato to root-knot nematode and occurrence of Pathogenic Biotypes Phytopathology. 49: 716-784.
- Roberts, P.A. (1990).** Resistance to nematodes: Definitions, concepts, and consequences. pp. 1-15 in J.L. Starr, ed. Methods for evaluating plant species for resistance to plant-parasitic nematodes. Society of Nematologists.
- Routaray, B.N., H. Sahoo and S.N. Das (1988).** Evaluation of some chilli cultivars against *Meloidogyne incognita* and *Rotylenchulus reniformis*. J. Ind. Bot. Soc . 1988, 67: 1-2, 220-221.
- Sadykin, A.V. and N.S. Okophyi (1985).** Rapid method of evaluating the nematode resistance of tomato. Pl. Abstr. 55 (12)10029.

- Santos, H.S., S.R.S. Wilcken and R. Goto (2002).** Reproduction of *Meloidogyne incognita* race 2 on different sweet-peppers (*Capsicum annuum* L.). *Nematologia-Brasileira*. 26: 2, 209-211.
- Sasser, J.N. (1977).** World wide dissemination and importance of the root – knot nematodes *Meloidogyne* spp. *J. Nematol.* 9 : 26-29
- Sidhu, G.S. and J.M. Webster (1981).** The genetics of plant-nematode parasitic svstems. *Botanical Review* 47:387-419.
- Simmonds, N.W. (1991).** Genetics of horizontal resistance to disease of crops . *Biological review* . 66 : 189-241.
- Snell, F.D. and C.T. Snell (1953).** Clorimetric methods of Analysis including some turbidimetric methods. D.Van. Nostrand compony Inc. Tornto-New York. London Vol III orgnic-1, pp 606.
- Souza, F., W.R. Maluf, L.A.A. Gomes and V.P. Campos (2002).** Inheritance of resistance to *Meloidogyne incognita* race 2 in the hot pepper cultivar Carolina Cayenne (*Capsicum annuum* L.). *Genetics and Molecular Research*. 1: 3, 271-279.
- Stegemann, H., A.E.T. Shehata and M. Hamza (1985).** Broad bean proteins (*Vicia Faba* L.): Electrophoretic studies

on seeds of some German and Egyptian cultivars.
Zeitschrift für Acker und Pflanzenbau 149 : 447-453.

Taylor, A.L. and J.N. Sasser (1978). Biology, identification and control of root knot nematodes (*Meloidogyne spp.*)
Coob pub. Dep. Plant Pathol., North Carolina State Univ., and U.S. Agency Int. Dev. Raleigh, N.C. III PP.

Thies, J.A., J.D. Mueller and R.L. Fery (1997). Effectiveness of resistance to southern root knot nematode in Carolina Cayenne pepper in greenhouse, microplot, and field tests. J.Amer. Soc. Hort. Sci. 122 (2) : 200-204.

Thies, J.A., J.D. Mueller and R.L. Fery (1998). Use of a resistant pepper as a rotational crop to manage to root knot nematode. HortScience. 33(4): 716–718.

Thies, J.A. and R.L. Fery (2000) .Characterization of resistance conferred by the N gene to *Meloidogyne arenaria* races 1 and 2, *M. hapla*, and *M. javanica* in two sets of isogenic lines of *Capsicum annuum* L. J. Amer. Soc. Hort. Sci. 125: 1, 71-75.

Thies, A. and R.L.Fery (2003). Response of bell pepper cultivars Near–isogenic for the N Gene to *Meloidogyne incognita* in field Trails. HortScience 38(7) : 1394-1396.

- Thies, J.A., J.D. Mueller and R.L. Fery, M. Gilbert and V. Joseph (2003). Response of bell pepper cultivars near-isogenic fore the N Gene to Meloidogyne incognita in field Trials. HortScience 38 (7): 1394-1396.*
- Thomas, S.H. and M. Cardenas (1985). Relationships between preseason numbers of Meloidogyne incognita and yield losses in chile pepper cultivars. Phytopathology. 75: 11, 1304.*
- Wiliski, A., J. Giebel and J. Glowinkowska (1977). Phenolic compounds in roots of tomatoes susceptible and resistant to Meloidogyne spp. Proce Naukowe Instytutu Ochrony Roslin, 13:155-162.*
- Yadav, B.D. and B.N. Mathur, (1993). Reaction of chilli genotypes to root-knot nematodes Meloidogyne incognita. Indian Journal of Mycology, and plant pathology. 23 (1) : 92-93.*
- Zamora, E., and P.W. Bosland (1994). Carolina Cayenne as a source of resistance to Meloidogyne incognita races 1, 2, 3 and 4. HortSciene 29(10) : 1184-1185.*