




REFERENCES

6. REFERENCES

- Abdel-Fattah, A. and El-Sayed, M. (1999):** Alternative dietary protein sources for farmed tilapia, *Oreochromis spp.* Aquaculture, **179**:149-168.
- Abdel-Ghany, A. E. (1995):** Effect of winter feeding on the rate of growth, food conversion and survival of Nile tilapia (*Oreochromis niloticus* L.) and common carp (*Cyprinus carpio* L.) in Egypt. Egypt. J. Agric. Res., **73** (1): 291-306.
- Abdel-Hakim, N. F., and Moustafa, E. T. (2000):** Performance of Nile tilapia (*Oreochromis niloticus*) raised in cages as affected with stocking density and dietary protein level. Egypt. J. Aquat. Biol and Fish, Vol. 4, No. **2**:95-116.
- Abo-State, H. A. M. (1997):** Using different levels and sources of lipids in fish nutrition. M. Sc. Thesis, Fac. of Agric, Cairo Univ.
- Agouz, H. M. and Tonsy, H. D. (2003):** Evaluation of whole cryfish meal (*Procambrus clarkii*) as partial or complete replacement of fishmeal protein in polyculture commercial diets. Egyptian J. Nutrition and Feeds, **6** (Special Issue):315-330.
- Akiyama, D. M. (1988):** Soybean utilization in fish feeds. Presentation at the Korean Feed Association Conference, Seoul, Korea.

- Akiyama, D. M., Dominy, W. G. and Lawrence, A. I., (1992):** Penaeid shrimp nutrition. In: *Marine Shrimp Culture Principles and Practices* (ed. by E.W. Past and L.J. Lester). PP. 535-568. Elsevier Science Publishers, BV, Amsterdam, the Netherlands.
- Aksoy, M., Lim, C.E. and Li, M.H. (2006):** Interaction between Dietary Levels of Vitamin C and E on Growth and Immune Responses in Channel Catfish *Ictalurus*. Aquaculture America Conference. Las Vegas, NV.
- Al-Azab, A. A. (2005):** Effect of partial and complete replacement of fish meal with locally produced shrimp meal on growth performance of Nile tilapia (*Oreochromis niloticus*). Egyptian J. Nutrition and Feeds, 8(Special Issue):1145-1156.
- Alford, B. B., Liepa, G. U. and Vanbeber, A. D. (1996):** Cottonseed protein: what does the future hold?. Plant Foods Human Nutrition, **49**:1-11
- Anderson, C. E. and Smitherman, R. O. (1978):** Production of normal male and androgen sex-reversal *Tilapia aurea* and *T. nilotica* fed a commercial cat fish diet in ponds, pp. 34-39. In R. O. Smitherman, W. L. Shelton and J. H. Grover (eds.). Culture of Exotic Fish Symposium Proceeding. Fish Culture Section. American Fisheries Society, Auburn, Alabama.
- Anderson, J. S. and Sunderland, R. (2002).** Effect of extruder moisture and dryer processing temperature on vitamin C and E and astaxanthin stability. Aquaculture, **207**: 137-149.

- Anderson, J.; Jackson, A. J.; Matty, A. J. and Copper, B. S. (1984):** Effect of dietary carbohydrate and fiber on the tilapia, *Oreochromis niloticus*. *Aquaculture*, **37**: 303-314.
- Anderws, J. W. and Page, J. W. (1974):** Growth factors in fishmeal component of catfish diet. *J. Nutr.*, **104**: 1091-1096.
- Association of Official Analytical Chemists, AOAC (1990):** Official Methods of Analysis, 15th edn. 684 pp. Association of Official Analytical Chemists, Arlington, VA, USA.
- Barraza, M.I., Coppock, C.E., Brooks, K.N., Wilks, D.L., Saunders, R.G. and Latimer, Jr, G.W. (1991):** Iron sulfate and feed pelleting to detoxify free gossypol in cottonseed diets for dairy cattle. *J Dairy Sci.*, **74**: 3457-3467.
- Barros, M.M., Lim, C. and Klesius, P.H. (2002):** Effect of soybean meal replacement by cottonseed meal and iron supplementation on growth immune response and resistance of channel catfish (*Ictalurus punctatus*) to *Edwardsiella ictaluri* challenge. *Aquaculture*, **207**: 263-2798.
- Blom, J. H., Lee, K. J., Rinchard, J., Dabrowski, K. and Ottobre, J. (2001):** Reproductive efficiency and maternal offspring transfer of gossypol in rainbow trout (*Oncorhynchus mykiss*) fed diets containing cottonseed meal. *J. Anim. Sci.*, **79**: 1533-1539.

- Bowen, S. H. (1982):** Feeding, digestion and growth-qualitative considerations. *In: R. S. V. Pullin and R. H. Lowe-McConnell (editors). The Biology and Culture of Tilapias. International Center of Living Aquatic Resources Management, ICLARM Conference Proceeding 7, Manila Philippines. Pages 141-156*
- Bowman, D. (1977):** Comparison enter *Tilapia aurea* steindachnery *Tilapia massambicus* peters en estanques de El-Salvador. *FAO Fish. Rep. 159 (1): 78-90.* (In Spanish with English summary).
- Boyd, D. (1979):** Water Quality in Warmwater Fish Ponds. Auburn University Agricultural Experiment Station, Auburn, AL.
- Braham, J.E. and Bressani, R. (1975).** Effect of different levels of gossypol on transaminase activity, on nonessential to essential amino acid ratio, and on iron and nitrogen retention in rats. *J. Nutrition, 105:348-355.*
- Braham, J.E., Jarquin, R., Elias, L.G., Gonzalez, M. and Bressani, R. (1967).** Effect of calcium and gossypol on the performance of swine and on certain enzymes and other blood constituents. *J. Nutrition, 91:47-54.*
- Bratt, J. R. and Groves, T. D. (1979):** Bioenergetics and growth, edited by Hoar, W. C., Randall, D. J. and Brett, J. R. *Nutrition, in Fish Physiology. Academic Press, New York, San Francisco and London. Vol. VIII; 179-352.*

- Brocas, C., R. M. Rivera, F. F. Poula-Lopez, L. R. McDowell, M., C. Calhoun, C. R. Staples, N. S. Wilkinson, A. J. Boning, P. J. Chenoweth, and P. J. Hansen. (1997): Deleterious actions of gossypol on bovine. Oocytes, and embryos. Biol. Reprod., **57**:901-907.
- Cahu, C. and Fakhfakch, M. (1990): Effect of dietary vitamin E on reproduction of penaeid shrimp. I. Zootechnical results. Aquaculture 90. Halifax. Canada.
- Cahu, C., Villette, M., Quazuguel, P. and Guillaume, J. (1991): The effect of n-3 highly unsaturated fatty acid and vitamin E supplementation in broodstock feed on reproduction of *penaeus indicus*. In: Fish Nutrition in Practice. Biarritz (France). Vol. 61.PP.589-598. INRA. Paris.France.
- Carleton, M.A., Drury, R.A.B., Wallinton, C.H. (1967): Carletons histological technique. 4th Ed. Oxford Univ. Press, New York, Toronto.
- Castelli, M.E., Castillo, A.R., Gallardo, M.R. and Quano, O.R. (1993): Dairy cows on lucerne pasture supplemented with cottonseed: evaluation of some blood components. *Rev. Med. Vet., (BuenosAires)* **74**: 236-242.
- Chau, B. S. and Shiau, S.Y. (1996): Optimal lipid level for growth of juvenile hybrid tilapia, *Oreochromis niloticus*, *Oreochromis aureus*. Aquaculture, **143**: 185-195.

- Cheng Z.J. and Hardy, R.W. (2002):** Apparent digestibility coefficients and nutritional value of cottonseed meal for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, **212**:361-372.
- Chervinski, J. (1982):** Environmental Physiology of Tilapias. *In*: R.S.V. Pullin and R.H. Lowe Mc-Connell eds. The Biology and Culture of Tilapias ICLARM Conference Proceedings, 7. International Center for Living Aquatic Resources Management (ICLARM), Manila, Philippines, pp. 119-128.
- Cho, C. Y., Cowey, C. B. and Watanabe, T. (1985):** Fish nutrition in Asia: methodological approaches to research and development. Ottawa, IDRC.
- Coche, A. G., (1982):** Cage culture of tilapias. *In*: R. S. V. Pullin and R. H. Lowe-Mc Connell (editors). The Biology and Culture of Tilapias. International Center of Living Aquatic Resources Management, (ICLARM), Conference Proceedings 7, Manila Philippines pp. 205-246.
- Coppock, C.E., West, J.W., Moya, J.R., Have, D.H., LaBore. J.M., Thompson, K.G., Rowe, Jr, L.D. and Gates, C.E. (1985):** Effect of amount of whole cottonseed on intake, digestibility, and physiological responses of dairy cows. *J Dairy Sci.*, **68**:2248-2258.
- Cowey, C. B. (1978):** Protein and amino acid requirements of finfish. ELFAC Symp. Fin fish. Nutr. And Feed Technol., Hamburg. FLFC/78/symp. R 16.

- Dabrowski, K., Rinchar, J., Lee, K. J., Blom, J.H., Ciereszko, A. and Ottobre, J. (2000 a): Effects of diets containing gossypol on reproductive capacity of rainbow trout (*Oncorhynchus mykiss*). Biol. Reprod., 62:227–234.
- Dabrowski, K., Rinchar, J., Ciereszko, A., Lee, K. J. and Czesny, S. (2000 b): Effects of gossypol containing diets on the viability of male and female gametes in rainbow trout (*Oncorhynchus mykiss*). 130th Annual Meeting
- Dabrowski, K., Lee, K. J., Rinchar, J., Ciereszko, A., Blom, J. H. and Ottobre, J. (2001): Gossypol isomers bind specifically to blood plasma proteins and spermatozoa of rainbow trout fed diets containing cottonseed meal. Biochim. Biophys. Acta, Gen. Subj., 1525: 37–42.
- Dato-Cajegas, C. R. and Yakupitiyage, A. (1996): The need for dietary mineral supplementation for Nile tilapia *Oreochromis niloticus*, cultured in semi-intensive system. Aquaculture, 144: 227-237.
- De-LaHiguera, M., Garcia-Gallego, M., Sauz, A., Cardenete, G., Suarez, M. D. and Moyano, F. J. (1988): Evaluation of lupin seed meal as an alternative protein source in feeding of rainbow trout. Aquaculture, 91: 37-50.
- De-Silva, S. S., Gunasekera, R. M. and Shim, K. F. (1991): Interactions of varying dietary protein and lipid levels in young red tilapia evidence of protein sparing. Aquaculture, 95: 305-318.

- Dorsa, W. J., Robinette, H. R., Robinson, E. H. and Poe, W.E. (1982):** Effects of dietary cottonseed meal and gossypol on growth of young channel catfish. Trans. Am. Fish. Soc., **111**: 651- 655.
- Duncan, D. B. (1955):** Multiple range and multiple F-test. Biometrics, **11**:1-24.
- Dupree, H. K. and Huner, J. V. (1984):** Nutrition, feeds and feeding practices. P. 141-157. in Duoree, H.K. and J.V. Huner, eds. Third report to the fish farmers. USA Fish and Wildlife Service Washington, D.C.
- El-Banna, R.A. (1991).** Studies on some nutrient requirements of tilapia and its effect on performance. Ph.D.Thesis, Fac.of Vet.Med., Cairo Univ.
- El -Eraky, W., Saleh, G. and Gropp, J. M. (1995):** A short note on the vitamin E requirement of *Tilapia nilotica* (*Oreochromis niloticus*); effect on health and growth. J. Appl. Ichthyol., **11**: 375-377.
- El-Husseiny, O. M., Arafa, S. A., Shehata, T. M. and Abdel-Samee, A. M. (1993):** Effect of dietary energy sources on Tilapia performance. 4th Symp. Anim. Poult. Fish Nut, El Fayoum, 63-93.
- El-Saidy D. M. (1999):** Evaluation of cottonseed meal as partial and complete replacement of fish meal in practical diets of Nile tilapia (*Oreochromis niloticus*) fingerlings. Egyptian Journal of Aquatic Biology and Fisheries, **3**: 441-457.

- El-Saidy, D. M. and Gaber, M. M. (2004):** Use of cottonseed meal supplemented with iron for detoxification of gossypol as a total replacement of fish meal in Nile tilapia *Oreochromis niloticus* (L.) diets. *Aquaculture Research*, **35**: 859-865.
- El-Sayed, A. M. (1990):** Long-term evaluation of cottonseed meal as a protein source for Nile tilapia *Oreochromis niloticus* (Linn.). *Aquaculture*, **87**:315-320.
- El-Sayed, A. M. and Teshima, S. (1992):** Protein and energy requirements of Nile Tilapia *Oreochromus niloticus* fry. *Aquaculture*, **103**: 55-63.
- Ensminger, M. E., Oldfield, J.E. and Heinemann, W. W. (1990):** Feeds and nutrition. The Ensminger Publ. Co., California, 1169-1197.
- Fagbenro, O. A. and Jauncey, K. (1993):** Chemical and nutritional quality of stored fermented fish silage. *Biores. Technol.*, **46**:207-211.
- Fagbenro, O. A. and Jauncey, K. (1994):** Chemical and nutritional quality of dried fermented fish silage and their nutritive value for tilapia (*Oreochromis niloticus*). *Animal Feed Sci. Technol.*, **45**:167-176.
- Forster, L. A. and Cahloun, M. C. (1995):** Nutrient values for cottonseed product deserve new look. *Feedstuffs*, **67**: 1-5.
- Fowler, L. G. (1980):** Substitution of soybean and cottonseed products for fish meal in diets fed to chinook and coho salmon. *Prog. Fish-Cult.*, **42**: 87-91.

- Frigg, M., Prabucki, A.L., Rubell, E.U., (1990):** Effects of dietary vitamin E levels on oxidative stability of trout fillets. *Aquaculture*, **84**: 145-158.
- Garcia-Abiado, M. A. Mbahinzireki, G., Rinchar, J., Lee, K. J. and Dabrowski, K. (2004):** Effect of diets containing gossypol on blood parameters and spleen structure in tilapia, *Oreochromis sp.*, reared in a recirculating system. *Journal of Fish Diseases*, **27**: 359-368.
- Garcia-Abiado, M. A., Mbahinzireki, G., Rinchar, J., Lee, K. J. and Dabrowski, K. (2005):** Effect of diets containing gossypol on blood parameters and spleen structure in tilapia, *Oreochromis sp.*, reared in recirculating system. *Journal of Fish Diseases*, **27**:359-368.
- Gatta, P.P., Pirini, M., Testi, S., Vignola, G. and Monetti, P.G., (2000).** The influence of different levels of dietary vitamin E on sea bass *Dicentrarchus Labrax* fish quality. *Aquac. Nutr.*, **6**:47-52.
- Gimenez, A.V.F., Fenucci, J.L. and Petriella, A.M. (2004):** The effect of vitamin E on growth, survival and hepatopancreas structure of the Argentine red shrimp *pleoticus muelleri* Bate (crustacean, Penaeidea). *AquacultureResearch*, **35**:1172-1178.
- Goda, A. M. (1997):** Types of nutrition for fresh water fish. Ph.D. Thesis, Fac. of Agriculture, Cairo University.
- Gonathan, C. and Naill, B. (1988):** Intensive fish farming. Oxford, London.

REFERENCES

- Grover, J. J., Olia, B. L., Brien, M. and Wicklund, R. I. (1989): Food habits of florida red tilapia fry in manured seawater pools in the Bahamas. *Prog Fish-Cult.*, **51**: 152-156.
- Halver, J.E. (1976): Formulating practical diets for fish. *Res. BD.Con.*, **33**:1032.
- Halver, J .E. (1989): *Fish Nutrition*. 2nd ed. New York . Academic Press.
- Hanley, F. (1991): Effect of feeding supplementary diets containing varying levels of lipid on growth, food conversion and body composition of Nile tilapia. *Oreochromis niloticus* (L.). *Aquaculture*, **93**:323-334.
- Haran, N.S. and Fenucci, J.L. (1997): Accion de acidos grasos dietarios sobre el crecimiento del langostino argentino *Pleoticus muelleri* Bate. *Revista de investigaciones Marinas*, **18**: 155-161.
- Hawkins, G.E., Commins, K.A., Silverio, M. and Jilek, J.J. (1985): Physiological effects of whole cottonseed in the diet of lactating dairy cows. *J Dairy Sci.*, **68**: 2608-2614.
- He, H. and Lawrence, A.L. (1993): Vitamin E requirement of *Pemaeus vannamei*. *Aquaculture*, **118**: 245-255.
- Hegedüs, M.; Bokori, J.; Andra Sofzki, E. and Kovari, L. (1990): Optimizing protein quality of mixtures of blood meal, feather meal and bone meal. *Aota Veterinari Hungarica*, **38**: 143-152.

- Henry, M. H., Pesti, G. M., Bakalli, R., Lee, J., Toledo, R. T., Eitenmiller, R. R. and Phillips, R. D. (2001): The performance of broiler chicks fed diets containing extruded cottonseed meal supplemented with lysine. *Poultry Sci.*, **80**: 762-768.
- Hepher, B. (1988): Nutrition of pond fishes. Cambridge, Cambridge Univ. press.
- Hepher, B. and Pruginin, Y. (1981): Commercial fish farming with species references to fish culture in Isreal. J. Wiley, New York, 261pp.
- Herman, R. L. (1970): Effects of gossypol on rainbow trout (*Salmo gairdneri*). *Journal of Fish Biology*, **2**:293-297.
- Huang, C.H., Chang, R. J., Huang, S.L. and Chen, W. (2003): Dietary vitamin E supplementation affects tissue lipid peroxidation of hybrid tilapia, *Oreochromis niloticus* × *O. aureus*. *Comp Biochem. Physiol. B Biochem. Mol. Biol.*, **134**(2): 265-270.
- Huang, S.L., and Huang, C.H. (2004): Lipid peroxidation in sarcoplasmic reticulum and muscle of tilapia is inhibited by dietary vitamin E supplementation. *J. Food Biochem.* **28**: 101-111.
- Hughes, S. G. (1988): Assessment of lupin flour as a diet ingredient for rainbow trout. *Aquaculture*, **71**: 379-385.
- Hughes, S. G. (1991): Use of lupin flour as replacement for full-fat soy in diets for rainbow trout. *Aquaculture*, **93**: 57-62.

- Hung, S.S.O.Cho, C.Y. and Slinger, S.J. (1981):** Effect of oxidized fish oil, ^{DL}- α - tocopheryl acetate and ethoxyquin supplementation on the vitamin E nutrition of rainbow trout (*Salmo gairdneri*) fed practical diets. Journal of Nutrition, **111**: 648-657.
- Jackson, A. J., Capper, B. S. and Matty, A. J. (1982):** Evaluation of some plant proteins in complete diets for tilapia *Sarotherodon mossambicus*. Aquaculture, **27**:97-109.
- Jauncey, K. (1982):** The effect of varying dietary protein level on the growth, food conversion, protein utilization and body composition of juvenil tilapias, *Sarotherodm mossambicus*. Aquaculture, **27**: 43-54.
- Jauncey, K. and Ross, B. R. (1982):** A guide to tilapia feeds and feeding. Institute of Aquaculture, University of Stirling, FKA, Scotland, V. K. L., **9**:68.
- Jobling, M. (1994):** Fish Bioenergetics. Chapman and Hall, New York, 309 pp.
- Jones, L.A. (1987):** Recent advances in using cottonseed products. Proceedings of the Florida Nutrition Conference, 12-13 March 1987, Daytona Beach, FL, pp. 119-138.
- Kanazawa, A. (1985):** Nutrition of penaid prawn shrimp. In: Proceedings of the First International Conference on Culture of Penaid Prawn Shrimp (ed.by Y. Taki., I. H. Primavera and J.A. Lobrera). PP. 123-130. Aquaculture Department. SEAFDEC, Iloilo, Philippines.

- Kaneko, J.J. (1989).** Clinical Biochemistry of Domestic Animals, 4th edn, Academic Press, New York.
- Ketola, H.G. (1975):** Mineral supplement action of diets containing soybean meal as a source of protein of rainbow trout. Prog. Fish-culture, **37**:73-75
- Kirk, R. G. (1972):** A review of recent developments in Tilapia culture, with special references to fish farming in the heated effluent of power stations. Aquaculture, **1**: 45-60.
- Kompiang, I. P. (1981):** Fish silage, its prospect and future in Indonesia. Indonesia Agric. Res. And Develop. J., **3**:1-12.
- Krogh, A. (1989):** Alternative protein sources from plants contain anti-nutrients affecting digestion in salmonids. Proceedings of Third International Symposium on Feeding and Nutrition in Fish. Tokyo Univ. of Fisheries, Tokyo.
- Krogh, A., Aea, T. B. and Rudeva, J. J. (1994):** Soybean proteinase inhibitors affect intestinal trypsin activities and amino acid digestibility in rainbow trout. Comparative Biochemistry and Physiology, **107**:215-219.
- Lee, K. J. and Dabrowski, K. (2002):** Tissue gossypol and gossypolone isomers in rainbow trout fed low and high levels of dietary cottonseed meal. J. Agric. Food Chem., **50**:3056-3061.

- Lee, K. J., Dabrowski, K., Blom, J. H., Bai, S. C. and Stromberg, P. C. (2002). A mixture of gossypol containing cottonseed meal, soybean meal and animal by-products can be used as a fish meal substitute in diets for juvenile rainbow trout (*Oncorhynchus mykiss*). J. Anim. Physiol. Anim. Nutr. , 86: 201-213.
- Lee, K. J., Rinchard, J., Dabrowski, K., Babiak, I., Ottobre, J. S. and Christensen, J. E. (2006): Long-term effects of dietary cottonseed meal on growth and reproductive performance of rainbow trout: three-year study. Animal Feed Science and Technology, 126:93-106.
- Lim, C. and Sessa, D. J. (1995). Nutrition and Utilization Technology in Aquaculture. AOCS Press, Champaign, IL.
- Lim, C., Ako, H., Brown, C. L. and Hahn, K. (1997): Growth response and fatty acid composition of Juvenile *Peneaus vannamei* fed different sources of dietary lipid. Aquaculture, 151:143-153.
- Linkongwe, J. S., Stecko, T. D., Stavffer, J. R. and Calin, R. F. (1996): Combination effects of water temperature and salinity on growth and feed utilization of juvenile Nile tilapia, *Oreochromis niloticus*. Aquaculture, 146:37-46.
- Lovell, R.T., (1978): Dietary phosphorus requirements for channel catfish, *Ictalurus punctatus*. Trans. Am. Fish Sac., 107:617.
- Lovell, R. T. (1989): Nutrition and feeding of fish. Van Nostrand Reinhold, New York.

- Lovell, R.T. and Limsuwan, T. (1982).** Intestinal synthesis and dietary non-essentiality of vitamin B₁₂ for tilapia. *Trans. Am. Fish Soc.*, **111**: 485
- Lovell, R. T., Prather, E .E.; Tres Disk, G. and Lm, C. (1974):** Inter- relationship between quality and quantity of protein in feeds for channel catfish in intensive pond culture. *Proc. Annu. Conf. Southeast Assoc. Game fish Comme*, **28**:426-461.
- Luo, L., Xue, X., Cai, H., Cao and Liang, Y. (2006):** Partial or total replacement of fishmeal by solvent-extracted cottonseed meal in diets for juvenile rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Nutrition*, **12**: 418-424.
- Magouz, F. J. (1990):** Studies on optimal protein and energy supply for tilapia (*Oreochromis niloticus*) in intensive culture. Ph. D. Thesis, Fac. Agric, Georg August Univ., Göttingen.
- Makinde, M.O. Akingbemi, B.T. and Airee, T.A. (1997):** Gossypol-induced erythrocyte fragility in the rat is ameliorated by ethanol intake. *South African Journal of Science*, **93**: 141-143.
- Maria, L. N., E. Dapkevicius, I. Batista, M. J. Nout, F. M. Rombouts and J. H. Houben (1998):** Lipid and protein changes during the ensilage of blue whiting (*Micromesistius poutassou* Risso) by acid and biological methods. *Food Chemistry*, **63**: 97-102.

- Maria, L. N., E. Dapkevicius, M. J. Nout, F. M. Rombouts, J. H. Houben and W. Wymenga (2000):** Biogenic amine formation and degradation by potential fish silage starter microorganisms. *Inter. J. Food Microb.*, **57**:107-114.
- Martin S.D. (1990):** Gossypol effects in animal feeding can be controlled. *Feedstuffs*, **62**: 14-17.
- Mbahinzireki, G.B., Dabrowski, K.J., EL-Saidy, D. and Wisner, E.R. (2001)** Growth, feed utilization, and body composition of tilapia (*Oreochromis sp.*) fed cottonseed meals-based diets in a recirculating system. *Aquaculture Nutrition*, **7**: 189-200.
- McCoy, H. D. (1990):** Fishmeal the critical ingredients aquaculture feeds. *Aquaculture magazine*, **16**:43-50.
- Middleton, T. F., P. R. Ferket, L. C. Boyd, H. V. Daniels and M. L. Gallagher (2001):** An evaluation of co-extruded poultry silage and culled jewel sweet potatoes as a feed ingredient for hybrid tilapia (*Oreochromis niloticus* × *O. mossambicus*). *Aquaculture*, **198**: 269-280.
- Millamena, O. M. (2002):** Replacement of fish meal by animal by-product meals in a practical diet for grow-out culture of grouper *Epinephelus coioides*. *Aquaculture*, **204**:75-84.
- Mires, D. (1995):** The tilapia. In: C.E. Nach and A.J. Novotny (Editors) *Production of aquatic animals*, PP: 133-152.
- Morgan, P. R. (1972):** Causes of mortality endemic tilapia of Lake Chilwa (Malawi). *Hydrobiologia*, **40**:101 –119.

- Murai, T. and Andrews, J.W. (1974):** Interactions of dietary α -tocopherol, oxidized menhaden oil and etoxyquin on channel catfish (*Ictalurus punctatus*). *Jornal of Nutrition*, **104**:1416-1431.
- Namatipour, G. R.; Brown M. L. and Gatlin, D. M. (1992):** Effect of dietary energy protein ratio on growth characteristic and body composition of hybrid striped bass (*Morone-Chrsops* females $\times$ *M. saxatilis* males). *Aquaculture*, **107**:359-368.
- National Research Council, NRC (1983):** Nutrient Requirements of warm water fish and shellfish. National Academy of Sciences, Washington D.C.
- National Research Council, NRC (1993):** Nutrient Requirements of Fish. National Academy Press, Washington, D.C.
- Nilsson, L. (1992):** Genetic parameters of growth and sexual maturity in Arabic Char (*Salvelinus alpinus*). *Aquaculture*, **106**:9-19.
- Nomeir A.A and Abou-Donia M (1982).** Gossypol-high-performance liquid chromatographic analysis and stability in various solvents. *J. Am. Oil Chem. Sac.*, **59**: 546-549.
- O'Connell, J.P. and Gatlin, D.M. (1994):** Effects of dietary calcium and vitamin D₃ on weight gain and mineral composition of blue tilapia (*Oreochromis Niloticus*) in low-calcium water. *Aquaculture*, **124**: 107-117.

- Ofojekwu, P. C. and Ejike, C. (1984): Growth response and feed utilization in the tropical cichlid, *Oreochromis niloticus* (Linn) fed on cottonseed-based diets. *Aquaculture*, **42**: 27-36.
- Okerman, H. W. and Hansen, C. L. (1994): Subproductos de origen animal. Editorial Acribia, S. A., Zaragoza, Spain.
- Oliva-Teles, A., Gouveia, J., Gomes, E. F. and Rema, P. (1994): The effect of different processing treatments on soybean meal utilization by rainbow trout. *Aquaculture*, **124**:343-349.
- Onibi, G.E., Scaife, J.R., Fletcher, T.C. and Houlihan, D.F., (1996): Influence of α -tocopherol acetate in high lipid diets quality of refrigerated Atlantic salmon (*Salmo salar*) fillets. Refrigeration and Aquaculture Proceedings of the Conference of IIR. Commission C2, 20-22 March, Paris, France. International Institute of Refrigeration, Paris, France, PP: 145-152.
- Ostrowski, A.C. and Garling, D.L. (1988): Influence of anabolic hormone treatment and dietary protein ratio on condition and muscle deposition of rainbow trout. *Prog. Fish Cult.*, **50**:133-140.
- Paul, B. N., S. Nandi, S. Sarkar, and P. K. Mukhopahdyay, (1997): Effects of feeding unconventional animal protein sources on the nitrogen metabolism in rohu *Labeo rohita* (Hamilton). *Bamidgeh*, **49** (4):183-192.

- Pauly, D. (1976):** The biology fishery and potential of *tilapia melanotheron* in a small West African lagoon. *Aquaculture*, 7:33-49
- Philippart, J. C. L. and Ruwet, J. C. (1980):** Ecology and distribution of tilapias. *In:* R.S.V. Pullin and R.H. Lowe-McConnell (Eds.), the Biology and Culture of Tilapias. ICLARM Conference Proceedings 7, International Center for Living Aquatic Resources Management, Manila, Philippines, pp: 15-59.
- Pillay, T.V.R. (1988):** Aquaculture Principles and Practices Fishing. News book, pp: 360-376.
- Ponganeerat, J. and Watanabe, T. (1991):** Nutritive value of protein of feed ingredients for carp, *Nippon suisam*. *Gaksishi*, 57:503-510.
- Popper, D. and Lichatowitch, T. (1975):** Preliminary success in predator control of *tilapia mossambica*. *Aquaculture*, 5:13-214
- Post, G. (1987):** Textbook of Fish Health. T.F.H. Publications, Neptune City, NJ, USA.
- Power, D.M. (1990):** The physiology of growth. *Trout News*, 10:20-30.
- Pullin, R. S. V. (1983):** Choice of Tilapia species for aquaculture.
L. Fishelson and Z. Yaron (Editors). International Symposium on Tilapia in Aquaculture. Green fields (Philippines) 4 (10): 24-40.

- Robinson, E. H., Rawles, S. D., Oldenburg, P. W. and Stickney, R. R. (1984 a):** Effects of feeding glandless or glanded cottonseed products and gossypol to *Tilapia aurea*. *Aquaculture*, **28**: 145–154.
- Robinson, E. H., Rawles, S.D. and Stickney, R. R. (1984 b):** Evaluation of glanded and glandless cottonseed products in catfish diets. *Prog. Fish-Cult.*, **46**:92–97.
- Rodriguez-Serna, M.; Olvera-Novoa, M. A. and Carmona-Osalde, C. (1996):** Nutritional value of animal by products meal in practical diets for Nile tilapia (*Oreochromis niloticus* L.) fry. *Aquacult. Res.*, **27** (1): 67-73.
- Roehm, J.N., Lee, D.J., Simnhuber, R.O. (1967):** Accumulation and elimination of dietary gossypol in the organs of rainbow trout. *J. Nutr.*, **92**: 425-428.
- Roem, A. J., Kohler, C. C. and Stickney, R. R. (1990).** Vitamin E requirements of blue tilapia, *Oreochromis aureus*(Steindachner), in relation to dietary lipid level. *Aquaculture*, **87**: 155-164.
- Ruff, N., Fitz Gerald, R. D., Cross, T.F. and Kerry, J.P. (2002):** Fillet shelf-life of Atlantic halibut *Hippoglossus hippoglossus* L., fed elevated levels of α -tocopherol acetate. *Aquac. Res.*, **33**:1059-1071.
- Santiago, C. B. (1985):** Amino acids requirements of Nile tilapia. Ph.D Thesis, Auburn University.

- Sarig, S. (1983):** A review on tilapia culture in Israel. International Symposium on Tilapia in Aquaculture Proceeding, pp: 116-122, Nazareth ,Israel.
- SAS, (1996).** SAS Procedure Guide "version 6.12 Ed". SAS Institute Inc., Cary, NC, USA.
- Satoh, S., Takeuchi, T., and Watanabe, T. (1987):** Requirement of tilapia for α -tocopherol. Nippon Suisan Gakkaishi, **53**:119-124.
- Schroeder, G. L. (1983):** The role of natural food in tilapia growth. A study based on stable isotope analysis. In: L. Fishelson and Z. Yaron (Editors), International Symposium on Tilapia Aquaculture Proceeding, PP: 313 - 322 Nazareth,Israel.
- Schwarz, F.J., Kirch Gessner, M., Steinhart, H. and Runge, G. (1988):** Influence of different fast with varying additions of α -tocopherol acetate on growth and body composition of carp (*Cyprinus carpio* L.). Aquaculture, **69**: 57-67.
- Sealey, W.M., Lim, C. and Klesius, P.H. (1997):** Influence of the dietary level of iron from iron methionine and iron sulfate on immune response and resistance of channel catfish to *Edwardsiella ictaluri*. J. World Aquacult. Soc., **28**: 142-149.
- Shiau S. Y. (1997):** Utilization of carbohydrates in warm water fish with particular reference to tilapia *O. niloticus* $\times$ *O. aureus*. Aquaculture, **121**: 79-96.

- Shiau, S. Y. and Huang, S. L. (1990):** Influence of varying energy level with two protein concentrations in diets for hybrid Tilapia, *Oreochromis nilotoicus* × *O. aureus* reared in seawater. *Aquaculture*, **91**: 143-152.
- Shiau, S. Y. and Shiau, L. F. (2001):** Re-evaluation of the vitamin E requirement of juvenile tilapia, *Oreochromis niloticus* × *O. aureus*. *Anim. Sci.*, **72**: 529-534.
- Siddiqui, A. and Al-Harbi, A. (1995):** Evaluation of three species of tilapia, red tilapia and hybrid tilapia as culture species in Saudi Arabia. *Aquaculture*, **138**: 147- 127.
- Siddiqui, A. Q.; Howladar, M. S. and Adam, A. A. (1988):** Effect of dietary protein levels on growth, feed conversion and protein utilization in fry and young Nile tilapia, *Oreochromis miloticus*. *Aquaculture*, **75**: 63-73.
- Spataru, P. (1982):** A contribution to the study of natural food of sarotherodon hybrids grown under condition of polyculture, supplemental feeding and intensive fertilization. *Bamidgeh*, **34**:144 – 157.
- Steffen, W. (1984):** Energy requirement and utilization in fresh water fish. *Zeitschrift Furide Binnen Fischerei der DDR*, **31**, (5) 138-143.
- Steffen, W. and Albricht, M. L. (1994):** Saving protein by increasing the proportion of dietary fat in food for rainbow trout. *Arch, Tierernahrung*, **23**:711 – 717 Berlin.

- Stephan, G., Guillame, J. and Lamour, F. (1995):** Lipid peroxidation in trout (*Scphthalmus maximus*) tissue: effect of dietary vitamin E and dietary n-6 or n-3 polyunsaturated fatty acids. *Aquaculture*, **130**:251-268.
- Stickney, R. R. and Lovell, R. T. (1972):** Nutrition and feed of channel catfish. *South. Coop. Ser. Bull.*, **218**: 67 PP.
- Stickney, R. R. and McGeachin, R. B. (1984):** Growth, food conversion and survival of fingerling *Tilapia aurea* fed differing levels of dietary beef tallow. *Progressive fish culturist*, **64**: 102-105.
- Stickney, R. R.; McGeachin, R. B.; Lewis, D. H. and Mrks, J. (1982):** Response of young channel catfish to diets containing purified fatty acids. *Trans. Am. Fish Soc.*, **11**(2): 665 – 669.
- Sun, L. T., Chen, G. R. and Chang, C. F. (1995):** Acure responses of blood parameters and comatose effects in salt-acclimated tilapia exposed to low temperatures. *Journal of Thermal Biology*, **20**: 299-306.
- Takeuchi, M. and Nakazol, J. (1981):** Effect of dietary phosphorous on lipid content and its composition in carp. *Bull. Jap. Soc. Sci. Fish*, **47**:348.
- Takeuchi, F.; Watanabe, F. and Ogimo, C. (1979):** Availability of carbohydrate and lipids as energy sources for carp. *Bull. Jpn. Soc. Sci. Fish*, **45**: 977 – 982.

- Talbot, F. H. and Newell, B. S. (1957):** A preliminary note on the breeding and growth of Tilapia in marine fish ponds on Zanzibar Island. *E. Afr. Agric. J.*, **23**: 118-121.
- Teshima, S.; Gonzalez, G. M. O. and Kamazow, A. (1978):** Nutritional requirements of Tilapia: utilization of dietary protein by tilapia zillii. *Mem. Fac., Fish Kagoshima Univ.*, **27**(1): 49-57.
- USDA, (2000) USDA—National Agricultural Statistics Service.**
- Velasquez-Pereira, J., P.J. Chenoweth, L. R. McDowell , C. A. Risco, C. A. Staples, D. Prichard, F. G. Martin, M. C. Calhoun, S. N. Williams, and N. S. Wilkinson. (1998):** Reproductive effects of feeding gossypol and vitamin E to bulls. *J. Anim. Sci.*, **76**: 2894-2904.
- Velasquez-Pereira, J., C. A. Risco, L. R. McDowell, C. R. Staples, D. Prichard, P. J. Chenoweth, F. G. Martin, S. N. Williams, L. X. Rojas, M. C. Calhoun, and N. S. Wilkinson. (1999):** Long-Term Effect of Feeding Gossypol and Vitamin E to Dietary Calves. *Journal of Dairy Science*, **82**:1240-1251.
- Velasquez-Pereira, J., Arechiga, C. F., McDowell, L. R., Hansen, P. J., Chenoweth, P. J., Calhoun, M. C., Risco, C. A., Batra, T. R., Williams, S. N. and Wilkinson, N. S. (2002):** Effects of gossypol from cottonseed meal and dietary vitamin E on the reproductive characteristics of super-ovulated beef heifers. *J. Anim. Sci.*, **80**: 2485-2492.

- Vidotti, R. M. and D. J. Carneiro (2002):** Acid and fermented silage characterization and determination of apparent digestibility coefficient for pacu, *Piaractus mesopotamicus*. J. World Aquacult. Soc., **33**:57-62.
- Viola, S.; Arieli, Y. and Zohar, G. (1988):** Unusual feedstuffs (tropica and lupin) as ingridients for carp and tilapia feed in intensive culture. Bamidgeh, **40**: 29-34.
- Wang, M .Z., Wang, J. S. and Li, B. L. (1985):** Determination of gossypol with high performance liquid chromatography. Acta Phama. Sinica, **20**: 682-687.
- Wassef, E. A. (1990):** Experiments on preparation and nutritional properties of fish silage. Comm. Sci. and Res., (Fac. Agric., Alexandria Univ.), **31**:143-160.
- Wassef, E. A. (1991):** Development of diets for gilthead bream *Sparus aurata* L. cultured in Egypt. J. King Abdul Aziz Univ., Mar. Sci., **2**:101-109.
- Wassef, E. A. (2002):** Alternative protein sources for fish feeds in Egypt. Paper presented to "Workshop Mediterranean Fish Nutrition". Rhodes (Greece), 1-2 June 2002. NCMR and CIHEAM, (Eds.):29-30.
- Wassef, E. A., M. A. Sweilam, and R. F. Attalah, (2003):** The use of fermented fish silage as a replacement for fish meal in Nile tilapia (*Oreochromis niloticus*) diets. Egypt. J. Nut. Feed, **6** (Special Issue):357-370.
- Watanabe, T. (1995):** Nutrition and growth. In: C.J. Shepherd and N. R. Bromin, Intensive fish farming .P.P. 154-197: Hartnolls Ltd, Bodnin, Corn Wall, Great Biriton.

- Watanabe, T., Takeuchi, T. and Wada, M. (1981).** Dietary lipid levels and α -tocopherol requirements of carp. Bulletin of the Japanese Society of Scientific Fisheries, 47:935-946.
- Watanabe, W.O., Clark, J. H., Dunham, J. B., Wicklund, R.I. and Olla, B.L. (1990):** Culture of Florida red Tilapia in marine cages. The effect of stocking density and dietary protein on growth. Aquaculture, 90: 123-134.
- Wilson, R.P., Robinson, E.H. and Poe, W.E. (1981):** Apparent and true availability of amino acids from common feed ingredients for channel catfish. J. Nutr., 111: 923-929.
- Wohlfarth, G. W.; and Hulata, G. (1983):** Applied genetic of Tilapias. International Center for Living Aquatic Resources Management (ICLARM), Manila, Philippines.
- Wohlfarth, G. W.; Hualata, G.; Rothbard, S.; Itzkowich, J. and Halvey, A. (1983):** Comparisons between interspecific tilapia hybrids for some production traits. In: L. Fishelson and Z. Yaron (Editors), International Symposium on Tilapia in Aquaculture, Proceeding, 559-569. Nazareth, Israel.
- Yang, W. and Xiang, S. (1995):** The separation of gossypol enantiomers by μ Bondapak C₁₈ column; Chim. J. Chromatogr., 13: 264-266.

- Yildirim, M., Lim, C., Wan, P. and Klesius, P. (2003):** Growth performance and immune response of channel catfish (*Ictalurus punctatus*) fed diets containing graded levels of gossypol-acetic acid. *Aquaculture*, **219**: 751-768.
- Yousif, O. M. (1996):** Effects of rearing conditions in closed recirculatory systems on growth, feed utilization, carcass composition and nutrient digestibility of juvenile tilapia (*Oreochromis niloticus* L.) Ph.D. Thesis, Institute of Animal Physiology and Animal Nutrition. Univ. of Göttingen, Germany.