REFERENCES

- Abdel-Gawad, E.M. (1970). Heritability estimates and types of gene action of some economic traits in "Dokki 4". 2. Heritability and types of gene action on 8 and 12 week body weight. Agric. Res. Rev. Cairo. 48:53-62.
- Ahlawat, S.P.S; Chaudhary, R.P. and Singh, B.P. (1980). Genetic and phenotypic correlation for some productive traits in the Babcock strain of White Leghorn. Anim. Breeding Abs. 49: 2947.
- Al-Mufti, A.J. (1978). Improving broiler weight in the White Baladi on chicken. M.Sc. Thesis. Cairo University. Egypt.
- Al-Soudi, K.A. and Bernier, P.E. (1973). Selection for growth in Coturnix Japanica on a low protein ration. Anim. Breeding Abs. 42:4004.
- Amer, M.A.F. (1965). Heritability of body weight in Fayoumi. Poultry Sci. 44:741-744.
- Anderson, R.L. and Bancroft, T.A. (1952). Statistical theory in research. 399 pp. McGrow-Hill Book Co., New York.
- Awad, S.H. (1978). Selection for body weight in chickens. M.Sc. Thesis. Cairo University. Egypt.
- Becker, W.A. (1975). Mannual of quantitative genetics. Washington State University, Press, Pullmon, Washington 99163.
- Carte, I.F. and Siegel, P.B. (1970). Scaling effects and the inheritance of jouvenile body weight in chickens. Anim. Breeding Abs. 41:882.
- Conkov, C.; Cicibaba, V.; Isaev, G.; Ingatov, S. and Konkov (1971).

 Some genetic parameters in Cornish and White Plymouth Rock lines
 Anim. Breeding Abs. 39:3978.
- Dillard, E.U.; Dickerson, G.E. and Lanorux, W.F. (1953). Heritabilities of egg and meat production qualities and their genetic and environmental relationships in New Hampshire pullets. Poultry Sci. 32:897-899.
- Duncan, B.D. (1955). Multiple range and multiple F. test. Biometrics. 2:1-42.
- Dutt, M.; Basu, S.B. and Sengar, O.P.S. (1971). Prediction equations for meat form live measurements in broilers. Indian J. of Poult. Sci. 6:7-12.
- El-Hossari, M.A. and Ragab, M.T. (1970). Correlated response to selection for high body weight at 8 weeks of age in a closed flock Fayoumi fowl. U.A.R. J. Animal Production 10:37-42.
- Erasmus, J.E. (1965). Selection for 12 week body weight in chickens. Anim. Breeding Abs. 34:1649.
- Falconer, D.S. (1953). A symmetrical response in selection experiments. I.LL. B.S. Symposium on population structure, pp 16-41.

Falconer, D.S. (1960). Introduction to quantitative genetics. Ronald Press Co., New York. -(1967). Introduction to quantitative genetics. Ronald Press Co., New York. - and King, J.W.B. (1953). A study of selection limits in the mouse. J. Genetics 51:561-581. Frairs, G.W. and Tiku, M.L. (1969). The effect of selection for early growth on the market grade of turkeys. Poultry Sci. 49:150-154. Goodman, B.L. (1973). Heritabilities and correlations of body weight and dressing percentages in broilers. Poultry Sci. 52: 379-381. - and Godfrey, G.F. (1956). Heritabilities of body weight in domestic fowl. Poultry Sci. 35:50-53. Gupta, P.C. and Agarwala, O.P. (1970). Relationship of dressed and live weights in broilers. Anim. Breeding Abs. 38:2984. Gyles, N.R. and Thomas, J.E. (1963). Divergent selection for eight week body weight in White Wyandottes. Poultry Sci. 42:1273-1274. Halaj, M. (1975). The relationship between body weight and some carcass parts in growing fowls. Anim. Breeding Abs. 9:4244. Hanafi, M.S. (1966). Estimation of genetic parameters under optimal and suboptimal environments for body weight in chickens using a diallel mating system. Ph. D. Thesis. Texas A&M University. Hazel, L.N. (1943). (Cited by Lerner, 1961). Hurry, H.F. and Nordskog, A.W. (1952). Statistical errors in grading chicks for feather growth. Poultry Sci. 31:819-821. - (1953). A genetic analysis of chicks feathering and its influence on growth rate. Poultry Sci. 32:18-25. Ideta, G. and Siegel, P.B. (1966). Selection for body weight at eight weeks of age. 4. Phenotypic, genetic and environmental correlations between selected and unselected traits. Poultry Sci. 45:923-933. Jaap, R.G. (1966). Response to selection for growth rate varies with strains. Anim. Breeding Abs. 35:3039. - (1968). Additive and non-additive genetic effects on growth rate in domestic fowl. Anim. Breeding Abs. 37:1964. - and Smith, J.H. (1959). Selection for rapidity of growth. Poultry Sci. 38:1214-1216. - and Goodman, B.L. (1962). A genetic analysis of growth and egg production in meat type chickens. Poultry Sci. 41:1446.

- Johanson, A.S., and Asmundson, V.S. (1957). Genetic and environmental factors affecting size of body and body parts of turkeys. 2. The relation of body weight and certain body measurements to pectoral and tibial muscle weights. Anim. Breeding Abs. 26:1016.
- Kan, J.; Krueger, W.F. and Quisenberry, J.H. (1959). Non-additive gene effects of six broiler traits as studied from a series of diallel matings. Poultry Sci. 38:972.
- Kumar, J. and Acharya, R.M. (1980). Genotypic and phenotypic parameters of growth and carcass yield of desi chickens. Anim. Breeding Abs. 49:2956.
- Lerner, I.M. (1954). Genetic homeostasis. Oiver and Boyd, Edinburg, Tweeddale court.
- Wiley & Sons, Inc., London.
- Lien, S. (1974 b). Genetic and phenotypic parameters for live weight and some carcass quality characters in broilers. Anim. Breeding Abs. 42:4010.
- Lush, J.L. (1845). Animal breeding plans. 3rd ed. Iowa State College Press, Ames, Iowa.
- (1948). The genetics of populations. Mimeographed Notes. Iowa State College, Ames, Iowa.
- Iowa, J. Agric. Sci. 40:3.
- Maloney, M.A. and Gilbreath, J.C. (1962). Selection for body size in broiler chickens. Poultry Sci. 41:1661-1663.
- selection for body weight in chickens. 2. The effect of selection for body weight on other traits. Poultry Sci. 42:338-344.
- Marks, H.L. (1970). Selection for four body weight in Japanese quail under two nutritional environments. Poultry Sci. 50:931-937.
- and 8 week body weight. Anim. Breeding Abs. 48:4263.
- _____(1980 b). Reverse selection in a Japanese quail line previously selected for 4-week body weight. Anim Breeding Abs. 49:1037.
- and Lepore, P.D. (1968). Growth rate inheritance in Japanese quail. 2. Early response to selection under different nutritional environments. Poultry Sci. 47:1540-1546.
- Martin, G.A.; Glazener, E.W. and Blow, W.L. (1953). Efficiency of selection for broiler growth at various ages. Poultry Sci. 32:716-720.
- Masic, B. (1972). Determination of body conformation and meatness, their

- inheritance and importance for selection in broilers. Anim. Breeding Abs. 41:904.
- Mather, K. and Harrison, B.T. (1949). (Cited by Lernet, 1961).
- Mc-Cartney, M.G.; Nestor, K.E. and Harvey, W.R. (1968). Genetics of growth and reproduction in the turkeys. 2. Selection for increased body weight and egg production. Poultry Sci. 47:981-990.
- McClung, M.R. (1958). Heritability and genetic correlations between growth rate and egg production in meat type chickens. Poultry Sci. 37:1225.
- Merritt, E.S. (1966). Estimates by sex of genetic parameters for body weight and skeletal dimensions in a randombred strain of meat type fowl. Poultry Sci. 45:118-125.
- reproductive traits in a randombred control strain of meat-type fowl. Poultry Sci. 47:190-199.
- Mukherjee, T.K. and Frairs, G.W. (1970). Heritability estimates and selection responses of some growth and reproductive traits in control and early growth selected strains of turkeys. Poultry Sci. 49:1215.
- Nestor, K.E.; McCartney, M.G. and Harvey, W.R. (1966). Genetics of growth and reproduction in the turkey. L. Genetic and non-genetic variation in hedy weight and body measurements. Poultry Sci. 46:1374.
- Reddy, P.M.; Sarma, P.L.N. and Reddy, C.V. (1978). Heritability estimates and genetic correlations among production traits in four White Leghorn strains. Anim. Breeding Abs. 49:3610.
- Ricard, F.H. and Rouvier, R. (1976). Genetic variation in weights of carcass parts in poultry and its application to selection. Anim. Breeding Abs. 46:5062.
- Saeki, Y.; Akita, T. and Onishi, N. (1969). Breeding chickens for meat production. VII. Age for efficient selection for rapid weight gain in broilers. Anim. Breeding Abs. 38:843.
- Siegel, P.B. (1962). Selection for breast angle at eight weeks of age.

 2. Correlated responses of feathering, body weights and reproductive characteristics. Poultry Sci. 42:437-449.
- breast angle at eight weeks of age in chickens. Genetics. 47:954-962.
- (1962 c). Selection for body weight at eight weeks of age.

 I. Short term response and heritability. Poultry Sci. 41:954-962.
- (1963 a). The correlated response of relative aggressiveness to selection for body weight and breast angle in chickens. Poultry Sci. 42:1208-1211.
- _____ (1963 b). Selection for body weight at eight weeks of age.

- 2. Correlated responses of feathering body weight and reproductive characteristics. Poultry Sci. 42:895-905.
- Siegel, P.B. (1970). Selection for juvenile body weight in chickens. Anim. Breeding Abs. 41:1892.
- sex differences and phenotypic correlation for six feathering characteristics. Poultry Sci. 36:621-628.
- of live measurement and eviscerated weight in broilers. Poultry Sci. 38:530-532.
- Singh, H.N. (1979). Estimates of various genetic and phenotypic parameters in the purebred and crossbred populations of two broiler breeds. Anim. Breeding Abs. 49:1684.
- cornish chicks. Anim. Breeding Abs. 48:7550.
- Smith, H.F. (1936). (Cited by Lerner, 1961).
- Tebb, G. (1957). Inbreeding effects contrary to the direction of selection on a poultry flock. Poultry Sci. 36:402.
- Tosovsky, R. and Jandac, J. (1975). Shank length and carcass quality of broilers. Anim. Breeding Abs. 46:1894.
- Wigan, L.G. and Mather, K. (1942). (Cited by Lerner, 1961).