

References

Abbott Laboratories (1982):

Toxicology Profile: Dipel, *Bacillus thuringiensis* Insecticide. Chemical and Agricultural Products Division, North Chicago, IL, .10-149.

Abdeen S. O.; Gad Allah, A. I.; Saleh W. S.; Hussein, N. M. and Abd El- Lateif, M. F. (1988):

Some Biochemical effects of diflubenzuron on the American boll worm, *Heliothis armigera* (Hbn.) Ann. Agric. Sci., Fac. Agric. Ain shams Univ., 3 (2); 1469 – 1478.

Abd El-Aal, A.E. (2003):

Effect of some insect growth regulators on certain biological, biochemical and histological aspects of the cotton leaf worm, *Spodoptera Littoralis* (Boisd) (Lepidoptera : Noctuidae) Ph. D. Thesis, Fac. of Science, Cairo Univ. Egypt.

Abd El-Aal, A.E and El Sheikh, T. A. (2005):

Effect of two Biocides and two Insect Growth Regulators on esterases activity of cotton leaf worm *Spodoptera Littoralis* (BOISD), Egypt. J. Agric. Res., 2(1):401-409.

Abd El-Aziz, S.H. (2000):

Physiopathological studies on bacterial infection of cotton leafworm, *Spodoptera Littoralis* .M.Sc.Thesis, Fac.of Scie.Ain shams Univ.

Abd El-Fattah, M. S; Shaaban, M. N. and Omar, M. M. (1986):

Effect of diflubenzuron and triflumuron on trehalase,

invertase and amylase enzymes activities in susceptible and profenfos-resistant strains of *Spodoptera littoralis* Bull. Ent. Soc. Egypt, Econ. Seri., 15: 207-219.

Abd El- Haleem, M. S. (1993):

Bioactivity of Dipel –2x, a commercial preparation of *B. thuringiensis* Berliner against the cotton leafworm *Spodoptera littoralis* (Boisd.). Egypt J. Agric. Res., 71 (1): 175-183.

Abd El- Haleem, M. S. (1997):

Effect of certain bio-control agents against the cotton leafworm *Spodoptera littoralis* (Boisd.) in comparison with methomyl. Egypt J. Biolog. Pest Control 7(1):35-39.

Abd El- Lateif, E.M. (2001):

Integrated pest management for cotton in Dakahlia governorate. Ph. D Thesis, Fac. of Agric, Mansoura Univ., Egypt.

Afifi, D. E. (2001):

Biochemical and toxicological studies on the effect of some citrus essential oils and biocides on the cotton the pink bollworm,=*Pectinophora gossypiella*. M.Sc. Thesis Fac. of Agric. Cairo Univ. Egypt.

Amr, M.M.; El-Batanouni, M.; Emara, A.; Mansour, N.; Zayet, H.; Atta, G.A. and Sanad, A. (1990):

Health profile among workers chronically exposed to pesticides. Project of Pesticide Intoxication, Egypt. 161-198.

Armando M., Frederic J.S., and Theodore, A. S. (2004):

Developmental effects of chlorpyrifos extend beyond neurotoxicity: critical periods for immediate and delayed-onset effects on cardiac and hepatic cell signaling. Environ Health Perspect. 112(2): 170–178.

Ashry, K.M.; Abu-Qare, A.W.; Saleem, F. R.; Hussein, Y. A.; Hamza, S.M.; Kishk, A.M. and Abou-Donia, M.B. (2002):

Inhibition and recovery of maternal and fetal cholinesterase enzymes following a single oral dose of chlorpyrifos in rats, Archives of Toxicol. 76(1): 30-39

Baker, R. F.; Abd El- Razak, N. A.; Hamed, M. S. and Gunedy, A. M. (1991):

Physiological effect of some insect growth regulator on the respirometric movements, total protein and free aminoacides of the housefly ,musca domestica, Ain shams science Bull, V. 28B: 169- 177.

Bakke, J. E.; Feil, V. J. and Price, C. E. (1987):

Rat urinary metabolites from O,O-diethyl-O-(3,5,6-trichloro-2-pyridyl) phosphorothioate. J. Environ. Sci. Health, part B 11, 225-230.

Bancroft, J. D. and Gamble, M. (2002):

Theory and Practice of Histology. 5th ed. Churchill Livingstone. Edinburgh, New York, London, 63-139.

Baronos, H. and Blackstock, J. (1973):

Estimation of lipids in marine animals and tissue: Detailed investigation of the sulphophosphovanillin method for total lipids. J.Exp.Mar.Biol.,Ecol.,12:103-118.

Bassel, T. T. and Ismail, I. E. (1985):

Acid and alkaline phosphatases of normal and juvenilized *Spodoptera Littoralis* during metamorphosis, Proc. Zoll. Soc. Egypt, 9:249-256.

Bearden, J. H. and Fuguay, J. W. (1980):

“Applied Animal Reproduction” Reston Publishing Co., Inc. Reston, Virginia, p. 158-165.

Bessey, O. A.; Lowry, O. H. and Brock, M. J. (1964):

A method for rapid determination of alkaline phosphatase with five cubic millimeters of serum. J. Biol. Chem., 164: 321-329.

Bishop, M. L.; Engelkrik, J. L. and Fody, E. P. (1996):

Clinical Chemistry: Principles, Procedures, Correlations. Lippincott. Philadelphia, New York.

British Columbia Ministry of Health (1992):

Bacillus thuringiensis. Report 1-46.

Blumbach, K. A.; Deger, H.M. and Dekant, W. (2000):

Biotransformation and male rat-specific renal toxicity of diethyl ethyl and dimethyl methylphosphonate. Toxicol sci.2000; 53(1):24-32.

Cappellozza, L.; Benedetti, R. and Cappellozza, S. (1995):

Effectiveness of tebufenozide (RH-5992) on different larval instars of *Bombyx mori* (Lepidoptera, Bombycida). Infrmatore Fitopatologico. 45(9): 52-56.

Coronado, GD; Thompson, B.; Strong, L.; Griffith, W.C. and Islas, I. (2004):

Agricultural task and exposure to organophosphate pesticides among farmworkers. Environ Health Perspect. 2004;112:142–147.

Cox C., (1997):

Subterranean Termites. Journal of Pesticide Reform 17(2): 1-20.

Clarke, B. S. and Jewess, P. J. (1990):

The inhibition of chitin synthesis in *S. littoralis* larvae by

Flufenoxuron, Teflubenzuron and Difiubenzuron. Pestic. Sci., 28:377- 388.

Cranshaw' W. S. (2006):

Bacillus thuringiensis. Colorado State University Cooperative Extension Entomologist and Professor, Bioagric. sciences and pest. manag. 11/99.

Curl, C.L.; Fenske, R.A. and Elgethun, K. (2003):

Organophosphorus pesticide exposure of urban and suburban preschool children with organic and conventional diets. Environ Health Perspect. 2003; 111: 377–382.

Dacie, S. J. and Lewis, S. M. (1986):

“Basic Hematological techniques” In Practical Hematology. International Student 6th ed. Churchill Livingstone. Edinburgh, London, Melbourne, and New York.

Daoust, A. R. (1993):

Commercialization of *Bacillus thuringiensis* and other bacterial Insecticides. In “The biopesticide *Bacillus thuringiensis* applications and in developing countries.” Salama, S. H.; Morris, N.O. and Rached, E. (Eds.). pp. 267-281. National Research Center, Cairo, Egypt and international Development Research Center (IDRC), Ottawa, Canada.

Degheele, D.; Yi, S. X. and Bai C. (1993):

Toxicity of benzylphenylureas to the African armyworm *Spodoptera exempta*. Crop protection, 12(1):35-38.

Dosoky, H. M. W. (1998):

Susceptibility of *Spodoptera littoralis* larvae to the microbial insecticide Delfin in the cotton and clover fields. J. Appl. Sci., 13(1): 242 –249.

Eaton, D.L. and Klaassen, C. D. (2001):

Principles of toxicology. In "Casarett and Doull's toxicology. The Basic science of Poisons". 6th ed. Mc Graw-Hill. Medical Publishing Division. New York, Chicago, London

Eid, M.A.; Saleh, W.S.; Ahmed, N.M. and El Sheikh, A.A. (1992):

Latent effects of some organophosphorus insecticides on *Spodoptera littoralis* (Boisd). Egypt J. Agric., 70 (2)., 435-442.

El- Dosoky, I. M (2000)

Biochemical studies of some insecticidal effect on environmental pollution. Ph.D Thesis, Fac. of Agric. Zagazig Univ.

El- Hamaky, M.A; Refaei A.F., Hegazy M.A. and Hussein N. M. (1990):

Knock-down and residual activity of certain insecticides Mededelingen Van De Faculteit Land bouwwetenschappen Rijksuniversiteit Gent, 55 (2b): 601 – 607.

El- Kashory, A.A. (1999):

Sub chronic toxicity studies of Imimacloprid, Profenofos and Carbosulfan and their mixtures on albino rats Ph.D. Thesis, Fac. of Science, Cairo Univ., Egypt.

El-Kattan, N. A. (1995):

Physiopathological studies on the Indian meal moth, *Plodia interpunctella* HB. (Pyralidae : Lepidoptera) infected with microbial entomopathogens. Ph.D. Thesis, Fac. of Agric., Ain Shams Univ., Egypt.

El- Negoly, A. E. (1998):

Studies on the effect of some bacterial strains locally producwd

on the cotton leaf worm. Msc. Thesis, Al- Azhar Univ., Egypt.

El-Nemaky, I. H. (2000):

Studies on the effect of integrated pest management elements on physiology of bollworms. M.Sc. Thesis, Fac. Agric., Al-Azhar Univ.,

El-Sayed, F.H. (2005):

Efficacy of some chitin synthesis inhibitors and their mixtures with mineral and vegetable oils and conventional insecticides against the Cotton leafworm, *Spodoptera littoralis* (Boisd), Ph.D. Thesis, Fac. of Agric, Cairo Univ.

El- Sayyad, H. I.; El- Sherbiny, S.A. and El- Gammal, L.H. (1995):

Studies on the toxic effects of diflubenzuron and Chlorfluazuron on testis of albino rats J. Egypt. Ger. Soc. Zool, 17(c):21-41.

EL-Sayyad, H.I. and Karim, S.A. (1998):

Effects of flufenoxuron-an environmental antimoulting insecticide on gonad function of albino rats; Egypt, Ger. Soc. Zool., Vol. 25(B), 45-64.

El- Sheikh, T. A. (2002):

Effects of application of selected insect growth regulators and plant extracts on some physiological aspects of the black cutworm, *agrotis ipsilon* (Hufn) Ph.D. Thesis, Fac. of Science., Ain Shams Univ.

El- Sheikh, T. A.; Abd El-Khalik' F. A. and Abd El-Aal, E. (2005):

Biochemical and histological effects of Flufenoxuron and two biocides on the cotton leafworm, *Spodoptera littoralis* (Boisd). Egypt. J. Agric. Res., 2(3):611- 672.

El -Sherbiny, S. A.; El Sayyad, H. I.; El- Gammal, H. L and El-Shershaby, E. M. (1995):

Studies on the toxic effects of Diflubenzuron and Chlorfluazuron on the testis of albino rat, Egypt. Ger. Soc. Zool., Vol. 17(C), 45-64.

El-Sweirky, F.I. (1994):

Integrated Management of the cotton leaf worm control with the minimizing environmental pollution by insecticides M.Sc. Thesis, Institute of Environ. Studies and Research, Ain Shams Univ., Egypt.

El-Sweirky, F.I. (2002):

Ovicidal activity of natural and synthetic compounds against Cotton leaf worm) ph. Thesis, Institute of Environ. Studies and Research, Ain Shams Univ., Egypt.

Ellman, G. L.; Courtney, K. D. and Andres, V. Jr. (1961):

A new and rapid colorimetric determination of acetyl cholinesterase activity. Biochem. Pharmacol. 7: 88-95.

Emara, A.S.; Moawad, M.G. and Gadallah, I. A. (1991)

Effect of three formulations of *Bacillus thuringiensis* on the American bollworm, *Heliothis armigera* (Hbn). 4th Arabic Conf. of Plant Protection, 1-5. Sydney. 11- 34.

Farag, A.M. (2001):

Biochemical studies on the effect of some Insect Growth Regulators on the cotton leaf worm, M.Sc. Thesis, Fac. of Agric., Cairo Univ.

Firdaus, K.; Shakoori, A. R.; Hafeez, M. A. and Ali, S. S. (2003):

Short term effects of chlorpyrifos on hematology and biochemical components of blood of the Sprague-Dawley rat.

Pakistan-Journal-of-Zoology. 2003; 35(3): 237-243.

Fisk, T. and Wright D. J. (1992):

Response of spodoptera exempta (walk) larvae to simulated field Spray applications of acylurea insect growth regulator with observation on cuticular uptake of acyl ureas . pesticide .sci, 35: 321- 330.

Fouad, H. (2004):

Toxicological studies some cotton insect pests. M.Sc. Thesis, Fac. of Agric.,Al azhar Univ.

Gad Allah, A. I.; Moawad, G. M.; Abbas, M. G. and Emara, S. A. (1990):

Biological and biochemical effects of the juvenile hormone mimic, S- 31183 on the American bollworm, *Heliothis armigera* (Hb.) (Lepidoptera: Noctuidae). Bull. Ent. Soc. Egypt, Econ, Ser., 18: 125-136.

Gallo, M.A. and Lawryk, N. J. (1991):

Organic phosphorus pesticides. In Hand book of pesticide toxicology Hayes, W. J., Jr. and laws, A.R., Jr., Eds. Academic Press New York Inc. San Diego, New York, London, Sydney, Tokyo, Toronto p.917-960.

Gartner, L. P. and Hiatt, J. L. (2001):

Color Textbook of Histology. 2nd ed. WB Saunders Co.

Philadelphia, New York, London. Gilbert, L. I.; W. Goodman.

Gilbert, L. I.; Goodman, W. and Bollenbacher, W. E. (1977):

Biochemistry of regulatory lipids and sterols in insects.

Biochemistry of lipids II, International Review of Biochem. (T. W. Goodwin, Ed.). 14: 1.

Goel, A.; Chauhan, D. P. and Dhawan, D. K. (2000):

Protective effects of zinc in chlorpyrifos induced hepatotoxicity. A biochemical and trace elemental study. Biological-Trace-Element- Research.; 74(2): 171-183.

Gomes, J.; Dawodu, A. H.; Lloyd, O. and Anilal, S.V. (1999):

Hepatic injury and disturbed amino acid metabolism in mice following prolonged exposure to organophosphorus pesticides, Human and Experimental Toxicology; 18(1): 33.

Haga, T.; Toki, T.; Koyangio, T. and Nishiyamo, R. (1984):

Structure-activity relationships of benzoylphenylurea. Internat. Cong. Entomol., Hamburg (FRG).

Harris, J. M. (1995):

Clinical Chemistry Interpretation and techniques. 4th ed.

Williams and Wilkins. Baltimore, Philadelphia, Hong Kong, London.

Hatem, A. E. (2006):

Comparacion de los efectos de insecticidas selectivos sobre el desarrollo y reproduccion de *S. littoralis* Lepidoptera : Noctuidae). { Comparison of the effects of insecticides on the *S. littoralis*} Thesis doctoral {Ph.D} Universidad {Univ.} de Cordoba .240.

Hegazy, G. (1990):

Comparative study of certain benzoylphenyl ureas on the integument ultra structure of the first instar larvae of *Spodoptera Littoralis* (Boisd). Ann. Agric. Sci, Special Issue, 521 – 529.

Hesham, F. H. and Hamouda, S. H. (1998):

Toxicity and residual activity of some pyrethroids in comparison with organophosphate and carbamate insecticides

against at three insect pests. J. Agric. Sci. Mansoura Univ., 23(11) : 5111- 5118.

Hessian, A. M.; Makkar, A .W.; Soker, A. L.; Allam, A. M. and Mustafa, S.A. (1996):

The bioactivity of certain benzoylphenylureas on the cotton leaf worm *S. littoralis* moth's survival longevity and fecundity. Ann. Agric. Science, Moshtohor, 34(3): 1263-1276.

Hossani, M. G.; Zimmermann, G. A.; Langen, B. and Vidal, S. (1998):

Screening of four bio-insecticides against the two cotton pests, *Spodoptera Littoralis* and *Helicoverpa armigera* (Lepidotera : Notuidae). Proceedings 5th International symposium on crop protection, Gent, Belgium, 5 may, 1998. Part II. Mededelingler faculteit land bouwkunding En Toegepaste. Biologische Wetenschappern Univ. Gent, 1998, 63 (2b): 461 – 645.

Ishaaya, I.; Moore, I. and Joseph, D. (1971):

Protease and amylase activity in larvae of the Egyptian cotton leafworm, *S. littoralis*. J. Insect Physiol., 17: 945-953.

Ivey, M. C.; Mann, H. D.; Oehler, D. D.; Claborn, H. V.; Eschle, j. L. and Hogan, B. F. (1988):

Chloropyrifos and its oxygen analogue: Residues in the body tissues of dipped cattle. J. Econ. Entomol. 65, 1647-1649.

Jabbar, A., Iqbal, A. and Malik, S. A. (1994):

Hepatotoxic effects of organophosphate insecticides.

Proceedings of Pakistan Congress of Zoology,-vol-12: 579-587.

Johnson, K.A. (2005):

Toxicology & Environmental Research and Consulting, Midland, MI; Laboratory Project Study 1-35.

Karim, S. A. (1998):

Patterns of developmental defects of rat fetus maternally treated with an environmental antimoulting insecticide Flufenoxuron; Egypt. Ger. Soc. Zool., Vol. 25(B), 65-81.

Kaur, S. P.; Sidhu, D. S.; Dhillon, S. S. and Kumar, N. (1985):

Transaminases during development and aging of the bruchid, *Zabrotes subfasciatus* (Coleoptera : Bruchidae). Insect Sci. Applic., 6: 585-590

Kidd, H. and James, D. R. (1991):

The Agrochemicals Handbook, Third Edition. Royal Society of Chemistry Information Services, Cambridge, UK. 10-20

Knowles, B. H. (1994):

Mechanism of action of *Bacillus thuringiensis* insecticidal delta-endotoxins. In Advances in Insect Physiology, Volume 24 (ed. PD Evans) pp. 275- 308. Academic Press, London.

Kumar, A. and Susheela, A. K. (1995):

Effects of chronic fluoride toxicity on the morphology of ductus epididymis and the maturation of spermatozoa of rabbit. New Delhi. Int J Exp Pathol 76(1):1-11

Laufer, H. and Schin, K. S. (1971):

Quatitative studies of hydrolytic enzymes activity in salivary gland of *Chironomus tentans* (Diptera : Chironomidae) during metamorphosis. Can. Entomol. 103:454-457.

Lemus, R. and Abd El- Ghany, A. (2000):

Chlorpyrifos: An unwelcome pesticide in our homes. Rev. Environ. Health. 15(4): 421-33.

Lores, E. M., Bradway, D. E. and Moseman, R. F. (1987):

Organophosphorus pesticide poisonings in humans:

Determination of residues and metabolites in tissues and urine.

Arch. Environ. Health 33, 270- 276.

Mann, H. S., Sohr A., Joginder S. and Sandahu S. S. (1995):

Biology and control of , *spodoptera Littoralis* (Boisd.) on

cotton. Jres. Punjab. Agric. Univ. 32(4): 375-378.

Marty, M.S.; Zablotny, C.L.; Liberacki, A.B.; Yano, B.L.

(2004):

Toxicology & Environmental Research and Consulting. The

Dow Chemical Company, Midland, MI; Laboratory Project

Study p.221;

Mastumura, F. (1985):

Toxicology of Insecticides. 2nd ed., Plenum Press, New York, p

267-270.

McGaughey, W.M. (1990):

Bacillus thuringiensis. In Baker, R.R. and Dunn, P.E. (eds.) New

directions. In biological control: Alternatives for suppressing

agricultural pests and diseases New York: Alan R. Liss, Inc. p.

483-498.

Merdan, A. (1993):

Application of biotechnology in pest control using *B.*

thuringiensis formulations. In “The biopesticide *B. thuringiensis*

and applications in developing countries.” Salama, S. H.;

Morris, N.O. and Rached, E. (Ed.). 197- 204. National Research

Center, Cairo, Egypt and International Development Research

Center, Ottawa, Canada.

Mohammed, F. R and Siddiqui, M. K. (2003):

Biochemical Enzyme Activity in Different Tissues of Rats

Exposed to a Novel Phosphorothionate Journal of Environmental Science and Health, Part B: Pesticides, Food Contaminants, and Agricultural Wastes Vol. 38, No. 1: 59 - 71.

Mohamed, M. H. (2003):

New approach for controlling the cotton leaf worm *S. littoralis*. M.Sc. Thesis, Fac. of Science, Mansoura Univ.

Mustafa, S. A. (1993):

Biochemical effects of some chemical compounds on *S. littoralis* (Boisd). Ph.D. Thesis. Fac. Agric, Al. Azhar Univ. Egypt.

Nasser, F. N. (1999):

New isolated *Bacillus Spp* against the cotton leaf worm, *S. littoralis* (Boisd). Egypt. J. Agric. Res. 77(4): 1573-1583.

Nath, B. S.; Suresh, B. M.; Varma, R. S. and kumar, R.P. (1997):

Bombyx mori (Lepidoptera: Bombycidae) in response to Organo- phosphorus insecticides toxicity. Ecotoxicol. and Envir. Safty, 36 : (2), 169-173.

NRC (National Research Council) (1995):

Nutrient Requirement of Laboratory Animals, 4th. Revised ed. National Academy Press. Washington, D.C.; 95-102.

Oncu M., Gultekin F.; Karaoz, E.; Altuntas, I.; and Delibas, N. (2002):

Nephrotoxicity in rats induced by chlorpyrifos-ethyl and ameliorating effects of antioxidants. Human and Experim. Toxicol. 21(4): 223-230.

PIP (Pesticide Information Project) (2006):

Bacillus thuringiensis. Report: Cornell Univ., USDA/Extension Service/National Agric. Pesticide Impact Assessment Program. 1- 37.

Prakash, M. and Arora, C. K. (1998):

Breeding in laboratory animals: 1sted. J. L. Kumar for Anmol Pub. Pvt. Ltd., New Delhi. P. 131-133.

Qiao, D.; Seidler, F.J. and Slotkin, T.A. (2005):

Oxidative mechanisms contributing to the developmental neurotoxicity of nicotine and chlorpyrifos. Toxicol. Appl. Pharmacol. 206(1):17-26.

Rao, N. V.; Rao, K. and Redy, A. S. (1994):

A note on the efficacy of IGR to caterpillar, *Helicoverpa armigera* (Hub.). J. Insect Sci., 5: 169-171.

Raslan, A. A. (1998):

Evaluation of Ecotech - Bio efficiency on the cotton leaf worm, *Spodoptera littoralis* (Boisd.) J. Appl. Sci., 13 (1): 235 – 241.

Ray, D. E., (1991):

Pesticides derived from plants and other organisms. In Handbook of Pesticide Toxicology. Hayes, W. J., Jr. and Laws, E. R., Jr., Eds. Academic Press, New York, 10-144.

Revathi, K. and Kumar, S. (2000):

Histopathological alterations induced by sub acute dosage of chlorpyrifos 40% E.C. in Swiss albino mice J. Exper. Zool.- India. 2000; 3(1): 45-47

Rizk, M .A. and Attia, B.M.(1990)

Effect of the chitin synthesis inhibitors on growth rates of the cotton leaf worm *spodoptera littoralis* Medelin genrande. Fac. land Rijks. Unvi. Gent, 55: 577-582.

Roe, R. M., (1991):

Vertebrate toxicology of the solubilized parasporal crystalline proteins of *B. thuringiensis israelensis*. In Reviews in Pesticide

Toxicology 1: Toxicological Studies of Risks and Benefits.
Hodgson, E., Roe, R. M. Eds. North Carolina State Univ.,
Raleigh NC, 10-148.

Ross, M. H. and Pawlina, W. (2005):

Histology. A text and atlas. 5th ed. Lippincott Williams and
Wilkins. New York, London, Tokyo.

**Russell, L. D.; Ettlin, R.; Sinha; H. and Clegg, E. D.
(1996):**

Histological and Histopathological evaluation of the testis.
Cache Riv. Pr. Florida. Cited In "Casarett and Doull's
toxicology. The Basic science of Poisons". 6th ed. Mc Graw-
Hill. Medical Publishing Division. N. Y., Chicago, London.

**Salamitou, S.; Ramisse, F.; Brehélin, M.; Bourguet, D.;
Gilois, N.; Gominet, M.; Hernandez, E. and Lereclus, D.
(2000):**

The plcR regulon is involved in the opportunistic properties of
B. thuringiensis and *B. cereus* in mice and insects. Microbiol.
146: 2825- 32.

Shaaban, M.N. and Mourad, E.I. (1994):

Effect of the insect growth inhibitor, flufenoxuron on
consumption and utilization of food in *Spodoptera Littoralis*
(Boisd). Zagazig J. Agric. Res., 21(5): 1547-1553.

**Shurab, E. H.; Emara, S. A.; Zohdy, N. Z. and Abdel-Aal, A.
E. (1999):**

Effect of four insect growth regulators on the black cutworm,
Agrotis ipsilon (Hufn.) (Lepidoptera: Noctuidae). The 2nd
Int. Conf. of Pest Control, Mansoura, Egypt, Sept., 1999
PP: 773-776.

Siest, G., Henny, J., Schiele, F. and Young, D. S. (1985):

Kinetic determination of creatinine. In Interpretation of Clinical Laboratory Tests. 220- 234.

Smith, G. N.; Watson, B. S. and F.S. (1985):

Investigations on Dursban insecticide. Metabolism of (³⁶Cl) O,O diethyl O-3,5,6-trichloro-2-pyridyl phosphorothioate in rats. J. Agric. Food Chem.15,132-138.

Sokar, L. A. (1995):

Possible alternatives to classical insecticides in management program of *S. littoralis* (Boisd). Ph.D. Thesis .Fac. Agric. Moshtohor, Zagazig Univ, Egypt.

Spiegel, J. P. and Shadduck, J. A., (1990):

Clearance of bacillus sphaericus and *B. thuringiensis israel.* from mammals. Econ. Entomol. 83: 347 55, .10- 150.

Srinivasan, R. and Panicker, K.N. (1994):

Life-table and intrinsic-rate of natural increase of *Dirhinus himalayanus* (Hymenoptera:Chalcididae), a pupal parasitoid of *Musca domestica* (Diptera:Muscidae). Biological Control Insect Pests:99-102.

Sviatnyi, I.M. (1992):

A hygienic evaluation of the insecticide Sonet and the characteristics of the mechanism of its toxic action in an experiment. Lik Sprava. (1): 84-7.

Taylor, A. J. and Vadgama, R. (1992):

Analytical reviews in clinical biochemistry: the estimation of urea. Ann. Clin. Biochem., 29-245.

Tasheva, M. and Hristeva, V. (1993):

Comparative study on the effects of five benzoylphenylurea

insecticides on hematological parameters in rats. J. Appl. Toxicol; 13 (1). 67- 68.

Theo, C. (2006):

A Case for Revisiting the Safety of Pesticides. Environ Health Perspect 114(1): 10–17.

Thomas, J. A. (2001):

Toxic response of the reproductive system. In "Casarett and Doull's toxicology. The Basic science of Poisons". 6th ed. Mc Graw-Hill. Medical Pub. Division. N. Y., Chicago, London.

Tsai,-S. F; Liao, J. W and Wang, S. C. (1995):

Clearance and distribution of *Bacillus thuringiensis subsp. kurstaki* from rat by oral administration. Plant Protect. Bull. Taipei.; 37(3): 265-270.

Tsumuki, H. and Kanehisa, K. (1984):

Phosphatases in the rice stem borer, *Chilo suppressalis* Walker (Lepidoptera Pyralidae) : Some properties and changes of the activities during hibernation. Cryobiology. 21: 177-182.

US EPA US Environmental Protection Agency, (1986):

Chemical Fact Sheet Number 26: Daminozide. Office of Pesticides and Toxic Substances, Washington, DC, .10-169.

US EPA US Environmental Protection Agency (1998):

Flufenoxuron. 738-R-98-023.

US EPA USEnvironmental Protection Agency. (2005):

Health Effects Division Chapter of the Tolerance Reassessment Eligibility Decision Document.

Valenzuela, A. I.; Pico, Y. and Font, G. (2001):

Determination of five pesticide residues in oranges by matrix solid- phase dispersion and liquid chromatography to estimate daily intake of consumers. J. AOAC Int. 84(3):901-9.

Verma, R. S.; Jatav, P.C. and Nalini, S. (2002):

Chlorpyrifos induced inhibition and reactivation of serum cholinesterase activity in rats. Indian J. Toxicol. 2002; 9(1): 29-33:

Vodela, J. K. and Dalvi, R. R. (1995):

Comparative toxicological studies of chlorpyrifos in rats and chickens. Veterinary and Human Toxicol. 37(1): 1-3.

Washington State Department of Health (1993):

Report of health surveillance activities: Asian gypsy moth control program. Olympia, WA., 13-76.

Watson, W.M; EL-Hamaky, M. A. and Radwan, S. M. (1988):

Evaluation of the effect of certain insecticides against different developmental stages of *Heliothis Armigera* HBN. Agric. Res. Review, 66(1): 39-46.

Whittaker, M., (1983):

“Methods of Enzymatic Analysis” V. Bergmeyer, J. Bergmeyer and M. Grabl eds. 3rd. ed., Verlag Chemie, Deerfield Beach, Florida 52-74.

Zhao, Z.L.; Wu, N.P. and Gao, W.H., (1995)

The influence of fluoride on the content of testosterone and cholesterol in rat, China Fluoride; 28(3): 128- 130