

6- REFERENCES

- Abawi, F.G., Sullivan; T.W. and Scheideler, S.E. (1985).
Interaction of dietary fat with levels of vitamins A
and E in broiler chicks.
Poultry Sci. 64: 1192- 1198.
- Allen, E.A.; Johnson, A.R.; Fogerty, A.C.; Pearson, J.A. and
Shenstone, F.S. (1967).
Inhibition by cyclopropene fatty acids of the
desaturation of stearic acid in hen liver.
Lipids, 2 : 419 - 423.
- A.O.C.S. (1975).
Official and tentative methods of analysis 2nd ed.
Chicago, Illianos, U.S.A.
- Appling, D.R.; and Chytil, F. (1981).
Evidenic of a role for retinoic acid (Vitamin A acid)
in the maintenance of testosterone production in male
rates.
Endocrinology, 108: 2120- 2123.
- Arnich, L.; and Arthur, V. (1980).
Interaction of fat soluble vitamins in hypervitaminoses
Ann. New York Acad. Sci. 35: 109- 177.
- Awad, S.T. (1981).
Some fat soluble vitamins in egg of hens according to
season, diet and breed.
Ph.D. Thesis, Faculty of Vet. Med. Cairo University.

- Bartov, I.; Lipstein, B. and Bornstein, S. (1974).
Different effects of dietary acdulated soybean oil soapstock, cottonseed soapstock and tallow on broiler carcass fat characteristics.
Poultry Sci. 53: 115- 124.
- Bartov, I.; and Bornstein, S. (1976 a).
Minimal levels of dietary oils, grain oils or supplementary oil, affecting the composition and stability of carcass fat and meat of broilers.
Poultry Sci. 55: 1936- 1046.
- Bartov, I.;and Bornstein, S. (1976 b).
Effect of degree of fatness in broilers on other carcass characteristics:
Relationship between fatness and the composition of carcass fat.
Br. Poultry Sci., 17: 17- 27.
- Bartov, I. and Bornstein, S. (1977).
Stability of abdominal fat and meat of broilers: the interrelationship between the effects of dietary fat and vitamin E supplements.
Br. Poultry Sci. 18: 47- 57.
- Bartov, I. and Bornstein, S. (1981).
Stability of abdominal fat and meat of broilers: Combined effect of dietary vitamin E and synthetic antioxidants .
Poultry Sci. 60: 1840- 1845 .
- Becker, W.A.; Spencer, J.V.; Mirosh, L.W. and Verstrate, J.A. (1979).
Prediction of fat and fat free live weight in broiler chickens using Backskin fat, abdominal fat and live body weight.
Poultry Sci. 58: 835- 842.
-

- Cherry, J.A.; Siegel, P.B.; and Beame, W.L. (1978) .
Genetic-nutritional relationship in growth and
carcass characteristics of broiler chickens.
Poultry Sci. 57: 1482- 1487.
- Combs, G.F. (1964).
Predicting amino acids requirements of chicks basal
on growth rate, body size and body composition.
Fed. Proc. 23: 46- 51.
- Combs, G.F.; Bossard, E.H.; Childs, G.R. , and
Blambery, B.L. (1964).
Effect of protein level and amino acid balance
on voluntary energy consumption and carcass
composition.
Poultry Sci. 43: 1309.
- Combs, G.F. (1978).
Studies on the utilization of vitamin E alcohol
and esters by the chick.
Poultry Sci. 57: 223 - 229.
- Couch, J.R.; Chang, W.Y. and Lyman, C.M. (1955).
The effect of free gossypol on chick growth.
Poultry Sci. 34: 178- 183.
- Cruickshank, E.M. (1934).
Studies in fat metabolism in the fowl.
The composition of the egg fat and depot fat
of the fowl as affected by the ingestion of
large amounts of different fats.
Biochem. J. 28: 965- 977.
- Curtin, J.V.; and Raper, J.T. (1956).
Feeding value of hydrolyzed vegetable fats in
broiler rations.
Poultry Sci. 35: 273- 278.

- Dam, H.; Nielsen, G.K.; Prange, I.; and Sondergaard (1958).
Influence of linaleic and linolenic acids on symptoms of vitamin E deficiency in chicks. Nature, 128: 802.
- Edwards, H.M.; Marion, J.E.; and Driggers, J.C. (1962).
Studies on fat and fatty acid requirements of poultry.
Proc. 12th World's Poultry Congress, 182.
- Edwards, H.M.; Ashour, A. and Nugara, D. (1973).
Carcass composition studies. 1. Influences of age, sex and Type of dietary fat supplementation on total carcass and fatty acids composition. Poultry Sci. 52: 934- 948.
- Edwards, H.M.; and Denman, Jr. (1975).
Carcass composition studies. 2- Influences of breed, sex and diet on gross composition of the carcass and fatty acid composition of the adipose tissue. Poultry Sci. 54: 1230- 1238.
- Essary, E.O. and Dawson, L.E. (1965).
Quality of fryer carcasses as related to protein and fat levels in the diet. 1- Fat deposition and moisture pack-up during chilling. Poultry Sci. 44: 7 - 15.
- Evans, R.J.; Bandemer, S.L.; Anderson, M.; and Davidson, J.A. (1962).
Fatty acid distribution in tissues from hen feed cottonseed oil or sterculia foelida seeds. J. Nutr. 76: 314- 319.
- Farr, A.J.; Hebert, A. and Johnson, W.A. (1977).
Studies on the effects of dietary energy levels and commercial broiler strains on live bird, dry carcass and abdominal fat weights. Poultry Sci. 56: 1713.

- Feigenbaum, A.S.; and Fisher, H. (1959).
The influence of dietary fat on the incorporation of fatty acids into body and egg fat of the hen.
Arch. Biochem. Biophys. 79: 302- 306.
- Fraps, G.S. (1943).
Relation of the protein, fat and energy of the ration to the composition of chickens.
Poultry Sci. 22: 421- 424.
- Ganguly, J.;Mehl;J.W. and Deuel, H.J. (1953).
Studies on carotenoid metabolism.
XII. The effect of dietary carotenoids on the carotenoid distribution in the tissue of chickens
J. Nutr. 50: 59- 72.
- Gallo- Torres, H.E. (1970).
Obligatory role of bile for the intestinal absorption of vitamin E.
Lipids 5: 379- 384.
- Goodman, Dew. S. (1980).
Vitamin A metabolism.
Fed. Am. Soc. Exp. Biol. 39: 2716- 2722.
- Griffiths, L.; Lesons, S.;and Summers, J.D. (1978).
Studies on abdominal fat with four commercial strains of male broiler chicken.
Poultly Sci. 57: 1198- 1203.
- Gyles, N.R.; Maeza, A.; and Goodwin, T.L. (1982).
Regression of abdominal fat in spent parents.
Poultry Sci. 61: 1809- 1814.
- Heywang, B.W. and Bird, H.R. (1955).
Relationship between the weight of chicks and levels of dietary free gossypol supplied by

different cottonseed products.
Poultry Sci. 34: 1239- 1247.

High, E.G. (1954).

Studies on the absorption, deposition and
depletion of vitamin A in the rat.
Arch. Biochem. Biophys. 49: 19- 29.

Hilditch, T.P.; Jones, E.C. and Rhead, A.J. (1934).

The body fats of the hen.
Biochem. J. 28: 786- 795.

Hill, F.W.; and Totuska, K. (1963).

Metabolizable energy of cottonseed meals for
chicks with particular reference to the effects
of gossypol.
Fed. Proc. 22: 200.

Hollander, D. (1981).

Intestinal absorption of vitamins A, E, D and K.
J. Lab. Clin. Med. 97, 449.

Horesign, W. ed. (1982).

Recent advances in animal nutrition.
Butterworths, London.

Insull, W.Jr.; and Ahrens, E.H.Jr. (1959).

The fatty acid of human milk from mothers on
diets taken ad- libilum .
Biochem. J. 72: 27- 33.

Jen, J.J.; Williams, W.P.Jr.; and Paynter, V.A. (1970).

The effect of dietary fats on the fatty acid
distribution of chicken adipose tissue.
Poultry Sci. 49: 1400- 1401.

Johnson, A.R.; Fogerty, A.C.; Pearson, J.A.; Shenstone,
F.C.; and Bersten, A.M. (1969).

Fatty acid desaturase systems of hen liver and
their inhibition by cyclopropene fatty acids.
Lipids, 4: 265- 269.

- Kubena, L.F.; Deaton, J.W.; Chen, T.C.; and Feece, F. (1974).
Factors influencing the quantity of abdominal fat
in broiler. I. Rearing temperature, sex, age or
weight and dietary choline chloride and inositol
supplementation.
Poultry Sci. 53: 211- 214.
- Leveille, G.A.; Romsos, D.R.; Yeh, Y.; and O'Hea, E.K.
(1975).
Lipid biosynthesis in the chick. A consideration
of site of synthesis, influence of diet and pos-
sible regulatory mechanisms.
Poultry Sci. 54: 1075- 1093.
- Lipstein, B. and Bornstein, S. (1963).
Studies with acidulated cottonseed oil soapstock,
1. Its use as a fat supplement in practical
broiler rations.
Poultry Sci. 42: 686- 693.
- Lipstein, B. and Bornstein, S. (1966).
Studies with acidulated cottonseed oil soapstock.
3. Its use as a fat supplemental in layer rations.
Poultry Sci. 45: 651- 661.
- Lipstein, B.; Bornstein, S. and Budowski, P. (1970).
Studies with acidulated cottonseed oil soapstock.
4- Its effect on the fatty acid composition of
broiler carcass lipids.
Poultry Sci. 49: 1631- 1638.
- Littlefield, L.H. (1972).
Strain difference in quantity of abdominal fat
in broilers.
Poultry Sci. 51: 1829.

Lyman, C. M. (1959).

Gossypol reactions with proteins. Proc. of
the Conf. on the chemical structure and rations
of gossypol and nongossypol pigments of cotton-
seeds.

National Cottonseed products Assn.

March, B. and Jacob, B. -(1956).

Fat studies in poultry, 5. The effect of dietary
fat level on the choline requirement of the chick.
Poultry Sci. 35: 545- 549.

Marion, J.E.; and Woodroof, J.G. (1963).

The fatty acid composition of breast, thigh and
skin tissues of chicken broilers as influenced
by dietary fats.

Poultry Sci. 42: 1202- 1207.

Marion, J.E.; and Woodroof, J.G. (1966).

Composition and stability of broiler carcasses
as affected by dietary protein and fat.

Poultry Sci. 45: 241- 247.

Marion, J.E.; Boggess, Jr., T.S.; and Woodroof, J.G.
(1967).

Effect of dietary fat and protein on lipid
composition and oxidation in chicken muscle.

J. Food Sci. 32: 426- 429.

Marusich, W.L.; Orginz, E.F.; Keating, J.; Mitrovic, M.
and Bunnell, R.H. (1975).

Effect of supplemental vitamin E in control of
rancidity in poultry meat.

Poultry Sci. 54: 831- 844.

Mc-Nair, H.M. and Bonelli, E.J. (1969).

Basic Gas Chromatography, 5th edition.

Lithographed by Consolidated Printers, Berkeley,
California.

Mead, J.F. (1961).

Synthesis and metabolism of polyunsaturated acids.

Fed. Proc. 4: 952- 955.

Mecchi, E.P.; Pool, M.F.; Nonaka, M.; and Klose, A.A.

(1956 a).

Further studies on tocopherol content and
stability of carcass fat of chicken and turkeys.

Poultry Sci. 35: 1246- 1251.

Mecchi, E.P.; Pool, M.F.; Behman, G.A.; Hamachi, M. and

Klose, A.A. (1956 b).

The role of tocopherol content in the compara-
tive stability of chicken and turkey fat.

Poultry Sci. 35: 1238- 1246.

Melton, S.L. (1983).

Methodology for following lipids oxidation
in muscle foods.

Food Technol. 37: 105- 111.

Merkely, J.W.; Littlefield, G.W.; Malone, G.W. and Chaloupka,

G.W. (1977).

Fresh eviscerated yields of five commercial broiler
strains.

Poultry Sci. 56: 1738.

- Middilknoop, J.H.; Kuit, A.R. and Zegwaard, A. (1977).
Genetic factors in broiler fat deposition.
Br. Poultry Sci., 18: 131- 143.
- Mirosh, L.W.; Becker, W.A.; Spencer, J.V. and Verstrate, J.A. (1980).
Production of abdominal fat in live broiler chickens.
Poultry Sci. 59: 945- 950.
- Morita, A. and Nakano, K. (1982).
Change in vitamin A content in tissues of rats fed on a vitamin A- free diet.
J. Nutr. Sci. Vitaminol, 28: 343- 350.
- Nair, J.H. (1964).
Dry soaps and other dry mixes. Food Dehydration Vol. II- Eds. W.B. Van. Arsdell and M.J. Copley.
The Ari Publishing Co., Inc., Westport, Conn.
- Neill, A.R.; Reichmann, K.G. and Connor, J.K. (1977).
Biochemical, Physiological and production indices related to fat metabolism in the laying fowl at various stages of physiological development.
Br. Poultry Sci. 18: 315- 324.
- Nir, I. and Ascarelli, I. (1967).
Effect of dietary protein level and thyroxine on vitamin A absorption and on plasma proteins in chicks.
Br. J. Nutr. 21: 167 - 180.
- Batrack, H. and Schaible, P.J. (1980).
Nutrition of fat-soluble vitamins.
Poultry feed and nutrition 2nd, 143.

Pepper, W.F.; Snyder, E.S.; Sibbald, J.R. and
Slinger, S.J. (1962).

The effect of cottonseed oil and cottonseed
derivatives on the quality of eggs stored at
30°C and 60°F. for varying periods of time.
Poultry Sci. 41: 1943- 1946.

Pereira, A.S.; Evans, R.W. and Stadelman, W.J. (1976).

The effect of processing on some characteris-
tics, including fatty acid composition of
chicken fat.

Poultry Sci. 55: 510- 515.

Pikul, J.; Leszczynski, D.E.; and Kummerow, F.A. (1985 a).

Influence of fat content and composition on
malonaldehyde concentration in chicken meat
and skin.

Poultry Sci. 64: 311- 317.

Pikul, J.; Leszczynski, D.E. and Kummerow, F.A. (1985 b).

Total lipids, fat composition and malonaldehyde
concentration in chicken liver, heart, adipose
tissue and plasma.

Poultry Sci. 64: 469- 475.

Porter, P.J. and Britton, W.M. (1974).

Fatty acid composition of chick fed full-fat
soybeans.

Poultry Sci. 53: 1137- 1141.

Rand, N.T.; Kummerow and Scott, H.M. (1957).

The relationship of dietary protein, fat and energy on the amount, composition and origin of chick carcass fat.
Poultry Sci. 36: 1151.

Reiser, R. (1950).

The metabolism of polyunsaturated fatty acids in growing chickens.
J. Nutr. 42: 325- 335.

Reiser, R. (1951).

The synthesis and interconversion of polyunsaturated fatty acid by the laying hen.
J. Nutr. 44: 159- 175.

Rinehart, K.E.; Green, D.E. and Williamson, J.L. (1975).

The influence of selected nutrition and management factors on broiler carcass composition.
PoultrySci. 54: 1809- 1810.

Salmon, R.E. and O'Neil, J.B. (1971).

The effect of the level and source of dietary fat on growth, feed efficiency, grade and carcass composition of turkeys.
Poultry Sci, 50: 1456- 1467.

Singsen, E.P.; Matterson, L.P.; Kozeff, A.; Bunell, R.H.; and Jungherr, E.G. (1954) .

Studies on encephalomalacia in the chick. The influence of vitamin E, deficiency on the performance of breeding hens and their chicks.
Poultry Sci. 33: 192.

- Tzeng, R.Y. and Becker, W.A. (1981).
Growth patterns of body and abdominal fat
weight in male broiler chickens.
Poultry Sci. 60: 1101 - 1106.
- Underwood, B.A.; Lorech, J.D. and Lewis, K.C. (1979).
Effect of dietary vitamin A deficiency,
retonic acid and protein quality and quantity
on serially obtained plasma and liver levels
of vitamin A.
J. Nutr. 109: 796- 806.
- Velu, J.G.; Baker, D.H. and Scott, H.M. (1972).
Regression equation for determing body
composition of young chicks.
Poultry Sci. 51: 698- 699.
- Vogel, A.R. (1975).
A textbook of practical organic chemistry
3rd ed. Book Society any Longman Group Ltd.
- Weeb, J.E.; Brunson, C.C.; and Yates, J.D. (1973).
Effect of feeding fish meal and tocopherol
on flavor of precooked, frozen turkey meat.
Poultry Sci. 52: 1029- 1043.

- Siieldon, B.W. (1984).
Effect of dietary tocopherol on the oxidative stability of turkey meat.
Poultry Sci. 63: 673- 681.
- Snedecor, G.W. and Cochran, W.G. (1967).
Statistical methods.
Six th Edition. Oxford and IBH.
- Summers, J.D.; Slinger, S.J. and Ashton, G.C. (1965).
The effect of dietary energy and protein on carcass composition with a note on a method for estimating carcass composition.
Poultry Sci. 44: 501- 509.
- Summers, J.D. and Lesson, S. (1979).
Composition of poultry meat as affected by nutritional factors.
Poultry Sci. 58: 536- 542.
- Thomas, O.P. and Combs, G.F. (1967).
Relationship between serum protein level and body composition in the chick.
J. Nutr. 91: 468- 472.
- Thompson, J.N.; and Scott, M.L. (1970).
Impaired lipid and vitamin E absorption related to a trophy of the pancereas in selenium chicks.
J. Nutre 100: 797- 809.
- Tove, S.B. (1960).
Milk lipid symposium the origin of depot fat.
J. Dariy Sci. 43: 1345- 1360.