

# CONTENTS

	Page
<b>I. INTRODUCTION .....</b>	<b>1</b>
<b>II. REVIEW OF LITERATURE .....</b>	<b>4</b>
<b>III. MATERIALS AND METHODS.....</b>	<b>36</b>
<b>IV. EXPERIMENTAL RESULTS.....</b>	<b>48</b>
1 Survey of potato viruses .....	48
2 Effect of different size of explant on growth rate and elimination of potato viruses.. 51	
2-1 Effect of size of explant on percentage of survived plantlets .....	51
2-2 Effect of size of explant on growth rate.....	54
2-3 Effect of size of explant on elimination of potato viruses .....	54
2-4 Effect of virus status of potato plantlets developed through tissue culture technique on production of microtubers .....	57
3 Effect of combination between Virazole treatment and size of explant.....	60
3-1 Effect of combination between Virazole treatment and size of explant on survivals.....	60
3-2 Effect of combination between Virazole treatment and size of explant on in vitro growth rate .....	61
3-3 Effect of Virazole on elimination of viruses .....	67
3-4 Effect of Virazole on microtubers production of virus-free and -infected plantlets in vitro .....	70
4 Effect of combination between size of explant and salicylic acid treatment .....	75
4-1 Effect of salicylic acid on survival percentage.....	75
4-2 Effect of salicylic acid on rate of growth.....	77
4-3 Effect of salicylic acid on elimination of potato viruses .....	82
4-4 Effect of salicylic acid on the in vitro potato microtubers production .....	84
5 Effect of combination between gamma radiation of infected potato tubers and size of explant .....	89
5-1 Effect of irradiation on survival percentage.....	90
5-2 Effect of irradiation on rate of growth.....	90
5-3 Effect of irradiation on elimination of potato viruses .....	96
5-4 Effect of pretreatment with irradiation on in vitro microtuber production .....	99
5-5 Effect of irradiation on protein pattern.....	103
6 Minutubers productivity of different virus-free plants under greenhouse conditions as affected by different pre-treatments .....	107
6-1 Minutubers productivity of virus-free plants of Spunta cultivar .....	108
6-2 Minutubers productivity of virus-free plants of Diamont cultivar .....	111
<b>V. DISCUSSION .....</b>	<b>115</b>
<b>VI. SUMMARY .....</b>	<b>132</b>
<b>VII. REFERENCES .....</b>	<b>143</b>
<b>ARABIC SUMMARY</b>	

<b>List of Tables</b>	<b>Page</b>
-----------------------	-------------

Table (1): Chemical ingredient of Murashige and Skoog's medium for potato meristem culture.	<b>40</b>
Table (2): Descriptive grades used for determination of the in vitro relative growth of potato explants	<b>41</b>
Table (3): Percentage of infection with PVX, PVY and PLRV and their combinations in potato cvs. Spunta and Diamont in different locations in Egypt.	<b>49</b>
Table (4): Effect of size of explant on number and percentage of survived potato plantlets cultured on MS-medium for 20 weeks.	<b>51</b>
Table (5): Effect of explant size on percentage of relative growth of in vitro potato cvs. Spunta and Diamont .in.	<b>54</b>
Table (6): Effect of explant size on number and "percentage" of virus-free or virus-infected plantlets of potato cvs. Spunta and Diamont developed from different size of explant.	<b>56</b>
Table (7): Comparison between in vitro microtuber production of the ELISA tested plantlets of potato cv. Spunta virus-free or infected with PVX, PVY and/or PLRV singly or in combination.	<b>58</b>
Table (8): Comparison between in vitro microtuber production of the ELISA tested plantlets of potato cv. Diamont virus-free or infected with PVX, PVY and/or PLRV singly or in combination.	<b>58</b>
Table (9): Effect of different concentrations of Virazole in MS-culture media on number percentage of survived plantlets developed from 0.2, 0.5 and 0.7 mm long explants for potato cultivars Spunta and Diamont.	<b>61</b>
Table (10): Effect of Virazole on the growth rate of potato cv. Spunta cultured in vitro through 20 weeks.	<b>63</b>
Table (11): Effect of Virazole on the growth and development (growth rate) of potato cv. Diamont cultured in vitro through 20 weeks.	<b>64</b>
Table (12): Effect of different concentration of Virazole on elimination of potato viruses using different size of explant in Spunta cultivar.	<b>68</b>
Table (13): Effect of different concentration of Virazole on elimination of potato viruses using different size of explant in Diamont cultivar	<b>69</b>
Table (14): Effect of different concentrations of Virazole on the in vitro microtuber production of plantlets of potato cv. Spunta either virus-free or infected with PVX, PVY and/or PLRV singly or in combination.	<b>71</b>
Table (15): Effect of different concentrations of Virazole on the in vitro microtuber production of plantlets of potato cv. Diamont virus-free or infected with PVX, PVY and/or PLRV singly or in combination.	<b>74</b>
Table (16): Effect of different concentrations of salicylic acid in MS-culture media on number percentage of survived plantlets developed from explants with different lengths (in mm) for potato cultivars Spunta and Diamont.	<b>77</b>

<b>Table (17):</b> <i>Effect of salicylic acid on the growth and development (growth rate) of potato cv. Spunta cultured in vitro through 20 weeks.</i>	<b>78</b>
<b>Table (18):</b> <i>Effect of salicylic acid on the growth and development (growth rate) of potato cv. Diamont cultured in vitro through 20 weeks.</i>	<b>79</b>
<b>Table (19):</b> <i>Effect of different concentration of salicylic acid on elimination of potato viruses using different size of explant in Spunta cultivar.</i>	<b>82</b>
<b>Table (20):</b> <i>Effect of different concentration of salicylic acid on elimination of potato viruses using different size of explant in Diamont cultivar.</i>	<b>84</b>
<b>Table (21):</b> <i>Effect of explant of different concentrations of salicylic acid on the in vitro microtuber production of plantlets of potato cvs. Spunta virus-free or infected with PVX, PVY and/or PLRV singly or in combination.</i>	<b>85</b>
<b>Table (22):</b> <i>Effect of explant of different concentrations of Salicylic acid on the in vitro microtuber production of plantlets of potato cvs. Diamont virus-free or infected with PVX, PVY and/or PLRV singly or in combination.</i>	<b>86</b>
<b>Table (23):</b> <i>Effect of irradiation and size of explant on percentage of survived plantlets for potato cultivars Spunta and Diamont.</i>	<b>89</b>
<b>Table (24):</b> <i>Effect of irradiation on the in vitro growth rate of potato cv. Spunta through 20 weeks of incubation.</i>	<b>92</b>
<b>Table (25):</b> <i>Effect of irradiation on the growth and development (growth rate) of potato cv. Diamont cultured in vitro through 20 weeks.</i>	<b>93</b>
<b>Table (26):</b> <i>Effect of different doses of gamma irradiation on elimination of potato viruses infected Spunta cultivar.</i>	<b>97</b>
<b>Table (27):</b> <i>Effect of different doses of gamma irradiation on elimination of potato viruses infected Diamont cultivar</i>	<b>98</b>
<b>Table (28):</b> <i>Effect different doses of gamma irradiation on the in vitro microtuber production of plantlets of potato cvs. Spunta virus-free or infected with PVX, PVY and/or PLRV singly or in combination.</i>	<b>100</b>
<b>Table (29):</b> <i>Effect different doses of gamma irradiation on the in vitro microtuber production of plantlets of potato cvs. Diamont virus-free or infected with PVX, PVY and/or PLRV singly or in combination.</i>	<b>101</b>
<b>Table (30):</b> <i>The effete of irradiation dosage on protein profile in the in vitro healthy and infected plantlets of Spunta and Diamont cultivars.</i>	<b>104</b>
<b>Table (31):</b> <i>Minitubers productivity under greenhouse conditions using in vitro micotubers of virus-free plantlets of Spunta cv., developed from different treatments.</i>	<b>109</b>
<b>Table (32):</b> <i>Minitubers productivity under greenhouse conditions using in vitro micotubers of virus-free plantlets of Diamont cv., developed from different treatments.</i>	<b>112</b>

<b>List of Figures</b>	<b>page</b>
<b>Fig. (1):</b> Percentage of infection with PVX, PVY and PLRV, singly or in combination in potato cultivars Spunta and Diamont during survey in Giza, Qalubia and Gharbia locations.	<b>50</b>
<b>Fig. (2):</b> Sprouted tubers and detached sprouts of two potato cultivars i.e. Spunta and Diamont.	<b>52</b>
<b>Fig. (3):</b> Excised meristem tips of potato cv., Spunta (explants) at 0.2, 0.5, and 0.7 mm long. <b>Fig. (4):</b> Effect of different size of explant on in vitro relative growth of potato plantlets of Spunta and Diamont cvs.	<b>53</b>
<b>Fig. (4):</b> Effect of different size of explant on growth rate of both Spunta and Diamont in vitro.	<b>55</b>
<b>Fig. (5):</b> Microtubers production of Spunta cv., plantlets virus-free (Free) or still infected with PVX, PVY (Y), PVX+PVY or PVY+PLRV, all developed from 0.5 mm long explant without any additional treatment.	<b>59</b>
<b>Fig. (6):</b> Effect of different concentrations of Virazole and size of explant on survival percentages of potato cvs. Spunta and Diamont.	<b>62</b>
<b>Fig. (7):</b> Effect of different concentrations of Virazole and size of explant on in vitro relative growth of potato cvs. Spunta and Diamont.	<b>65</b>
<b>Fig. (8):</b> Effect of Virazole treatment on growth rate of potato plantlets ( Spunta cv.) after 1-5 months in vitro.	<b>66</b>
<b>Fig. (9):</b> Effect of Virazole at 10 ppm (A), 20 ppm (B), and 40 ppm (C) on microtubers production of virus-free plantlets (Free) as well as those infected with PVX, PVY (Y), PLRV singly or in combination. (all plantlets developed from 0.5 mm long explant of Spunta cv.).	<b>72</b>
<b>Fig. (10):</b> Effect of different concentrations of salicylic acid and size of explant on survival percentages of potato cvs. Spunta and Diamont.	<b>76</b>
<b>Fig. (11):</b> Effect of different concentrations of salicylic acid and size of explant on in vitro relative growth of potato cvs. Spunta and Diamont.	<b>80</b>
<b>Fig. (12):</b> Effect of salicylic acid treatment on growth rate of potato plantlets (Spunta cv.) after 1-5 months in vitro.	<b>81</b>
<b>Fig. (13):</b> Effect of salicylic acid at 10 ppm (A), 20 ppm (B), and 40 ppm (C) on microtubers production of virus-free plantlets (Free) as well as those still infected with PVX, PVY (Y), PLRV singly or in combination. (all plantlets developed from 0.5 mm long explant of Spunta cv.).	<b>87</b>
<b>Fig. (14):</b> Effect of pretreatment with different doses of gamma irradiation and size of explant on survival percentages of potato cvs. Spunta and Diamont.	<b>91</b>
<b>Fig. (15):</b> Effect of pretreatment with different doses of gamma irradiation and size of explant on in vitro growth rate of potato cvs. Spunta and Diamont.	<b>94</b>
<b>Fig. (16):</b> Effect of pretreatment with different doses of gamma irradiation on growth of potato plantlets ( Spunta cv.), after 1-5 months in vitro.	<b>95</b>

**Fig. (17):** Effect of pretreatment with 15, 30, and 30 Gy of gamma irradiation on microtubers production of virus-free plantlets (Free) as well as those still infected with PVX, PVY (Y), PLRV singly or in combination. (all plantlets developed from 0.5 mm long explant of Spunta cv.). **102**

**Fig. (18):** SDS- PAGE for protein patterns in non-irradiated healthy and virus-infected plantlets of Spunta cultivar compared with those exposed to different doses of irradiation. (Molecular weight standard is indicated in KD at left side). ——— Lain M = protein Marker. ; Lain 6 = healthy Diamont cultivar. ; 7 = Infected Diamont cultivar. ; 8 = Irradiated plantlets at 15 GY. ; 9 = Irradiated plantlets at 30 GY. ; 10 = Irradiated plantlets at 45 GY. **105**

**Fig. (19):** SDS- PAGE for protein patterns in non-irradiated healthy and virus-infected plantlets of Diamont potato cv. compared with those exposed to different doses of irradiation. **106**

**Fig. (20):** Minutubers produced under greenhouse conditions from virus-free plants of potato cultivars Spunta and Diamont developed through tissue culture technique alone (control). **114**