

REFERENCES

- Abdel-Hamid, A.M.; Sadik, Fayzalla, E.A. (1985).** Preserving power of some additives against fungal invasion and mycotoxin production in stored crushed corn containing different levels of moisture. *Acta-Phytopathologica-Academia Scientarum-Hungaricae*, 20 (3/4): 309-320.
- Agrawal, A.K.; Kanjilal, S.C.; Thakre, B.D. and Bansode, P.C. (1990).** Storage of wheat seed in stone slab bin. *Seeds & Farms*, 15 (3-4): 37-38.
- Ali, H.Y. (1992).** Control of storage molds of moist corn grains during storage. *Bulletin of Fac. of Agric., Cairo Univ.*, 43 (1. Suppl.): 353-374.
- Anon, (1962).** Approved Method of American Association of Cereal Chemists. 7th Edt., The American Association, St. Paul Minn. (c.f. *Egypt. J. Food Sci.*, 10(1-2):3).
- Anon, (1975).** Official Method of Analysis. 12th Edt., by the Association of Official Analytical Chemist, Benjamin Franklin Station, Washington, DC 20044.
- Anon, (1980).** Official Method of Analysis of the Association Official Analytical Chemists. 3rd Edt., published by the Associative of Official Analytical Chemist. Box 540 Benjamin Franklin Station, Washington, Dc 20044.
- Anon, (1990).** Official Method of Analysis of the Association Official Analytical Chemists. 15th Edt., Kenneth Helrich edit., published by the Association of Official Analytical Chemist., Inc. Virginia, USA.
- Assawah, M.W. and Elarosi, H. (1960).** Fungi associated with wheat, barley and maize grains. *Egypt. J. Bot.*, 3 :153-194.

- Bangkok, B. (1986).** Aflatoxin in Thailand. Aflatoxin in maize. A proceedings of the workshop El-Batan, DF. Mexico, April 7-11. pp. 373-376.
- Botast, R.J.; Anderson, R.A.; Waraer, K. and Kwolek, W.F. (1981).** Effect of moisture and temperature on microbiological and sensory properties of wheat flour and corn meal during storage. *Cereal Chem.*, 58 : 309-311.
- Bottomley, R.A.; Christensen, C.M. and Geddes, W.F. (1952).** Grain storage studies, 10: the influence of aeration time and moisture content on fat acidity, nonreducing sugars and mold flora of stored yellow corn. *Cereal Chem.*, 29 : 53-64.
- Calvert, O.H.; Lilehoj, E.B. Kwolek, W.F. and Uber, M.S. (1978).** Aflatoxin B₁ and G₁ production in developing *Zea mays* kernels from mixed inocula of *Aspergillus flavus* and *A. parasiticus*. *Phytopathology*, 68 : 501-506.
- Cardona, T.D.; Noomhorn, A.; Ilangantilcke, S.G. and Guzaman, J.D. (1992).** Asmal scale chemical applicator to prevent aflatoxin contamination in maize cobs. *Postharvest Biology and Technology*, 1 (4): 327-342.
- Chatterjee, D.; Chattopadhyay, B.K. and Mukherje, S.K. (1990).** Storage deterioration of maize having pre-harvest infection with *Aspergillus flavus*. *Letters in Appl. Microbiol.*, 11 (1): 11-14.
- Choudhary, A.K. (1992).** Influence of microbial co-inhabitants on aflatoxin synthesis of *A. flavus* on maize kernels. *Letters in Applied Microbiology*, 14 (4): 143-147.
- Christensen C.M. (1948).** The mold flora of stored wheat and corn and its relation to heating of maize grain. *Cereal Chem.*, 25 : 4051.

- Christensen, C.M. (1957).** Deterioration of stored grains by fungi. *Botan. Rev.*, 23: 108-134.
- Christensen, C.M. (1970).** Moisture content, transfer and invasion of stored sorghum seeds by fungi. *Phytopathology*, 60 : 280-283.
- Christensen, C.M. (1972).** Storage tests with popcorn. *Plant Disease Reporter*, 56 : 1097-1099.
- Christensen, C.M. and Kaufman, H.H. (1965).** Deterioration of stored grains by fungi. *A. Rev. Phytopathology*, 3: 69-84.
- Clear, R.M. and Patrick, S.K. (1993).** Prevalence of some seed-borne fungi on soft white winter wheat seed from Ontario, Canada. *Canadian Plant Disease Survey*, 73 (2): 143-149.
- Cuero, R.G.; Duffus, E.; Osuji, G. and Pettit, R. (1991).** Aflatoxin control in preharvest maize effects of chitosan and two microbial agents. *J. of Agric. Science*, 117: 155-169.
- Dubois, M.; Cilles, K.A.; Hamilton, J.K. and Smith, F. (1956).** Colorimetric method for determination of sugars and related substances. *Analytical Chem.*, 28 (3): 350 - 356.
- Eisa (Nawal) A.; Abdel-Reheem, S.K.; Badr, A. E. and Abo-Elela, M.E. (1996).** Pathological studies on deterioration of yellow corn during storage and its control.
- El-Araby, Mervat S. (1985).** Studies on rots of *Zea maize* kernels. M. Sc. Thesis, Fac. Agric., Moshtohor, Zagazig Univ.
- El-Kady, L.A. and El-Maraghy, S.S.M. (1990).** Mycoflora and natural occurrence of mycotoxins of wheat (*Triticum vulgare* L.) grains in Egypt. *Egyptian Journal of Botany*, 33 (2): 153-167.

- Epstein, D.M.; Steiberg, P.; Nelson A.I. and Wei, L. (1970).** Aflatoxin production as affected by environmental conditions. J. of Food Sci., 35 : 289-391.
- Faraj, M.K.; Smtth, J.E. and Harran, G. (1991).** Interaction of water activity and temperature on aflatoxin production of *A. flavus* and *A. parasiticus* in irradiated maize seeds. Food Additives and Contaminants, 8 (6): 731-736.
- Gangopadhyay, S. and Wyllie, T.D. (1973).** Biochemical comparison of nine soybean varieties in relation to their susceptibility to *Macrophomina phaseolina*. Indian J. Mycol and Pl. Pathol., 3: 131-140.
- Glueck, J.A.; Roney, L.W.; Rosenow, D.T. and Miller, F.R. (1977) .** Physical and structural properties of field deteriorated (weathered) Sorghum grain. Sorghum diseases, A world review, Proceedings of the International Work Shop. International crops Research Institute for the semi-Arid tropics, India, pp. 469.
- Golubchuk, M.H.; Sorgar-Domenigg; Cuendet, L.S.; Christensen, C.M. and Geddes, W.F. (1956).** Influence of mold infestation and temperature on the deterioration of wheat during storage at approximately 12% moisture content. Cereal Chem., 33: 45-52.
- Golzar, H. (1989).** Studies on deteriorative abilities on some common storage fungi on wheat seeds. Iranian J. of Plant Pathology, 25 (1-4): 1-2.
- Grzelak, K. and Szyrmer, J. (1980).** Evaluation of the germination capacity and mycoflora of triticales seed. Hodawla Roslin Aklimatyzacjai Nasiennictwo, 24 (5):625-627.
- Gupta, R.B.L.; Majumdar, V.L. and Bhatnagar, G.C. (1990).** Influence of seed dressing fungicides on mycoflora and viability of wheat seed under storage. Seed Research, 18 (2): 157-159.

- Hansen, H.N. (1926).** A simple of obtaining single spores culture. Science, 64: 384, 1659.
- Hasan, H.A.H. (1993).** Differential action of cercoran and Topsin-M on sensitive and tolerant strains of toxigenic fungi. Cryptogamie Mycologie, 14 (1): 61-68 Botany Dept., Fac. of Science, Assiut University, Egypt.
- Hocking, A.D. and Banks, H.J. (1991).** Effects of phosphine fumigation on survival and growth of storage fungi in wheat. J.of Store Products Research, 27 (2):115-120.
- Hysek, J.; Tempirova, Z.; Vlk, J. and Curiova, S. (1986).** Mycoflora on stored cereal seed. Rostlinna Vgroba, 32 (9):963-970.
- Ilangatntilek, S.; Surapurk, P. and Escalant, F. (1987).** Frame-level chemical treatment to control aflatoxin development in crib stored maize cobs. edited by Meas, B.M.D., 1-5.
- Koksharova, T.A; Agmalova, S.R. and Nikitina, E. (1988).** Biochemical characteristics of embryotissues of wheat seeds stored for different periods. Fiziologiyi Ibiokhimiya Kul'turnykh Rastenii, 20 (3):288-291.
- Macek, J. and Celar, F. (1990).** Physiological properties of the fungus *Trichoderma longibrachiatum* Rifai and its pathogenicity for maize seedlings and young maize plants. Yugoslavia. J. of Phytopathology, 130 (3):241-248.
- Mehratore, A.B. and Dwnedi, P.K. (1980).** Fungi associated with wheat in the field. International Bioderterioration Bulletin, 16 (2):37-42.
- Meronuck, R.A. (1987).** The significance of fungi in cereal grains. Plant Disease, 71 (3):287-291.

- Moreno-Martinez, E. and Ramirez-Gonzalez, J. (1989).** Use of Fungicides in the preservation of viability of maize seed invaded by *Aspergillus glaucus*. *Anales del Instituto de Biologia, Serie Botanica*, 59 (1):55-62.
- Moreno-Martinez, E. and Vidai-Gaona, G. (1981).** Preserving the viability of stored maize seed with fungicides.. *Plant Disease*, 65 (3):260-261.
- Mostafa, M.M. (1980).** Studies on some rots of soybean plant. Ph.D. Thesis. Fac. of Agric., Al Azhar. Univ.
- New, J.H. (1988).** Studies on vacuum packing of seed. *Seed Science and Technology*, 16 (3):715-723.
- Olafson, J.H.; Christensen, C.M. and Geddes, W.F. (1954).** Grain storage studies. 15: Influence of moisture content, commercial grade, and maturity on the respiration and deterioration of corn. *Cereal Chem.*, 31:333-340.
- Osman, H.Y.A (1982).** Studies on fungi associated on the sorghum grains during storage. Ph. D. Thesis Fac. of Agric., Cairo Univ.
- Papavize, and Chirstensen, C.M. (1958).** Fungus invasion and deterioration of wheat stored at low temperatures and moisture content of 15 to 18 percent. *Cereal Chem.*, 35: 27-34.
- Paster, N.; Barkai-Golan, R. and Calderon, M. (1979).** Methyl bromide fumigation for the control of *Aspergillus* and penicillia in stored grains. *Annals of Applied Biology*, 92 (3):313 321.
- Paul, S.R.; Choudhury, A.K. and Dey, S.C. (1992).** A note on the effect of pre-sowing seed treatments with potassium salts on storage quality of wheat seeds. *J. of Potassium Research*, 8 (2):148-151.

- Penocic, and Viktorija, (1969).** The effect of *Nigrospora* (Khuskia) on the development of maize. Arh- Poljopr. Naukthe., 22 (11): 118-158 [c.f. Rev. of Pl. Path., 50 (9):526].
- Prasad, T. and Pathak, S.S. (1987).** Impact of various storage systems on biodeterioration of cereals. Indian Phytopathology, 40 (1):39-46.
- Qasem, S.A. (1959).** Studies on deterioration of stored ? by fungi. ?Dis Abstr. 20 : 1952-1953.
- Qasem, S.A. and Christensen, C.M. (1958).** Influence of moisture content, temperature, and time on the deterioration of stored corn by fungi. Phytopathology. 48 : 544-549.
- Reddy, V.K. and Reddy, S.M. (1990).** Elaboration of mycotoxins by seed-borne fungi of maize. National Academy Science Letters, 13 (1):11-13.
- Riker, A. J. and Riker, R.S. (1936) .** Introduction to Research on Plant Disease. Johns S. Swift Co. Inc. Sta. Louis, Chicago, New York. 117 pp.
- Rostom, Z.M.F. and Osman, S.A. (1994).** The temperature and moisture content of stored grain in two Saudi Arabian provinces. J. of the University of Kuwait, Sci., 21(1): 61-74. (c.f. CAB Abstr.)
- Roth, G.W. and Marshall, H.G. (1987).** Effect of Timing of nitrogen fertilization and a fungicide on stored winter wheat. Agronomy Journal, 79 (2): 197-200.
- Sahab, A.F.; Metwally, M.; Naguib, K. and El-Sayed, A. (1989).** Isolation and identification of associated fungi with wheat grains with special references to vomitoxin (DON). African J. of Agric.Sci., 16 (1-2):53-65.

- Sanderson, D.B.; Mulr, W.E.; Sinha, R.N.; Tuma, D. and Kttson, C.I. (1989).** Evaluation of a model of drying and deterioration of stored wheat at near-ambient conditions. J. of Agric. Engineering Res., 42 (3):219-233.
- Sauer, B.D. and Pomerany, Y. (1992).** Biochemical, function and nutritive changes during storage. (c.f. Sauer, B.D.; R.A. Meronuck and Christentsen, C.M. 1991). Storage of cereal grains and their products). Am. Ass. Cereal Chem. Inc. St. Paul. M. N. USA. pp.813
- Sauer, D.B and Tuite, J. (1986).** Condition that affect growth of *Aspergillus flavus* and production of aflatoxin in stored maize. Aflatoxin in Maize. A Proceedings of Workshop. El Batan, DF. Mexico, April 7- 11 P: 41-50.
- Sauer, D.B.; Storey, C.L.; Ecker, O. and Fulk, D. (1982).** Fungi in U.S. export wheat and corn. Phytopathology, 72 (11):1449-1452.
- Semeniuk, G. (1954) .**Microflora, pp.77-151. In J.A. Anderson and A.W. Alcock, (Eds.), St. Paul., Minn., USA.
- Shotwell, O.L.; Hesseltine, C.W.; Burmeister, H.R.; Kwolek, W.F.; Shanon, G.M. and Hall, H.H. (1969).** Survey of cereal grains and soybeans for the presence of aflatoxin. 10 wheat, grain sorghum and oats. Cereals Chem., 46: 446-454.
- Singh T.; Tyagi, R.P.S. and Ram, B. (1982).** Investigation of some effective fungicides for control of storage fungi. Pesticides, 16 (4):15-16.
- Singh, D.V. (1983).** Fungi associated with wheat seeds and their significance. Seed Research, 11(1):103-105.
- Singh, P.P.; Bedi, P.S. and Dheera Singh (1986).** Effect of storage microflora on the total lipid and free fatty acid contents of wheat

- Tanboon, E.K.P. (1989).** Control of aflatoxin in maize. Tropical Agriculture Research Series, 22: 49-55.
- Teich, A.H.; Sampson, D.R.; Shugar, L.; Smid, A.; Curnoe, W.E. and Kennema, C. (1987).** Yield, quality and disease response of soft white wheat cultivars to nitrogen fertilization in Ontario, Canada. Cereal-Research Communications, 15(4): 265-272.
- Toman, (1994) :** Effects of postharvest oil and fungicide application on storage fungi in corn following high- temperature grain drying. Plant Disease, 78 (1):38-43.
- Tvaruzek, L. and Frecer, R. (1993).** The development some *Fusarium* sp. in young winter wheat plants depending on temperature. Ochrana Rostlin, 29 (1):1-8.
- Warmelo, K.T.; Westhuizen, and Minne, J.A. (1968).** The production of aflatoxin in naturally infected high quality maize. Dept. of Agric. Tech. Serv. Tech. Comm No. 71.
- Welty, R.E.; Qasem, S.A. and Christensen, C.M. (1963).** Tests of corn stored four years in a commercial bin. Cereal Chemistry, 40 : 277-282.
- Wilson, D.M.; Guelaner, R.C. and Hill, R.A.(1983).** Effect of plant metabolites on the *Aspergillus flavus* group and aflatoxin production. Dikens, 13-15.
- Zuber, M.S.; Calvert, O.H.; Lillehoj, E.B. and Kwolek, W.F. (1976).** Preharvest Development of Aflatoxin B₁ in corn in the United States. Phytopathology, 66: 1120-1121.