

REFERENCES

- Abu Zeid, A. A. (1983): Production of virus-free potato plants. Ph. D. Thesis in Plant Virology. Fac. of Agric., Ain Shams Univ., PP. 98.
- Allam, E. K.; R. A. Omar and A. A. El-Amrety (1973). Comparative studies on three strains of PVX isolated from naturally infected potato in Egypt. I. Host range, Physical properties and transmission. Egypt. J. Phytopathol. Vol. 5, 19-30.
- Accatino, P. (1966). Papa corahila libre de virus mediante cultivo de meristemas. Agric. Tech. (Santiago) 26, 34-39. (c.f. Mellor and Stace-Smith, 1972).
- Amirouche, L. (1982). Factors influencing the multiplication of potato in vitro. M. Sc. Thesis, University of Aberdeen. (c.f. Miller et al., (1985).
- Anon, (1987). Technical report of Egyptian seed potato production committee in cooboration with Horticulture Research Institut. Ministry of Agriculture, A.R.E. (In Arabic).
- Appiano, A., and S. Pennazio (1972). The microscopy of potato meristem tips infected with potato virus X. J. Gen. Virol. 14, 273-276.
- Bajaj, Y. P. S., and L.A. Dionne (1966). Virus X-free roots from infected potato plants. Am. Potato J. 43: 384.
- Bang, U. (1979). Totally infected varieties freed from virus diseases by meristem culture. Vaxtskyddsrapporter 10: 1-26. (R.P.P. 59: 519, 1980).

- Beemster, A. B. R. and A. Rozendal (1972). Potato viruses: properties and symptoms. In: De Bokx, J. A. (ed.), viruses of potato and seed potato production. PP. 115-143. Wageningen Cent. Agric. Publ. Doc.
- Borissenko, S.; G. Schustar and W. Schmygla (1985). Obtaining a high percentage of explants with negative serological reactions against viruses by combining potato meristem culture with antiphytoviral chemotherapy. *Phytopath Z.*, 114 (2): 185-188.
- Cassells, A. C. (1987). In vitro induction of virus-free potatoes by chemotherapy. In Y.P.S. Bajaj (ed.) *Biotechnology in Agriculture and Forestry Vol. 3*: 40-50. Springer-Verlag Berlin Heidelberg.
- Cassels, A. C. and R. D. Long (1980): The regeneration of virus-free plants from cucumber mosaic virus and potato virus-Y infected tobacco explants cultures in the presence of virosole. *Z. Naturforsch.* 35, 350-351.
- Cassels, A. C. and R. D. Long (1982). The elimination of potato viruses X, Y, S and M in meristem and explant cultures of potato in the presence of virazol. *Potato Res.*, 25, 165-173.
- Chandra, R.; J. H. Dodds, and P. Tovar (1988). In vitro tuberization in potato (Solanum tuberosum L.). *International Association for Plant Tissue Culture. Newsletter No. 55, July 1, 1988.*
- Close, R. (1964). Some effects of other viruses and of temperature on the multiplication of potato virus X. *Ann. Appl. Biol.* 53, 151-164.

- Crowley, A. C. and J. B. Hanson (1960). The infection of apical meristems of tomato roots with tobacco mosaic virus after treatment with ethylenediamine tetraacetic acid. *Virology*, 12, 603-606.
- De Bokx, J. A. (1972). Test plants. In: Bokx, J. A., DE (Ed.): viruses of potato and seed-potato production. PP. 102-110. Wageningen. Cent. Agric. Publ. Doc.
- Dodds, J. H. (1988). Tissue culture technology practical application of sophisticated methods. *Amer-potato J.* 65: 167-180.
- Donnelly (Danielle) (1990). Tissue culture technology for crop production. Canada-Egypt McGill Agric. Response Program (CEMARP).
- El-Shehedi, A. A.; E. A. Salama, H. M. Mazyad, S. A. Eid and T. M. Wasr El-Din (1981). Effect of some chemical additives on growth of potato meristems and their freedom from PVX. *Agric. Res. Rev.* 59 (2): 241-252.
- Fernow, K. H.; L. C. Peterson and R. L. Plaisted (1962). Thermotherapy of potato leaf roll. *Am. Potato J.* 39, 446-451.
- Goodwin, P. B.; Y. C. Kim, and T. Adisarwanto (1980). Propagation of potato by shoot-tip culture. 1. Shoot multiplication. *Potato Res.*, 23: 9-18.
- Gregorini, G. and R. Lorenzo (1974). Meristem-tip culture of potato plants as a method of improving productivity. *Potato Res.* 17: 24-33.
- Hamid, A. and S. B. Locke (1961). Heat inactivation of leaf roll virus in potato tuber tissue. *Am. Potato J.*, 38, 304-310.

- Harris, R. E. and J. H. Stevenson (1982). Tissue culture of grapes. *Canada Agric.*, 27 (1): 29-33.
- Hollings, M. (1965). Disease control through virus-free stock. *Ann. Rev. Phytopathol.* 3: 367-396.
- Hollings, M., and O. M. Stone (1964). Investigation of carnation viruses. I. Carnation mottle. *Ann. Appl. Biol.*, 53: 103-118.
- Hollings, M.; and O. M. Stone (1968). Technique and problems in the production of virus-tested planting material. *Scient. Hort.*, 20: 57-72.
- Holmes, F. O. (1948). Elimination of spotted wilt from stock of dahlia. *Phytopathology* 38: 314 (Abstr.).
- Horvath, J. (1967). Separation and determination of viruses pathogenic to potatoes with special regard to potato virus Y. *Acta Phytopath.*, 2: 319-359.
- Hoyman, Wm. G. (1964). Red Pontiac vine and tuber yields as affected by virus X. *Am. Potato J.* 41: 208-211.
- Hussey, G. and N. J. Stacey (1981). In vitro propagation of potato Solanum tuberosum. *Ann. Bot.* 48: 787-796.
- Ingram, D. S. (1973). Growth of plant parasites in tissue culture. In: Street, H. E. (Ed.). *Plant tissue and cell culture*. PP. 392-421. Oxford-Blackwell Sci. Publ.
- Jayasinghe, M.; Chaquillanqui and L. F. Salazar (1989). Modified expression of virus resistance in potato in mixed virus infections. *Am. Potato J.*, 66: 137-144.

- Jordan, M.; G. Apablaza; and P. Lippi (1978). Obtaining potato plants free of viruses X and Y by the in vitro culture of shoot apices and serological detection by the ELISA test. *Ciencia e Investigacion Agraria*, 5 (4): 207-211.
- Kapitsa, O. S. and E. N. Andrejeva (1965). Disinfection of vegetatively produced plants from virus diseases (in Russian). *Turkey Inst. Genet.*, 35, PP. 18-35. (c.f. Ouf. (1969).
- Kassanis, B. (1950). Heat inactivation of leaf roll virus in potato tubers. *Ann. App. Biol.*, 37: 339-341.
- Kassanis, B. (1954). Heat-therapy of virus infected plants. *Ann. App. Biol.*, 41: 470-474.
- Kassanis, B. (1957). The use of tissue culture to produce virus free clones from infected potato varieties. *Ann. Appl. Biol.*, 45 (3), 422-427.
- Kassanis, B. (1967). Plant tissue culture PP. 537-566. In "Methods in virology", Vol. 1 Maramorasch and Koprowski, eds., Academic Press New York and London.
- Kassanis, B. and T. W. Tinsley (1957). The freeing of tobacco tissue culture from potato virus Y by 2-thiouracil. *Proc. of the 3rd Conf. on potato virus diseases*. Liss. Wageningen, 153-155.
- Kassanis, B.; and W. W. Schwabe (1961). The effect of paracrinkle virus on the growth of King Edward potato at different temperature and day length. *Ann. Appl. Biol.*, 49: 616-620.

Kassanis, B. and A. Varma (1967). The production of virus-free clones of some British potato varieties. *Ann. Appl. Biol.*, 59, 447-450.

Klein, R. E. and C. H. Livingston (1982). Eradication of potato virus X from potato by ribavirin treatment of cultured potato shoot tips. *Am. Potato J.*, 59, 359-365.

Klein, R. E., and C. H. Livingston (1983). Eradication of potato virus X and S from potato shoot-tip cultures with ribaverin. *Phytopathology*, 73: 1049-1050.

Krylova, N. V., and V. I. Stepanenko (1971). Do viruses penetrate the apical meristem of plant. *Tr. Biol. Poch. Inst. SSSR*, 4, 1225-226 (R.P.P. 52: 309, 1972).

Krylova N. V.; I. Stepanenko and V. G. Reifman (1973). Potato virus X in potato apical meristems. *Acta. Virol.* 17, 172.

Lecklerc, Y. (1989). Acclimatization, field performance and micro tubernation of tissue cultured potato (Solanum tuberosum). M. Sc. Thesis. MacDonald College of McGill Univ. Canada.

Limassat, P.; and P. Cornuet (1949). Recherch du virus de la mosaïqua du tabaccolans 125 meristems des plantes infectees. *c.r. hebd. Seanc. Acad. Sci. Paris*, 228, 1971. (c.f. Kassanis 1957).

Lindsay, G. C. (1987). In vitro improvement potatoes: The New Zealand approach. In Y.P.S. Bajaj (ed.) *Bio-technology in Agriculture and Forestry*. Vol. 3 Potato. P. 52-61 Springer-Verlag Berlin Heidelberg.

- Lizarraga, R., V. Espinosa, U. Jayasinghe and J. H. Dodds (1989). Tissue culture for elimination of pathogens CIP. Research Guide 3. International potato center, Lima, Peru. 21 PP.
- Lozoya, Saldano, H. and O. Marlin-Lora (1984). Thermotherapy and tissue culture for elimination of potato virus X (PVX) in Mexican potato cultivars resistant to late blight. Am. Potato J., 61: 735-739.
- Lozoya-Saldana, H.; and W. D. Dawson (1982). Constant and alternating temperature regimes and tissue culture to obtain PVS-free potato plants. Am. Potato J. 59: 221-230.
- MacDonald, D. N., (1973). Heat treatment and meristem culture as means of freeing potato varieties from viruses X and S. Potato Res. 16: 263-269.
- Manzer, F. E. (1959). Potato meristem culture and virus X. Am. Potato J., 36, 191-195.
- Mauk, S. C., and A. R. Langille (1978). Physiology of tuberization in Solanum tuberosum L. cis-Zeatin riboside in the potato plants, its identification and changes in endogenous levels as influenced by temperature and photoperiod. Plant Physiol. 62: 438-442.
- Mellor, F. C. and R. Stace-Smith (1967). Eradication of potato virus X by thermotherapy. Phytopathology 57: 674-673.
- Mellor, E. C. and R. Stace-Smith (1969). Development of excised potato buds in nutrient culture. Can. J. Bot., 47, 1617.

- Mellor, F. C. and R. Stace-Smith (1970). Virus strain differences in eradication of potato viruses X and S. *Phytopathology*, Vol. 60, 1587-1590.
- Mellor, F. C., and R. Stace-Smith (1971). Heat-stable strains of potato leaf roll virus. *Phytopathology*, 61: 246-247.
- Mellor, F. C., and R. Stace Smith (1977). Virus-free potatoes by tissue culture in J. Reinert and Y.P.S. Bajaj. (ed.), *Applied and fundamental aspects of plant cell tissue and organ culture*. Springer-Verlag, Berlin Heidelberg, New York. PP. 616-635.
- Mellor, F. C. and R. Stace-Smith (1987). Virus-free potatoes through meristem culture. In Y.P.S. Bajaj (ed.) *Biotechnology in Agriculture and Forestry*, Vol. 3:30-39.
- Miller, P. R.; L. Amirouche; T. Stachbury and S. Matthews (1985). The use of plant growth regulators in micro-propagation of slow-growing potato cultivars. *Potato Res.* 23: 479-486.
- Morel, G. (1967). International symposium plant pathology. New Delhi PP. 100-101. Indian Phytopathology Soc. Publ. (c.f. Smith, 1974).
- Morel, G., and C. Martin (1952). Guérison de dahlias atteints d'une maladie A virus. *c.r. hebdomadaire des séances Acad. Sci., Paris* 235: 1324-1325.
- Morel, G. and C. Martin (1955). Cure of potatoes infected by virus diseases. *c.r. hebdomadaire Acad. Agric. Fr.* 41: 472-475. (*R.A.M.* 53: 215, 1956).
- Morel, G.; C. Martin and J. F. Muller (1968). La guérison des pommes de terre atteintes de maladies à virus. *Ann. Physiol. Vegetale*. 10, 113-139.

- Morel, G., and J. F. Muller (1964). La culture in vitro du meristem apical de la pomme de terre. c.v. hedd Scanc. Acad. Sci. Paris 258: 5250-5252. (c.f. Puak, 1972).
- Mori, K. (1971). Production of virus free plants by means of meristem culture. JARQ Vol. 6 (1), 1-7.
- Mori, K. (1977). Localisation of viruses in apical meristems and production of virus-free plants by means of meristems and tissue culture. Acta Horticulturae, 78, 389-396.
- Munro, J. (1961). The importance of potato virus X. Am. Potato J., 38, 440-447.
- Murashige, T., and F. Skoog (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiologia Plantarum. 15: 473-397.
- Norris, D. O. (1953). The effect of virus X on yield of potatoes. J. Aust. Inst. Agric. Sci. 19, 251-256.
- Norris, D. O. (1954). Development of virus-free stock of Green-Mountain potato by treatment with Malachite green. Aust. J. Agric., 5, 658-663.
- Novak, F. J. and J. Zadina (1987). In vitro propagation of potato progress in Czechoslovakia. In Biotechnology in Agriculture and forestry. Vol. 3 potato (ed. by Y.P.S. Bajaji, (1987). Springer-Verlag, Berlin Heidelberg PP. 23-29.
- Nowak, J., and D. Colborne (1989). In vitro tuberization and tuber proteins as indicators of heat stress tolerance in potato. Am. Potato J., 66: 35-45.

- Nyland, G. and A. C. Goheen (1969). Heat therapy of virus diseases of perennial plants. *Ann. Rev. Phytopathol.*, 7: 331-354.
- Okasawa, Y.; N. Katsura and T. Tagawa (1967). Effect of auxin and kinetin on the development and differentiation of potato tissue cultured in vitro *Physiol. Plant.* 20: 826-829.
- Omar, R. A.; N. Shaheen; M. A. Abdel-Hadi and M. M. Abu El-Naga (1983-a): Potato viruses in Egypt and factors affecting their spread within potato fields. *Egypt. Soc. Appl. Microbiol. Proc. V. Microbiol.*, Cairo, May 1983. Vol. III Plant Pathology, PP. 404-433.
- Omar, R. A.; F. Fadel; M. A. Abdel-Hadi and M. M. Abu El-Naga (1983-b). The influence of viruses on potato yield in Egypt and some measures of control. *Egypt. Appl. Microbiol. Proc. V. Conf. Microbiol.*, Cairo, May 1983, Vol. III Plant Pathology, 435-463.
- Oshima, N. and C. H. Livingston (1961). The effects of anti-viral chemicals on potato virus X. *Am. Potato J.* 38, 294-299.
- Oshima, N. and C. H. Livingston (1963). In vitro culture of potato tissue, *Am. Potato J.*, 40: 9-16.
- Ouf, M. F. (1964). Comparative studies on some methods for obtaining virus-free seed potatoes (in Russian). *Avtoreferat Kand. diss.*, TSXA; Moskva. (c.f. Ouf, 1969).
- Ouf, M. F. (1969). The use of apical meristematic tissue cultures of potato etiolated sprouts to produce virus X-free clones from infected varieties. *Bull. Sci. and Technol. (Asyut Univ.)* Vol. 12 (1) 229-247.

- Palmer, C. E. and O. E. Smith (1969). Cytokinin and tuber initiation in potato solanum tuberosum. Nature 221-280.
- Palmer, C. E., and O. E. Smith (1970). Effect of kinetin on tuber formation on isolated stolons of Solanum tuberosum L. cultured in vitro. Plant Cell Physiol. 11, 303-314.
- Palmer, C. E.; and W. G. Barker (1972). Changes in enzyme activity during elongation and tuberization of stolons of Solanum tuberosum L. cultured in vitro. Plant and Cell Physiology 13: 681-688.
- Pennazio, S. (1971). Treatment of virus disease of potato Solanum tuberosum L., Meristem tip culture combined with heat treatment. Ottoflorofruttic. Ital. 55: 446-452 (R.P.P. 52: 213, 1973).
- Pennazio, S. (1973). Effects of four antimetabolites on PVK inhibition in infected potato tips cultured on artificial substrate. Rivista di Patologia vegetale, IV 9 (1): 3-10.
- Pennazio, J. and P. Redalfi (1974). Potato virus X eradication in cultured potato meristem tips. Potato Res., 17, 333-335.
- Pennazio, S.; and M. Vacchiati (1978). Potato virus X eradication from potato meristem tips held at 30°C. Potato Res. 21: 19-22.
- Phillips, D. J. (1968). Carnation shoot-tip culture. Tech. Bull. 102, Colorado State Univ. Exp. Station.
- Quak, (Fredrerikea) (1961): Heat treatment and substances inhibiting virus multiplication in meristem culture to obtain virus-free plants. In "Advances in Hort, Sci. and their applications". Vol. 1.

- Quak, (Fredrika), (1972): Therapy culture, PP. 158-166. In "Viruses of potatoes and seed potato production". (J.A. de Bokx, ed.). Wageningen Center for Agric. Publishing Documentation.
- Quak, F. (1977). Meristem culture and virus-free plants. Plant cell tissue and organ culture (Ed. by Reinert and V.F.S. Bajaj,) PP. 598-615. Springer Verlag, Berlin.
- Rawlins, T. E. and C. M. Tompkins (1936). Studies on the effect of carborandum as abrassive in plant virus inoculation. *Phytopathology*, 26: 578.
- Roberts, A. A.; F. M. Boldgett and R. E. Wilkinson (1952). Potato virus X: Inoculation of potato varieties tolerant to virus Y. *Am. Potato J.* 29: 212-220.
- Rosell, G.; F. G. De Bertoldi and R. Tizio (1987). In vitro mass tuberization as a contribution to potato micro-propagation. *Potato Res.*, 30: 111-116.
- Rozenberg, V. (1972). Freeing regional potato varieties from S, X, M and Y by meristem culture in the Estanion SSSR, In *Kratk. Doki povopt Zashch. Rest. VIII. Pribalt knof. pozashch. Rast. ch. I Kaunsas 1972*, 106-109 (Ru) from *Referatiunyi Zhurnal* (1972) 10: 55-84-84 (R.P.P. 52: 340, 1973).
- Sheffield, F. M. L. (1942). Presence of virus in primordial meristem. *Ann Appl. Biol.* 29: 16-17.
- Shepard, J. F. and L. Claflin (1975). Critical analysis of the principles of seed potato certification. *Ann. Rev. Phytopathol.*, 13: 271-293.

- Sip, V. (1972). Eradication of potato viruses A and S by thermotherapy and sprout tip culture. *Potato Res.*, 15, 270-273.
- Smith, K. M. (1974). *Plant viruses*. Chapman and Hall, London, 211, P.
- Stace-Smith, R. (1985). Virus-free clones through plant tissue culture. *Comprehensive Biotechnology Vol. 3, Chapter 10*, PP. 29.
- Stace-Smith, R., and F. C. Mellor (1968). Eradication of potato virus X and S by thermotherapy and axillary bud culture. *Phytopathology* 58: 199-203.
- Stace-Smith, R. and F. C. Mellor (1970). Eradication of potato spindle tuber virus by thermotherapy and axillary bud culture. *Phytopathology*, 60: 1857-1858.
- Stalknecht, F. G. and S. Farnsworth (1982). General characteristics coumarin-induced tuberization of axillary shoots of *Solanum tuberosum* cultured in vitro. *Am. Potato J.* 59: 17-32.
- Svobodova, J. (1964). Spread of Y-virus through tobacco callus tissue into shoot developed from those tissues. PP. 65-67. In: "Host-parasite Relations in plant pathology" Ministry of Agric. the Hungarian Academy of Science.
- Sward, R. J. and N. D. Hallam (1976). Changes in fine structure of the potato meristem following heat treatment for virus eradication. *Aust. J. Pot.* 24, 597-605.
- Tao, G.; Y. Wei; G. Goupu and C. Cui (1987). In vitro production and release of potato varieties in China. In Y.P.S. Bajaj (ed.). *Biotechnology in Agriculture and Forestry*, Vol. 3: 62-79.

- Tapio, E. (1972). Virus-free clones of the potato varieties Pito and Tammiston Aikainen. *Ann. Agric. Fenn.* 11: 115-118.
- Thieme, R. and B. Patt (1982). Production of tubers in vitro and their use in the establishment of a potato store. *Arch. Zuechtungsforsch.* 12, 257-262. (c.f. Wang and Hu. 1985).
- Thomson, A. D. (1956). Studies on the effect of malachite green on potato viruses X and Y. *Aust. J. Agr. Res.*, 7: 428-434.
- Thomson, A. D. (1958): The elimination of viruses from potato tissue. *Proc. 3rd Conf. Potato virus diseases.* Lisse-Wageningen, 156-159.
- Thriumalacher, M. J. (1954). Inactivation of potato leaf roll by high temperature storage of seed tubers in Indian plains. *Phytopathol. Z.*, 22: 429-436.
- Vasti, S. M. (1973). Effect of antiviral chemicals on production of virus X-free potato tubers. *Pakistan J. of Botany*, 5 (2): 139-142.
- Walkey, D. G. A. (1976). High temperature inactivation of cucumber and alfalfa mosaic viruses in Nicotiana rustica cultures. *Ann. Appl. Biol.*, 84: 183-192.
- Walkey, D. G. A. (1980). Production of virus free-plants by tissue culture. In tissue culture methods for plant pathologists by Ingram, D. S. and J. P. Helgeson, P. 109-117.

- Wang, P. J. and C. Y. Hu (1982). In vitro mass tuberization and virus free potato production in Taiwan. Am. potato J. 59: 33-37.
- Wang, P. J. and C. Hu (1985). Potato tissue culture and its application in agriculture. In Potato Physiology. Edited by P.H.Li. Academic Press Orlando, P.P. 503-577.
- Wang, P. J. and L. C. Huang (1975). Callus culture from potato tissue and the exclusion of potato virus X from plants regenerated from stem tips. Can. J. Bot. 53: 2565-2567.
- Wattimena, G. B. (1983). Micropropagation as an alternative technology for potato production in Indonesia. Dissertation Abstract 44: 07: 2040.
- White, P. R. (1934). Multiplication of viruses of tobacco and acuba mosaic in growing excised tomato root tips. Phytopathology. 24: 100-101.
- Wright, N. S. (1970). Combined effects of potato virus X and S on yield of Netted Gem and White Rose potatoes. Am. potato J. 47, 475-478.
- Wright, N. S. (1977). The effect of separate infections by potato virus X and S on Netted Gem potato. Am. potato J. 54:147-149.
- Wright, N. S. (1983). Uniformity among virus-free clones of ten potato cultivars. Am. Potato J. 40: 381-388.
- Yora, K.; and Tsuchizaki (1962): The elimination of latent viruses from potato of Danskaku variety by use of tissue culture. Ann. Phytopathol. Soc., Japan 27: 219-221. (R.P.P. 42: 1213-1963).

Zhuk, I. R. (1978). Culture of apical meristem as a method of elimination of virus disease of potato Sel. Biologia. 13: 512-516. (R.P.P. 58: 166, 1979).
