

REFERENCES

Abdel-Sabour, M. S.; Bekhit, M. M. M.; Hassan, A. M. and Sherif, H. S. (1993):

Genetic studies in hexaploid wheat, diploid rye and some promising BC5 of wheat X rye. 2. Yield components.

Egypt. J. Genet. Cytol., 22 (1): 57-71.

Afanas'eva, N.I. (1985):

Inheritance of plant height in hybrids from crosses of spring wheat with rye. Genetice Selektzii. 73(3):57-59.

Attia, T. and Lelley. T. (1986):

production of 6 x triticales containing wild rye species. Egyptian J. Genet. Cytol. 15(2):285-296.

Baum, M. (1991)

Rye-wheat hybrids: The production of wheat chromosome additions to rye. Genome 34: 840 - 844.

Beatty, K, D., Rupert, E.A. and Nancy Dehgan (1976):

Doubling chromosome number of wheat-rye amphiploids with colchicine, DMSO, and cold treatment. Wheat Information Service, Dec. 1976, 43:10 - 12.

Bekhit, M.M.M. (1990):

Genetical and cytological studies on some triticales hybrids. PH.D. Thesis, Fac. of Agric. Moshtohor, Zagazig univ.

Bekhit, M. M. M.; Abdel-Sabour, M. S.; Sherif, H. S. and Hassan, A. M. (1993):

Genetic studies in hexaploid wheat, diploid rye and some promising BC5 of wheat X rye. 2. Yield

components. Egypt. J. Genet. Cytol., 22 (1): 51-56.

Cakmak, I.; Cakmak, O.; Eker, S.; Ozdemir, A.; Watanabe, N. and Braun, H. J. (1999a):

Expression of high zinc efficiency of *Aegilops tauschii* and *Triticum monococcum* in synthetic hexaploid wheats. Recent progress in plant nutrition. Contributions from an International Colloquium on plant nutrition, Stuttgart, Germany, 13-14 February 1998. Plant-and-Soil, 215: 2, 203-209.

Cakmak, I.; Kalayci, M.; Ekiz, H.; Braun, H. J.; Kilinc, Y.; Yilmaz, A.; Welch, R. M. (ed.); Graham, R. D. (1999b):

Zinc deficiency as a practical problem in plant and human nutrition in Turkey: a NATO-science for stability project. Special issue. Sustainable field crop systems for enhancing human health: agricultural approaches to balanced micronutrient nutrition. Field Crops Research., 60: 1-2, 175-188.

Cakmak, I.; Tolay, I.; Ozdemir, A.; Ozkan, H.; Ozturk, L. and Kling, C. I. (1999c):

Differences in zinc efficiency among and within diploid, tetraploid and hexaploid wheats. Annals of Botany. 1999, 84: 2, 163-171.

Carrillo, M.; Monteagudo, A. and Sanchez-Monge, E. (1983):

Inheritance of yield components and their relationship to plant height in hexaploid triticales. Z. Pflanzenziiditg 90: 153 - 165

Deo, R.; Mathur, G. M.; Deo-R; Gissel, N. G. (ed.) and Jensen, A. (1999):

Zinc status of arid soils, its response and uptake by wheat. Plant nutrition - molecular biology and genetics. Proceedings of the Sixth International Symposium on Genetics and Molecular Biology of Plant Nutrition, Elsinore, Denmark, 17-21 August 1998., 193-197.

Erenoglu, B.; Cakmak, I.; Romheld, V.; Derici, R. and Rengel, Z. (1999):

Uptake of zinc by rye, bread wheat and durum wheat cultivars differing in zinc efficiency. Plant and Soil, 209: 2, 245-252.

Fageria, N. K. (1990):

Iron requirement of cereals and legumes in solution culture. Plant nutrition-physiology and applications., 213-217.

Friebe, B. and Larter. E. N. (1988):

Identification of a complete set of isogenic wheat/rye D-genome substitution lines by means of Giemsa C-banding. Theor. of Appl. Genet. 76:473-479.

Genc, Y.; McDonald, G. K.; Rengel, Z. and Graham, R. D. (1999):

Genotypic variation in the response of barley to zinc deficiency. Plant nutrition - molecular biology and genetics., 205-221.

Gerlach, W. L. (1977):

N-banded karyotypes of wheat species. Chromosoma, 62 : 49-56.

- Gill, B. S.; Friebe, B. and Endo, T. R. (1991):**
Standard karyotype and nomenclature system for description of chromosome bands and structural aberrations in wheat (*Triticum aestivum*). *Genome*, 34 (5): 830-839.
- Giraldez, R.; Cermeno, M. C. and Orellana, J. (1979):**
C-banding pattern in the chromosome of inbred lines and open pollinating varieties of rye. *Z. pflanzenzüchtg.* 83:40-48.
- Graham, R. D. (1981):**
Genetics of the copper efficiency factor in rye. In: J. F. Loneragan, A. P. Robson, and R. D. Graham (eds.), *Copper in Soils and plants*, Proc. Golden jubilee. Int. Symp. On "Copper in soils and plants", Murdoch University, Perth, 1981, 358. Academic Press, Sydney, New York, London, Toronto, San Francisco.
- Graham, R. D. (1984):**
Breeding for nutritional characteristics in cereals. *Adv., Plant Nutr.*, 1: 57 – 102.
- Graham, R. D.; Ascher, J. S. and Hynes, S. C. C. (1993):**
Selecting zinc-efficient cereal genotypes for soil of low zinc status. Genetic aspects of plant mineral nutrition., 349-358.
- Hafiz, H. M. I. and Larik, A. S. (1982).**
Frequency of aneuploids in euploid progenies of hexaploid triticales. wheat information service 55:13-14.
- Hansson, M.; Gough, S. P.; Kannangara, C. G. and Wettstein, D. (1998):**
Chromosomal locations of six barley genes encoding enzymes of chlorophyll and heme biosynthesis and the sequence of the

ferrochelatase gene identify two regulatory genes.
Plant Phys. And Bioch. Paris., 36: 8, 545-554.

Ismail, M.; Sinha, R. B. and Sakal, R. (1995):
Screening of wheat varieties for their differential response to zinc application in calcareous soil. Journal of the Indian Society of Soil Science, 43: 3, 481-483.

Jamjod, S.; Mann, C. E. and Rerkasem, B. (1993):
Combining ability of the response to boron deficiency in wheat. Genetic aspects of plant mineral nutrition., 359-361.

Kalayci, M.; Torun, B.; Eker, S.; Aydin, M.; Ozturk, L. and Cakmak, I. (1999):
Grain yield, zinc efficiency and zinc concentration of wheat cultivars grown in a zinc-deficient calcareous soil in field and greenhouse. Field Crops Research, 63: 1, 87-98.

Khabaz, S. H.; Barker, S. J.; Graham, R. D.; Rathjen, A. J.; Gissel, N. G. (ed.) and Jensen, A. (1999):
The application of amplified fragment length polymorphism (AFLP) for breeding Mn efficiency in durum wheat (*Triticum turgidum* L. var. durum). Plant nutrition - molecular biology and genetics. Proceedings of the Sixth International Symposium on Genetics and Molecular Biology of Plant Nutrition, Elsinore, Denmark, 17-21 August 1998., 409-416.

Kynast, R. G.; Röder, M. S. and Römheld, V. (1994):
Physical mapping of micronutritional genes in wheat-rye translocations. Submitted to 2nd Triticeae Symp., Logon.

- Lelley, T.; Josifek, K. and Kaltsikes, P. J. (1978):**
Polymorphism in the Giemsa C-banding pattern of rye chromosomes. *Can. J. Genet. cytol.* 20:307-312
- Lukaszewski, A., J., Apoliarska, B. and Gustafsan, J.P. (1987):**
Introduction of the D-genome chromosomes from bread wheat into hexaploid tritiale with a complete rye genome. *Genome*, 29 : 425-430. *Can. J. Genet. Cytol.* 24:83-92.
- Lukaszewski, A. J. and Gustafson, J. P (1983).**
Translocation and modifications of chromosomes in tritcale x wheat hybrids. *Theor. Appl. Genet.* 64: 239-248.
- May, E. Cedric (1983).**
Triticale x wheat hybrids and the interaction of specklet leaf blotch resistance to wheat. *Proc. 6th Inter. wheat Genet. symp., Kyoto. Japan.* PP.175-179.
- McGrath, S. P.; Chaudri, A. M.; Thacker, D.; Salmon, S. E.; Little, K.; Douglas, S. and Cauvain, S. P. (1995):**
Chemical composition of wheat grain: I. Survey results; II. Effects of copper amendments on bread quality. *HGCA-Project-Report, No. 102*, 31 pp..
- Merry, L. J.; Graham, R. D. and Schachtman, D. P. (1999):**
Triticum tauschii – A potential source of genes for the improvement of zinc efficiency in bread wheat. *Plant nutrition – Molecular Biology and Genetics.*: 235-244.
- Moudry, J. and Dvoracek, V. (1999):**
Chemical composition of grain of different spelt (*Triticum spelta* L.) varieties.

Chemicke slozeni zrna ruznych odrud psenice spaldy (*Triticum spelta* L.). Rostlinna-Vyroba., 45: 12, 533-538.

Mujeeb-Kazi, A. and Kimber, G. (1985):

The production, Cytology and Practicality of Wide hybrids in the triticeae. Cereal Research communications, 13 (2-3): 111 – 124.

Naranjo, T. and Lacadena, J. R. (1982):

C-banding pattern and meiotic pairing in five rye chromosome of hexaploid triticales. Theor. Appl. Genet. 61:233-237.

Oettler, G. (1983):

Crossability and embryo development in wheat rye hybrids. Euphytica 32:593-600.

Orellana, T., Cermeno, Mc & hacadena. JR. (1983):

Meiotic pairing in wheat-rye addition and substitution lines. Can. J. Genet. cyto. 26:25-33.

Papa, C., Morris. R. M. and Schmidt, J. W. (1990):

Rye C-banding patterns and meiotic stability of hexaploid triticales (x. *tritico secale*) selections differing in Kernel shriveling. Genome 33:686-689.

Pearson, J. N. and Rengel, Z. (1997):

Genotypic differences in the production and partitioning of carbohydrates between roots and shoots of wheat grown under zinc or manganese deficiency. Annals-of-Botany., 80: 6, 803-808.

Podlesak, W.; Wener, T.; Grun, M.; Schlegel, R. and Hugenhof, E. (1990):

Genetic differences in the copper efficiency of cereals. Plant nutrition – Physiology and applications., 297-301.

Rengel, Z. and Graham, R. D. (1995a):

Importance of seed Zn content for wheat grown on Zn-deficient soil. I. Vegetative growth. Plant-and-Soil, 173: 2, 259-266.

Rengel, Z. and Graham, R. D. (1995b):

Importance of seed Zn content for wheat growth on Zn-deficient soil. II. Grain yield. Plant-and-Soil, 173: 2, 267-274.

Rengel, Z. and Graham, R. D. (1995c):

Wheat genotypes differ in Zn efficiency when grown in chelate-buffered nutrient solution. I. Growth. Plant and Soil, 176: 2, 307-316.

Rengel, Z. and Graham, R. D. (1995d):

Wheat genotypes differ in Zn efficiency when grown in chelate-buffered nutrient solution. II. Nutrient uptake. Plant and Soil, 176: 2, 317-324.

Rengel, Z.; Graham, R. D. and Pedler, J. F. (1994):

Time-course of biosynthesis of phenolics and lignin in roots of wheat genotypes differing in manganese efficiency and resistance to take-all fungus. Annals-of-Botany, 74: 5, 471-477.

Saberi, H. K.; Graham, R. D. and Rathjen, A. J. (1999):

Inheritance of manganese efficiency in durum wheat. Journal of Plant Nutrition, 22: 1, 11-21.

Saberi, K. H.; Graham, R. D.; Ascher, J. S. and Rathjen, A. J. (2000):

Quantification of the confounding effect of seed manganese content in screening for manganese efficiency in durum wheat (*Triticum turgidum* L. var. durum).

Journal of Plant Nutrition, 23: 7, 855-866.

Seal, A. G. (1982):

C-banded wheat chromosomes in wheat and triticale.

Theor. Appl. Genet. 63: 39-47.

Seal, A. C. and bennett, M. D. (1981):

The rye genome in winter hexaploid triticale.

Can.J.Cenet.Cyto.23.647.653.

Schlegel, R.; Kynast, R.; Schwarzacher, T.; Romheld, V. and Walter, A. (1993):

Mapping of genes for copper efficiency in rye and the relationship between copper and iron efficiency. Plant and Soil, 154: 61-65.

Schlegel, R.; Cakmak, I. and Atanasova, M. (1999):

The relationship between straw length, root dry weight and zinc efficiency as well as shoot zinc content in hexaploid wheat, *Triticum aestivum* L., Plant and Soil, submitted.

Schlegel, R.; Cakmak, I.; Ceoloni, C. (ed.) and Worland, A. J. (1998a):

Zinc efficiency in nulli-tetrasomics of hexaploid wheat 'Chinese Spring'. Proceedings of the 10th EWAC meeting, Viterbo, Italy, 16-19 June 1997. EWAC-Newsletter, 22-25.

Schlegel, R.; Ozdemir, I.; Cakmak, H.; Saberi, H. and Atanasova, M. (1998b):

Localization of genes for zinc and manganese efficiency in wheat and rye. Plant and Soil, submitted.

- Schlegel, R.; Werner, T. and Jacob, F. (1991a):**
Mineral nutrition and genetical control in cereals.
Proc. EUCARPIA Symp. Cereal Section,
Schwerin (Germany), Vortr. F. Pflanzen-
zuechtung 20, 79-84.
- Schlegel, R.; Werner, T. and Hugenhof, F. (1991b):**
Confirmation of a 4 BL/5RL, wheat-rye
chromosome translocation line in the wheat
cultivar "Viking" showing high copper efficiency.
Plant breeding, 107: 226-234.
- Shang, X. M.; Jackson, R. C. and Nguyen, H. T. (1988):**
A new technique for chromosomes of wheat
(*triticum*) and its relatives. Cereal Research and
communications 16 (3-4):169-174.
- Singh, R. J. and Robbelen, G. (1975):**
Giemsa banding technique reveals deletions
within rye chromosomes in addition lines. Z.
Pflanzenzuecht. 76: 11-18.
- Snowden, K. C.; Richards, K. D. and Gardner, R. C.
(1995):**
Aluminum-induced genes. Induction by toxic
metals, low calcium, and wounding and pattern of
expression in root tips. Plant-Physiology, 107: 2,
341-348.
- Surrikov, I. M. and Kissel, N. I. (1985):**
Crossability of bread wheat with diploid and
traploid hybrids. Theor. Appl. Genet. 71 (3):467-
471.
- Taylor, G. J. and Stadt, K. J. (1990):**

Interactive effects of cadmium, copper, manganese, nickel and Zinc on root growth of wheat (*Triticum aestivum*) in solution culture. Plant nutrition-Physiology and applications., 317-322.

Zeller, F. J. and Hsamsai, L. K. (1983):

Broadening the genetic variability of cultivated wheat by utilizing rye chromatin PROC.6th inter.wheat Genet .sympo.Kyoto,Japan,116-173.
Theor. Of Appl. Genet. 76: 473 - 479.