

REFERENCES

- Abd EL-Mogeid, S.A. (1995). Inheritance of yield, yield compnents and some morphological characteristics in spring wheat crosses. Ph.D. Thesis, Fac. Agric., El-Minia Univ., Egypt.
- Abd EL-Rahman, M.F.S. (1991). Gene action in the inheritance of yield and its components in wheat. M.Sc. Thesis, Fac. Agric., Ain Shams Univ., Egypt.
- Abdel Hak T. M., N. A. El-Sherif, A. A. Bassiouny, I. I. Shafik and Y. El-Dauadi, (1980). Control of wheat leaf rust by systemic fungicides. Proceeding of the Fifth European and Mediterranean Cereal Rusts Conference, Bari, Italy, 255-266.
- Abdel-Aty, M. S. M. and Y. S. Katta, (2000). Genetic analysis and heterosis of grain yield and related traits in bread wheat (*T. aestivum* L.). J. Agric. Res. Tanta Univ., 28(2):287-300.
- Abul-Nass, A.A., A.A. EL-Hosary and M.M. Saker. (1986). Genetical studies in common wheat (*Triticum aestivum* L.). Egypt J. Agron. 11: 93-102.
- Afiah, S. A. N., (1999). Combining ability, association and path coefficient analysis of some wheat (*T. aestivum* L.) diallel crosses under desert conditions. J. Agric. Sci. Mansoura Univ., 24(4): 1583-1596.
- Ageez, A. A. and G. A. R. El-Sherbeny, (1998). Heterosis in relation to additive genetic variance for yield and its components in bread wheat (*T. aestivum* L.). J. Agric. Sci. Mansoura Univ., 23(12): 5287-5295.

- Aglan, M. A. A., (2003). Breeding studies on wheat resistance to leaf and stem rusts and its association with yield and its components. M.Sc. Thesis Fac. Of Agric. Minufiya Univ., Egypt.
- Agrama, H.A and M.E. Moussa. (1996). Mapping QTLs in breeding for drought tolerance in Maize (*Zea Maize* L.). *Euphytica* 91, 89-97.
- Agrama, H.A and M.R. Tuinstra. (2003). Phylogenetic diversity and relationships among sorghum accessions using SSRs and RAPDs. *Afric J Biotech* 2: 334-340.
- Akanda, S. I., and C. C. Mundt, (1996). Path coefficient analysis of the effects of stripe rust and cultivar mixtures on yield and yield components of winter wheat. *Theor. App. Genet.* 92:666-672.
- Amin M. R.; M. M. Hoque, M. A. Shaheed, A. K. D. Sarker, and Z. Kabir, (1990). Genetic variability character association and path analysis in wheat [in Bangladesh]. *Bangladesh J. of Agric. Res. (Bangladesh)*. v.15 (2): 1-5.
- Anderson, J. A., Y. Ogihara, M. E. Sorreles and S. D. Tanksely, (1992). Development of a chromosomal arm map for wheat based on RFLP markers. *Theor. Appl. Genet.* 83(8):1035-1043.
- Ansari, K.A. (1990). Heterosis studies in some interspecific hybrids of wheat (*T. aestivum* L.) *Pak. J. Agric., Agril. Engg., Vet Sc.* 6: 37-40.

- Atiq-ur-Rehman, I. Khaliq, M. A. Khan and R. I. Khushnood, (2002). Combining ability studies for polygenic characters in *Aestivum* species. *International Journal of Agriculture and Biology*, 4(1):171-174.
- Awaad, H. A., (2002). Assessment of some genetic parameters using diallel cross fashion and their implications in breeding programs of bread wheat (*T. aestivum* L.). *Zagazig J. Agric. Res.*, 29(4): 1123-1141.
- Ayres, N. M., A. M. McClung and P. D. Larkin, (1997). Microsatellites and a single-nucleotide polymorphism differentiate apparent amylose classes in extended pedigree of US rice germplasm. *Theor. Appl. Genet.* 91:773-781.
- Bahrman, N., J. LE Gous, D. Harini, L. Guilbaud and L. Jestin, (1999). Genetic diversity of old French six-rowed barley varieties assessed with molecular, biochemical and morphological markers and its relation to BaMWV resistance. *Heredity* 83:568-574.
- Barakat, M. N., M. I. Motawei, S. I. Milad, M.A. Moustafa and Y. H. El-Daoudi, (2000). Using RAPD markers for evaluating genetic relationship among wheat cultivars. *Proc. 9th Conf. Agron. Minu. University, Egypt.*
- Barma, N. C. D.; S. H. Khan, M. A. K. Mian, and A. Islam, (1990). Variability and interrelationships of eight quantitative characters in bread wheat (*Triticum aestivum* L.) [in Bangladesh]. *Bangladesh J. of Plant Breeding and Genetics* (Bangladesh). v.3 (1,2) p.71-75.

- Bekheit, B.R., M.G. Mossad, M.A. EL-Morshidy and A. Tamam (1989). Inheritance of yield in some durum wheat crosses. *Assiut J. of Agric. Sci.* 20: 220-234.
- Bhowmik A., M. S. Ali, Z. Sadeque, and K. Saifuddin, (1989). Correlation and path analysis in wheat (*Triticum aestivum* L.) [in Bangladesh]. *Bangladesh J. of Plant Breeding and Genetics* (Bangladesh). Vol 2 (1,2): 23-26.
- Bhowmik, A., Z. Sadeque and M.S. Ali, (1991). Combining ability analysis in wheat (*T. aestivum* L.). *Annals of Bangladesh Agriculture*, 1 (1): 13-18.
- Borghi, B. and M. Perenzin, (1994). Diallel analysis to predict heterosis and combining ability of grain yield, yield components and bread-making quality in bread wheat (*T.aestivum* L.). *Theor. Appl.Genet.*, 89:975-981.
- Botstein, D., R. I. White, M. H. Sholnick and R. W. Davies, (1989). Construction of a genetic map in man using RFLP. *Theor. Appl. Genet.* 32:314-321.
- Boulot, O. A. and E. A. M. El-Sayed, (2001). Quantitative genetic studies on leaf rust resistance and its components in a 7-parents diallel cross of wheat. *J. Agric. Sci. Mansoura Univ.*, 26(1):133-146.
- Barcellos, A. L., A. P. Roelfs and M. I. B. de Moraes-Fernandes, (2000). Inheritance of adult plant leaf rust resistance in the Brazilian wheat cultivar Toropi. *Plant Disease*, 48(1): 90-93.

- Branlard, G. and C.I. Chevalier, (1984). Oiveraity of bread wheats cultivated in France. *Agronomic* 4: 933-938.
- Briggle, L.W., E.L. Cox and R.M. Hays (1967). Performance of a spring wheat hybrid , F₂, F₃ and parent varieties at five population levels. *Crop Sci.* 7:485-490.
- Budak N. and M.B. Yildirim, (1995). Harvest index, biomass production and their relationships with grain yield in wheat. *Pl. Breed. Abst.* 67(7).
- Budak, N., (2001). Heterosis and combining ability in a 8 x 8 diallel durum wheat population. *Ege- niversitesi Ziraat Fakültesi Dergisi*, 38(2/3):55-62.
- Cochran, W.G. and G.M Cox, (1957). *Experimental design*. 2nd ed. John Wiley, N.Y., U.S.A.
- Chague, V., T. Fahima, A. Dahan, G. L. Sun, A. B. Korol, Y. I. Ronin, A. Grama, M. S. Roder and E. Nevo, (1999). Isolation of microsatellite and RAPD markers flanking the Yr15 gene of wheat using NILs and bulked segregant analysis. *Genome* 42(6):1050-1056.
- Chao, S., P. J. Sharp, A. J. Worland, E. J. Warham and R. M. D. Koebner, (1989). RFLP based genetic maps of wheat homoeologous group-7 chromosomes. *Theor. Appl. Genet.* 78:493-504.
- Chen, X., R.F. Line Z. Shi and H. Leung, (1998). Genetics of wheat resistance to stripe rust. *Proc. Of the 9th International Wheat Genetics Symposium*, Vol: 237-239.

- Clark, J.A., J.H. Martin and C.R. Ball, (1922). Classification of American wheat varieties. USOA bulletin 1074. Washington, DC.
- Corbellini, M., M. Perenzin, M. Accerbi, P. Vaccino, B. Borghi, (2002). Genetic diversity in bread wheat, as revealed by coefficient of parentage and molecular markers, and its relationship to hybrid performance. *Euphytica*, 123(2):273-285.
- Cowen, N. M. and K. J. Frey, (1987). Relationships between three measures of genetic distance and breeding behavior in Oat (*Avena Sativa* L.). *Genome* 22:97-106.
- Cox, T.S-, and J.P. Murphy, (1990). The effect of parental divergence on F₂ heterosis in winter wheat crosses- *Theor. Appl. -Genet.* 79: 241-250.
- Darwish, I. H. I., (1992). Breeding studies on wheat. M.Sc. Thesis Fac. Of Agric. Menofiya Univ. Egypt.
- Das, M. K., S. Rajaram, C. C. Mundt and W. E. Kronstad, 1992. Inheritance of slow rusting resistance to leaf rust in wheat. *Crop Science*, 32(6): 1452-1456.
- Dawam, H.A and F. Hendawey, (1990). Inheritance of yield and its components in common wheat (*T. aestivum* L) *Proc . 4th Conf.Agron . Cario . 15-16 Sept.*
- Daya, C., V. Chawla, O. P. Luthra and S. Uppal, (1994). Genetic studies of latent period of leaf rust on some wheat crosses. *Crop Research Hisar*, 7(2):269-272.

- Dedryver, F., M. F. Jubier, J. Thouvenin and H. Goyean, (1996).
Molecular markers linked to leaf rust resistance gene Lr24 in
different wheat cultivars. *Genome*, 39:830-835.
- Delaney, D., S. Nasuda, T. R. Endo and B. S. Gill, (1995).
Cytologically based physical maps of the group-2
chromosomes of wheat. *Theor. Appl. Genet.* 81:245-252.
- Dencic S., R. Kastori, B. Kobiljski, and B. Duggan, (2000).
Evaluation of grain yield and its components in wheat
cultivars and landraces under near optimal and drought
conditions. *Euphytica*. 113: (1): 43-52.
- Deshmukh, P.B., S.B. Alale, M.K. Piinde, S.R. Golhar, S.A.
Sadanshiv, (1999). Genetic divergence in durum wheat. *J.*
Soils Crop. 9: 104-107.
- Devos, K. M., G. J. Bryan, A. J. Collins and P. Stephenson, (1995).
application of two microsatellite sequences in wheat storage
proteins as molecular markers. *Theor. Appl. Genet.* 90:247-
252.
- Dobrotvorskaya, T. V., A. I. Sedlovskii and S. P. Martynov, (1986).
Use of multivariate statistical analysis to select parental pairs:
experimental testing on spring wheat. *Tsitologiya Genetika*
20:452-457.
- Dudely, J. W., (1993). Molecular markers in plant improvement:
Manipulation of genes affecting quantitative traits. *Crop Sci.*,
33:660-668.

- Duncan, D.B. (1955). Multiple range and multiple F-test. *Biometrics*, 11: 1-42.
- Dyck, P. L., (1991). Genetics of adult plant leaf rust resistance in Chinese Spring and Sturdy wheats. *Crop Sci.*, 31: 309-311.
- El-Adl, A.; M. Kosba; Z. M. El-Diasty and G. A. Rizk, (1996). The relative importance of additive and non-additive gene variances in new varieties of wheat (*T. aestivum* L.). *J. Agric. Sci. Mansoura Univ.*, 21: 1717-1733.
- El-Desoky M. A.; A. Ghallab; and E. A. Teama, (2000). Efficient use and uptake of nitrogen for wheat yield in a clay soil in relation to split nitrogen application. *Assiut J. Agric. Sci.* 31(5) 153-167.
- El-Hity, M. A., (1994). Diallel analysis of genetic variation for yield and its components in wheat. *J. Agric. Res. Tanta Univ.*, 20: 490-499.
- El-Maghraby, M. A., (1998). Genetic evaluation of yield and its components in wheat (*T. aestivum* L.) under different environments. M.Sc. Thesis, Fac. Of Agric., Alex. Univ., Egypt.
- El-Maghraby, M. A., M.E. Moussa, N.S. Hana and H.A. Agrama, (2005). Combining ability under drought stress relative to SSR diversity in common wheat. *Euphytica*, 141: 301-308.
- EL-Marakaby, A.M.; O.S. Khalil; A.A. Mohamed and M.F. Abd El-Rahman. (1992). Estimation of the relative importance of characters contributing in yield of a diallel cross of bread

- wheat. Ann. Agric. Sci. A. Shams Univ. Cairo 37: 69-75.
- El-Marakby, A. M., A. A. Mohamed and M. F. Abd El-Rhaman. (1993). Studies on types of gene action and heritability in bread wheat crosses. Ann . Agric. Sci. Ain Shams Unvi. Cario, 37: 151-159.
- El-Saied, E. A. M., (1996). Breeding wheat for high yield. Ph.D. Thesis Fac. Of Agric., Al-Azhar Univ., Egypt.
- El-Shami, M. M., T. M. Shehab El-Din, A. H. Abdel-Latif and M. S. Sharsher, (1996). Heterosis and combining ability for grain yield and some related characters in bread wheat. J. Agric. Sci. Mansoura Univ., 21: 2789-2796.
- Esmail, R. M., (2001). Correlation and path coefficient analysis of some quantitative traits with grain yield in bread wheat (*Triticum aestivum* L.). Bulletin of the National Research Centre (Cairo), 26(3): 395-408.
- Ezzahiri, B. and A. B. Roelfs, (1989). Inheritance and expression of adult plant resistance to leaf rust in Era wheat. Plant disease, 73:549-551.
- Farshadfar, M.,I. Kissimcn and J. Sutka. (1995). Genetic distance between *Triticum timopheevi* Zhuk., *T. araraticum* Jukubz. and *T. aestivum* L. Plant Breeding 114:401-405
- Feuillet C. and B. Keller, (1998). Molecular aspects of biotic stress resistance in wheat. In: Slinkard AE (ed). Proc 9th Int Wheat Genet Symp vol 1. University Extension Press, Saskatoon, Sask. pp 171-177.

- Frei, O. M., C. W. Sluber and M. M. Goodman, (1986). Use of allozymes as genetic markers for predicting performance in maize self cross hybrids. *Crop Sci.* 26:37-72.
- Gale, M. D., M. D. Akthinson, C. N. Chinoy, R. I Harcourt, J. Jia, Q. Y. Li and K. M. Devos, (1995). Genetic maps of hexaploid wheat. *Proc 8th Int. Wheat Genet. Symp.*, China Agric. Sciencetech Press, Beijing, PP 29-40.
- Garc a del Moral L. F., Y. Rharrabtia, D. Villegasb and C. Royob, (2003). Evaluation of grain yield and its components in durum wheat under Mediterranean conditions. An Ontogenic Approach. *Agron. J.* 95:266-274.
- Garland, S. H., L. Lewin, M. Abedinia, R. Henry and A. Blakeney, (1999). The use of microsatellite polymorphism for the identification of Australian breeding lines of rice (*Oryza sativa* L.). *Euphytica*, 108:53-63.
- Goldstein, D. B., A. R. Linares, L. L. Cavalli-Sforza and M. W. Feldman, (1995). An evaluation of genetic resources for use with microsatellite loci. *Genetics*, 139:463-471.
- Goodman, M. M., (1972). Distance analysis in biology. *Syst. Zool.*, 21:174-186.
- Griffing, B., (1956). Concept of general and specific combining ability in relation to diallel crossing systems. *Aus J Biol Sci* 9: 463-493.
- Gupta, P.K. and R.K. Vershney, (2000). The development and use of microsatellite markers for genetic analysis and plant breeding

- with emphasis on bread wheat. *Euphytica*, 113:163-185.
- Gutierrez, O. A., S. Basu, S. Saha, J.N. Jenkins, B.D. Shoemaker, C. L. Cheatham and J.C. McGarty, (2002). Genetic distance among selected cotton genotypes and its relationship with F_2 Performance. *Crop Sci.* 42: 1841-1847.
- Hamada, A.A. (1988). Determination of genetic variance components under different environmental condition in wheat, Ph.D Thesis Fac. Agric. Moshtohor Zagazig Univ., Egypt.
- Hameed, M. Y. and A. H. Adary, (1986). Diallel analysis for grain yield protein content and leaf rust infection in semi-winter wheat: I- Estimates of heterosis and combining ability. *Zanco*, 4(2): 23-36.
- He, S., H. Ohm and S. Mackenzie, (1992). Detection of DNA sequence polymorphism among wheat varieties. *Theor. Appl. Genet.* 84:573-578.
- Helguera, M., I.A. Khan and J. Dubcovsky, (2000). Development of PCR markers for the wheat leaf rust resistance gene Lr47. *Theor. Appl. Genet.* 100:1137-1143.
- Hohmann, U., T. R. Endo, K. S. Gill and B. S. Gill, (1994). Comparison of genetic and physical maps of group-7 chromosomes from *T.aestivum* L. *Molecular and General Genetics.* 245(5):644-653.
- Hosaka, K., M. Mori and K. Ogawa, (1994). Genetic relationships of Japanese potato cultivars assessed by RAPD analysis, *Am. Potato.J.* 71:535-546.

- Ibrahim, K. I. M., (1994). Heterotic performance and combining ability in factorial crosses of bread wheat. *Annals of Agric. Sci., Moshtohor*, 32: 1167-1179.
- Jaccard, P., (1908). Nouvelles recherches sur la distribution locale. *Bull. Soc. Vaud. Sci. Nat.* 44: 223-270.
- Jaint, S.K., C. O. Qualst, G.M. Bhatt and K.K. Wu, (1975). Geographical patterns of phenotypic diversity in world collection of durum wheats, *Crop, Sci.* 15: 700-704.
- Jaradat, A.A. (1991). Phenotypic divergence for morphological and yield related traits among landrace genotypes of durum wheat from Jordan. *Euphytica*. 52: 155-164.
- Jagadev, P.N., (1993). Divergence in wheat under barani condition. *Madras Agric. J.* 80:457-460.
- Jian, R.S., Q. Song, S. Singh, J. Lewis, R. W. ward, P. B. Cregan and B. S. Gill, (2003). Genetic and physical maps of *Xbarc* SSR loci in wheat. *National Fusarium Head Blight Forum Proceeding* 41-45.
- Johanson R., F.G.H. Lupton, (1987). Breeding for disease resistance. In: Lupton FGH (ed) *Wheat breeding. Its scientific basis.* Chapman and Hall, Cambridge. pp 369-418
- Joshi, C. P and H.T. Nguyen, (1993a). Application of RAPD technique for the detection of polymorphism among wild and cultivated tetraploid wheats. *Genome*. 36(3):602-609.
- Joshi, C. P and H.T. Nguyen, (1993b). RAPD (random amplified polymorphic DNA) analysis based intervarietal genetic

- relationships among hexaploid wheats. *Plant Science* 93: 95-103.
- Joshi, S. K. ; S. N. Sharma, D. L. Singhania and R. S. Sain, (2003). Hybrid vigor over environments in a ten-parent diallel cross in common wheat. *SABRAO Journal of Breeding and Genetics*. 35 (2): 81-91
- Kaul, K., R. K. Singh, O. P. Luthra and V. (Chawla, 1991). Genetic analysis of leaf rust resistance and yield attributes in some intervarietal crosses of wheat. *Agricultural Science Digest*, 11 (2): 77-80.
- Khaleeqe, M. H. and A. Alam, (1995). Heritability and number of genes controlling leaf rust resistance in 5 cultivars of wheat in Pakistan. *Pakistan Journal of Phytopathology*, 7(1):5-8.
- Khalifa, M. A., A. A. Ismail, G. R. El-Nagar and I. A. Amin, (1998). Genetical studies on earliness, grain yield and its components in bread wheat. *Assuit Journal of Agricultural Science*, 229(5): 59-69.
- Knott D.R., (1989). The wheat rusts - breeding for resistance. Monographs on theoretical and applied genetics, vol 12 Springer, Berlin Heidelberg New York.
- Koller, B., A. Lehmann, J.M. Medwott and C. Gessler, (1993). Identification of apple cultivars using RAPD markers. *Theor. Appl. Genet.* 85: 901-904.
- Korzun, V., M.S. Roder, K. Wendehake, A. Pasqualone, C. Lotti, M. W. Ganal and A. Blanco, (1999). Integration of dinucleotide

- microsatellite from hexaploid bread wheat into a genetic map of durum wheat. *Theor. Appl. Genet.* 98:1202-1207.
- Larbi, A., A. Mekliche, R. Abed, M. Badis, C. Royo, M.M. Nachit, N. Fonzo, and J.L. Araus, (2000). Effect of water deficit on the yield of two varieties of durum wheat (*Triticum turgidum* var. *durum*) in a semi-arid region. *Proceeding of a seminar, Zaragoza, Spain, 12-14 April 2000.* 40: 295-297.
- Leonardi, A., C. Damerval, Y. Hebert, A. Gallais & D. de Vienne, (1991). Association of protein amount polymorphic (PAP) among maize lines with performance of their hybrids. *Theor Appl Genet* 82: 552-560.
- Li BoQun, Yu GuoDong, Shi YouMing and Tan ChangHua, (2003). Heterosis, combining ability and genetic distance in hybrid wheat involving thermo-photo-sensitive genic male sterility. *Journal of Triticeae Crops*, 23(3): 17-21.
- Ling, H.-Q., Y. Zhu and B. Keller, (2003). High-resolution mapping of the leaf rust disease resistance gene *Lr1* in wheat and characterization of BAC clones from the *Lr1* locus. *Theor. Appl. Genet.* 106:875-882.
- Liu, X., K. Ishiki and W. Wang, (2002). Identification of AFLP markers favorable to heterosis in hybrid rice. *Breeding science* 52:201-206.
- Liu, Y. G. and K. Tsunewaki, (1991). RFLP analysis in wheat. II. Linkage maps of the RFLP sites in common wheat. *Genetics* 53:487-493.

- Liu, Z.Q., Pei. Yand and Pu. ZJ., (1999). Relationship between hybrid performance and genetic diversity based on RAPD markers in wheat, *Triticum aestivum* L. Plant Breeding. 118:119-123.
- Lettering, J. M., A. M. Botha and F.J. Kloppers, (2002). AFLP and RAPD markers linked to leaf rust resistance gene *Lr41* in wheat. South-African-Journal-of Plant-and-Soil. 19(1):17-22.
- Lush, J.L., (1949). Heritability of quantitative characters in farm animals. Eigh Intern Congr. Genetics Proc. (Hereditas Suppl. Vol.) P 356.
- Mahdy, E.E. (1988). Inheritance of some agronomic traits in wheat (*T. aestivum* L). Assiut J. Agric. Sci., 9: 203-212.
- Martin, J.M., L.E. Talbert , S.P. Lanning and N.K. Blake, (1995). Hybrid performance in wheat as related to parental diversity. Crop Sci. 35:104-108.
- Mavi, G.S., G.S Nanda, V.S. Sohu, S. Sharma and Sukhnandan Kaur, (2003). Combining ability analysis for yield and its components in bread wheat (*Triticum aestivum* L.) in two nitrogen regimes. Crop Improvement. 30 (1): 50-57.
- McIntosh R.A., C.R. Wellings. R.F. Park. (1995). Wheat rust: an atlas of resistance genes. CSIRO Australia and Kluwer. Dordrecht.
- McIntosh, R. A., H. S. Barian, R. F. Park and C. R. Wellings, (2001). Aspects of wheat rust research in Australia. Euphytica, 119:115-120.

- McIntosh, R.A., (1992). Close genetic linkage of gene conferring adult-plant resistance to leaf rust and stripe rust in wheat. *Plant Pathology*. 41:523-527.
- Meer, J., H. C. Robert and F. M. Kenneth, (2002). Map Manager version0.22.
<http://www.mapmgr.roswellpark.org/mmQTX.html>.
- Melchinger, A.E., A. Graner, M. Singh & M.M. Messmer, (1994). Relationships among European barley germplasm: I. Genetic diversity among winter and spring cultivars revealed by RFLPs. *Crop Sci* 34: 1191-1199.
- Melchinger, A.E., M. Lee, K.R. Lamkey and W.L. Woodman, (1990). Genetic diversity for restriction fragment length polymorphisms: Relation to estimated genetic effects in maize inbreds. *Crop Sci* 30: 1033-1040.
- Michelmore, R.W., I. Paran and R.V. Kasseli, (1991). Identification of markers linked to disease resistance genes by bulked segregant analysis: A rapid method to detect markers in specific genome regions by using segregating populations. *Proc. Acad. Sci. U.S.A.* 88: 9852.
- Mickelson-Young, L., T. R. Endo, B. S. Gill, (1995). A cytogenetic ladder-map of the wheat homoeologous group-4 chromosomes. *Theor. Appl. Genet.* 90(7/8):1007-1011.
- Mitkees, R.A. and H.N. EL-Rssas, (1986). Gene action in the inheritance of some characters of bread wheat . *Egypt J. Genet. Cytol.*, 15: 135-146.

- Mitkees, R.A. and N.F. Dawla, (1983). Genetic analysis of a bread wheat cross. Ann. of Agric. Sci. Moshtohor, 20: 225-236.
- Mohamed, A.A., A.M. Ismail and S.A. Mohamed, (1987). Genetic behaviour of some characters in a wheat cross under salinity condition. Ann. of Agric. Sci. Moshtohor, 25: 141-158.
- Moreno, M. and J. Gonzalez, (1992). Estimates of markers-associated QTLs effects in Montana Carlo back cross generations using multiple regression. Theor. And Appl. Genet. 85: 423-434.
- Moser, H. and M. Lee, (1994). RFLP variation and genealogical distance multivariate distance, heterosis, and genetic variance in oats. Theor Appl Genet 87: 947-956.
- Motawei, M. I., M. N. Barakat, S. I. Milad, M. A. Moustafa and Y. H. El-Daoudi, (2001). Molecular markers linked to leaf rust resistance gene *Lr29* in F₂ wheat population. Alex. J. Agric. Res. 46:57-62.
- MSTAT-C, (1991). A Software Program for the Design, Management, and Analysis of Agronomic Research Experiments. Michigan State University, East Lansing, USA.
- Mukhtar, M. S., Mehboob-ur-Rahman, Y. Zafar, (2002). Assessment of genetic diversity among wheat (*T. aestivum* L.) cultivars from a range of localities across Pakistan using RAPD analysis. Euphytica 128(3):417-425.
- Murray, M. and W.F. Thompson, (1980). Rapid isolation of high molecular-weight plant DNA. Nucleic Acids Res 8: 4321-4325.

- Naik, S., K S Gill, V S Prakasa Rao, V S Gupta, S A Tamhankar, S Pujar, B S Gill and P K Ranjekar, (1998). Identification of a STS marker linked to the *Aegilops speltoides*-derived leaf rust resistance gene *Lr28* in wheat. *Theor. Appl. Genet.* 97:535-540.
- Nanda, G.S, P.S. Virk and K.S. Gill, (1983). Diallel analysis over environments in wheat-plant characters and harvest index. *Indian J Genet* 43: 21-27.
- Nei, M., (1972). Genetic distance between populations. *Am. Nat.* 106: 283-292.
- Parentoni, S.N., J.V. Magalhães, C.A.P. Pacheco, M.X. Santos, T. Abadie, E.G. Gama, P.E.O. Guimarães, W.F. Meirelles, M.A. Lopes, M.J.V. Vasconcelos and E. Paiva, (2001). Heterotic groups based on yield-specific combining ability data and phylogenetic relationship determined by RAPD markers for 28 tropical maize open pollinated varieties. *Euphytica* 121: 197-208.
- Paull, J.G., K. J. Chalmers, A. Karakousis, J. M. Kretschmer, S. Manning and P. Langridge, (1998). Genetic diversity in Australian wheat varieties and breeding material based on RFLP data. *Theor. Appl. Genet.* 96:435-446.
- Perenzin, M., M. Corbellini, M. Accerbi, P. Vaccino and B. Borghi, (1998). Bread wheat: F₁ hybrid performance and parental diversity estimates using molecular markers. *Euphytica*, 100:273-279.

- Plaschke, J., A. Borner, K. Wendehake, M. W. Ganal and M. S. Roder, (1996). The use of wheat aneuploids for the chromosomal assignment of microsatellite loci. *Euphytica* 89: 33-40.
- Plaschke, J., M. W. Ganal, and M. S. Roder, (1995). Detection of genetic diversity in closely related bread wheat using microsatellite markers. *Theor. Appl. Genet.* 91:1001-1007.
- Powell, W., C.G. Machray and J. Provan, (1996). Polymorphism revealed by simple sequence repeats. *Plant Science* 1:215-221.
- Prabhu, K. V., S. K. Gupta, A. Charpe and S. Koul, (2004). SCAR marker tagged to the alien leaf rust resistance gene *Lr19* uniquely marking the *Agropyron elongatum*-derived gene *Lr24* in wheat. *Plant Breeding*, 123: 417-420.
- Pujar, S., S.A. Tamhankar, V.S. Rao, V.S. Gupta, S. Naik and P.K. Ranjekar, (1999). Arbitrally primed PCR based diversity assessment reflects hierarchical grouping of Indian tetraploid wheat genotype. 99: 868-876.
- Robert, O. F. Dedryver, B. Rolland, C. Abelard and B. Jaudean, (2000). Relationships between molecular identification of gene *Yr17* and adult plant resistance against stripe rust and leaf rust in bread wheat varieties. *Acta Phytopathologica et Entomologica Hungarica*. 35:59-63.
- Roder, M. S., V. Korzun, K. Wendehake, J. Plaschke, M.H. Tixier, P. Leroy and M. Ganal, (1998). A microsatellite map of wheat. *Genetics* 149:2007-2023.

- Rolhf, F.J., (2000). NTSYS-pc: Numerical taxonomy and multivariate analysis system. Version 2.02 Exeter Software, Setauket, N.Y.
- Roy, J.K. , H.S. Balyan, M. Prasad and P.K. Gupta, (2002). Use of SAMPL for a study of DNA polymorphism, genetic diversity and possible gene tagging in bread wheat. Theor. Appl. Genet. 465-472.
- Sagahi-Maroo, M., K. Soliman, R. Jorgensens and R. Allard, (1984). Ribosomal DNA spacer length polymorphisms in barley: Mendelian inheritance, chromosomal location and population dynamic. Proc. Nat. Acad. Sci. 81: 8018, U.S.A.
- Sairam R.K.; and K.K. Singh, (1989). N-use efficiency, N assimilation, and morphophysiological traits in barley. ICARDA. Barley and Wheat Newsletter. V.8(2) p. 26-28.
- Salama, S. M., (2000). Partial diallel analysis and heterosis in bread wheat (*T. aestivum* L.). Zagazig J. Agric. Res., 27:1371-1384.
- Sant, V.J., A.G. Patankar, N.D. Sarode, L.B. Mhase, M.N. Sainani, R.B. Deshmukh, P.K. Ranjekar & V. S. Gupta, (1999). Potential of DNA markers in detecting divergence and in analyzing heterosis in Indian elite chickpea cultivars. Theor Appl Genet 98: 1217-1225.
- SAS, (1985). SAS/STAT guide for personal computers. Version 6 edn. SAS end. SAS Institute. Cary, N. C.
- Sayre K. D., S. Rajaram and R. A. Fischer, (1997). Yield potential progress in short bread wheat in Northwest Mexico. Crop Sci.

37: 36-42.

- Semeena, S., and L. Singh, (2001). Studies on path co-efficient analysis of harvest index and its related traits in wheat. *Indian Journal of Agricultural Research*, 35(2): 127-129.
- Seyfarth, R., C Feuillet, G Schachermayr, M Winzeler and B Keller, (1999). Development of a molecular marker for the adult plant leaf rust resistance gene *Lr35* in wheat. *Theor. Appl. Genet.* 99:554-560.
- Shamsuddin, A.K.M., (1985). Genetic diversity in relation to heterosis and combining ability in spring wheat. *Theor. Appl. Genet.* 70: 306-308
- Shehab El-Din, T. M., M. M. El-Shami and A. H. Abdel-Latif, (1996). Qualitative and Quantitative genetic studies on *Triticum aestivum*: *Puccinia recondite* interaction. *J. Agric. Sci. Mansoura Univ.*, 21(11): 3769-3778.
- Sherif, H., N.F. Dawla and A.M. Abdel-Safi, (1985). Genetic studies on some quantitative characters in wheat (*T. aestivum* L.). *Ann. of Agric. Sci. Moshtohor*, 23: 134-141.
- Shikano, T. and N. Taniguchi, (2002). Using microsatellite and RAPD markers to estimate the amount of heterosis in various strain combinations in the guppy (*Poecilia reticulata*) as a fish model. *Aquaculture*, 204:271-281
- Singh, D., R. F. Park and R. A. McIntosh, (1999). Genetic relationship between the adult plant resistance gene *Lr12* and the complementary gene *Lr31* for seedling resistance to leaf

- rust in common wheat. *Plant Pathology*, 48:567-573.
- Singh, P.K., (1994). Genetic diversity in durum wheat (*Triticum durum* Desf.) germplasm. *Ann. Agric. Res.* 15: 418-422.
- Smith, H.H., (1952). Fixing transgressive vigor in *Nicotiana rustia*. Iowa State College Press, Ames, Iowa, U.S.A.
- Smith, J.S.C., (1984). Genetic variability within U.S. hybrids using multivariate analysis of isozymes data. *Crop Sci.* 24: 1041-1045.
- Smocek, J. and P. Martinek., (1997). Divergence between winter wheat parents in yield per ear. *Sbornik UVTIZ Genetika a Slechteni.* 23: 265-271.
- Snedecore, G.W. and W.G.Cochran, (1967). Statistical methods. 6th ed. Iowa State University Press.
- Solomon, K. F. and M. T. Labuschagne, (2004). Inheritance of evapotranspiration and transpiration efficiencies in diallel F₁ hybrids of durum wheat (*Triticum turgidum* L. var. *durum*). *Euphytica*, 136(1): 69-79.
- Solomon, K. F., M. T. Labuschagne and A.T.P. Bennie, (2003). Responses of Ethiopian durum wheat (*Triticum turgidum* var. *durum* L.) genotypes to drought stress. *South African Journal of Plant and Soil.* 20: 2, 54-58.
- Song Ge., Giancalo, C.X. Oliveria , Barbara, A. Schaal , Gao Li-Zhi and Hong DE-Ydan, (1999). RAPD variation within and between natural population of the wild rice *Oryza rufipogon* from China and Brazil. *Heredity*, 82:638-644.

- Sornprach, T. A., (1988). Phenotypic studies in wheat and triticale. M.Sc Thesis in Agric. Bangkok (Thailand).
- Souza, E., and M.E. Sorrelles, (1991a). Prediction of progeny variation in oat from parental genetic relationships. *Theor Appl Genet.* 82: 233-241.
- Souza, E., and M.E. Sorrelles, (1991b). Relationship among 70 North American Oat germplasms: II cluster analysis using qualitative characters. *Crop Sci.* 31:605-612.
- Spagnoletti, P. L., and C. O. Qualset, (1987). Geographical diversity for quantitative spike characters in a world collection of durum wheat. *Crop Science*, 27:235-241.
- Stachel, M., T. Lelley, H. Grausgruber and J. Vollmann, (2000). Application of microsatellite in wheat (*T. aestivum* L.) for studying genetic differentiation caused by selection for adaptation and use. *Theor. Appl. Genet.* 100:242-248.
- Stephenson, P., G. Bryan, J. Kirby, A. Collins, K. Devos, C. Busso and M. Gale, (1998). Fifty new microsatellite loci for the wheat genetic map. *Theor. Appl. Genet.* 97:946-949.
- Stiles, J.I., C. Lemme, S. Sondur, M. B. Morshidi and R. Manshardt, (1993). Using randomly amplified polymorphic DNA for evaluating relationships among Papaya cultivars. *Theor. Appl. Genet.* 85:697-701.
- Stubbs, R.W., J.M Prescott, E.E. Saari and H.J. Dubin, (1986). Cereal Disease Methodology Manual. CIMMYT, Mexico pp. 22.

- Sun, Q., Y. Wei, Z. Ni, C. Xie, T. Yang, (2002). Microsatellite marker for yellow rust resistance gene *Yr5* in wheat introgressed from spelt wheat. *Plant Breeding* 121(6):539-541.
- Talbert, L. E., S. P. Lanning, R. L. Murphy and J. M. Martin, (2001). Grain filling duration in twelve hard red spring wheat crosses: Genetical variation and association with other agronomic traits. *Crop Sci.*, 41: 1390-1395.
- Tanksley, S. D., (1983). Molecular markers in plant breeding. *Plant Mol. Bio. Rep.* 1:3-8.
- Trethowan, R. M., O. Abdalla, and W.H. Pfeiffer, (1991). Evaluation of the rate and duration of grain filling in triticale and its association with agronomic traits. *Proceedings of the second International Triticale Symposium. CIMMYT, 1991* P.128-130.
- Tsegaye, S., G. Belay, A.A. Jaradat, (1998). Variation for isozymes and agronomic traits in Ethiopian tetraploid wheat (*Triticum durum* L.) landraces. *Triticeae III. Proc. of 3th International Triticeae Symposium, Aleppo, Syria, 4-8 May 1991* 191-198.
- Uddin, N. M., F. W. Ellison, L. O. Brien and B. D. H. Latter, (1992). Heterosis in F_1 hybrids derived from crosses of adapted Austrian wheats. *Aust. J. Agric. Res.*, 43: 907-917.
- Van Beuningen, L. T. and R. H. Busch, (1997). Genetic diversity among North American spring wheat cultivars: III. Cluster analysis based on quantitative morphological traits. *Crop Science*. 37: 981-988.

- Vershney, R. K., P. C. Sharma, P. K. Gupta, H. S. Balyan, B. Ramesh, J. K. Roy, A. Kumar and A. Sen, (1998). Low level of polymorphism detected by SSR probes in bread wheat. *Plant Breeding* 117:182-184.
- Vierling, R. A., and R. Ngugen, (1992). Use of RAPD markers to determine genetic diversity of wheat genotypes. *Theor. Appl. Genet.* 84:835-838.
- Wang CuiLing, Gao HaiTao, Wang ShuZi, Duan GuoHui, Wu ShaoHui, Zhang XuePin and Lu ShuZuo, (2003). Genetic analysis of quality and yield characters in winter wheat. *Journal of Triticeae Crops*, 23 (3):26-28.
- Wang, L. F., J. X. Ma, R. H. Zhou, X. M. wang, J. Z. Jia, (2002). Molecular tagging of the yellow rust resistance gene *Yr10* in common wheat, P.I.178383 (*T. aestivum* L.). *Euphytica* 124(1):71-73.
- Wang, Y.B., (1988). Study on the changes of genetic distance under different conditions. *Acta. Agric. Univ. Jilinensis*, 10:18-22.
- Wang, Z.Y., G. Second and S.D. Tanksley, (1992). Polymorphism and phylogenetic relationships among species in the genus *Oryza* as determined by analysis on nuclear RFLPs. *Theor. Appl. Genet.* 83: 565-581.
- Welsh, J. and M. McClelland, (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.* 18:7213-7218.

- Whittaker, J. C., R. N. Curnow, C. S. Haley and R. Thompson, (1995). Using marker-maps in marker-assisted selection. *Genet. Res. Camb.* 66:255-265.
- Wilkie, S.E, P.G. Isaac and R.J. Slater, (1993). Random amplified polymorphic DNA for genetic analysis in *Allium*. *Theor. Appl. Genet.*, 86: 497-504.
- William, H. M., D. Hoisington, R. P. Singh and D. Gonzalez-de-Leon, (1997). Detection of quantitative trait loci associated with leaf rust resistance in bread wheat. *Genome* 40(2):253-260.
- Williams, M., R. P. Singh, J. Huerta-Espino, S. Ortiz-Isilas, D. Hoisington, (2003). Molecular marker mapping of leaf rust resistance gene *Lr46* and its association with stripe rust resistance gene *Yr29* in wheat. *Phytopathology* 93(2):153-159.
- Xiong, L.Z., G.P. Yang, C.G. Xu, Qifa Zhang and M.A. Sagahi Maroof, (1998). Relationships of differential gene expression in leaves with heterosis and heterozygosity in a rice diallel cross. *Molecular Breeding* 4:129-136.
- Yadav, B., C. S. Tyagi and D. Singh, (1998). Genetical studies and transgressive segregation for yield resistance to leaf rust of wheat. *Wheat Information Service*, 87: 15-21.
- Zalewski, D., (2001). Estimation of general and specific combining ability of quantitative traits of winter wheat. *Biuletyn Instytutu Hodowli i Aklimatyzacji Roslin*, 216(2): 267-272.
- Zeller, F. J., (1973). 1B/1R wheat-rye chromosome substitutions and

- translocation. In "Proceeding of the Fourth International Wheat Genetics Symposium". (Eds ER Sears and LMS Sears.) pp. 209-221.
- Zeven, A. C. and R. Schachl, (1989). Groups of bread wheat landraces in Austrian Alps. *Euphytica* 41: 235-246.
- Zhao, M.F., X.H. Li, J. B. Yang, C.G. Xu, R.Y. Hu, D.J. Liu and Q. Zhang, (1999). Relationship between molecular marker heterozygosity and hybrid performance in intra-and inter-subspecific crosses of rice. *Plant Breeding* 118:139-144.