

CONTENTS	Page No.
INTRODUCTION	1
REVIEW OF LITERATURE	4
1. Plant parasitic nematodes associated with sugarbeet plants.....	4
Plant parasitic nematodes-agrometrological factors relations	10
2. Arthropod fauna associated with cultivated soil.....	13
Soil mites- ogrometerological factors relations.....	22
3. Relationship between plant parasitic nematode and soil organisms.....	24
3.1. Plant parasitic nematodes-soil predaceous mites relations.....	24
3.2. Plant-parasitic nematodes- soil fungi relations...	27
3.3. Plant parasitic nematodes-soil bacteria relations.	32
4. The effect of organic matter on the population dynamics of the nematophagous fungi.....	36
5. Susceptibility of sugarbeet varieties to plant parasitic nematodes.....	40
6. Sugarbeet yield losses as response of nematode densities.....	45
MATERIALS AND METHODS	49
1. Soil organisms survey.....	49
1.1.Arthropod fauna.....	49

1.1.1. Sampling, extraction, enumeration and identification of arthropod fauna associated with sugarbeet plants	49
1.2. Nematode fauna.....	50
1.2.1. Sampling, extraction, enumeration and identification of nematodes fauna associated with sugarbeet plants.....	50
1.2.3. The " importance " and " prominence " values of nematodes.....	51
1.2.3.1. Measuring of distribution uniformity or rate of occurrence.....	51
1.2.3.2. Relative density (<i>Rd</i>) of each species was calculated as percentage.....	52
1.2.3.3. Relative biomass (<i>Rb</i>).....	52
1.2.3.4. The importance of nematode genus or species in the community.....	53
1.2.3.5. Prominence value (<i>Pv</i>).....	53
1.2.4. Field population dynamics of root-knot nematode, <i>Meloidogyne incognita</i> on sugarbeet plants.....	53
1.3. Microorganisms	54
1.3.1. Fungi and bacteria recovered from soil of sugarbeet fields.....	54
1.3.2. Fungi recovered from nematode eggs.....	54

1.3.3. Fungi recovered from nematode.....	55
1.3.4. Purification of different microorganism isolates for pure culture.....	55
1.3.4.1. Identification of microorganisms.....	56
1.3.4.1.1. Fungal isolates.....	56
1.3.4.1.2. Bacterial isolates.....	56
2. Effect of soil microorganisms on the suppression of the root-knot nematode infecting sugarbeet plants.....	56
2.1. Effect of different soil fungi on <i>Meliodogyne</i> <i>incognita</i> larvae	
2.1.1. Laboratory bioassay test.....	57
2.1.1.1. Fungal filtrate preparation.....	57
2.1.1.2. Bioassay test.....	58
2.1.2. Greenhouse tests.....	59
2.1.2.1. Fungal inoculum preparation.....	59
2.1.2.2. Experiment.....	59
2.1.2.3. Determination of sugarbeet yields and quality	60
2.1.2.4. Statistical analysis of Data.....	61
2.2. Effect of different soil bacteria on <i>Meliodogyne</i> <i>incognita</i> larvae	61
2.2.1. Bacterial isolates.....	61
2.2.1.1. Bacterial filtrates preparation.....	61
2.2.1.2. Bioassay test.....	62
2.2.2. Greenhouse test.....	62

2.3.Effect of predacious soil mites against root-knot nematode, <i>M. incognita</i>	64
2.3.1. Laboratory experiment.....	64
2.3.2. Greenhouse experiment.....	66
3. Interrelationship among nematophagous fungus, predator nematode and plant parasitic nematode	68
3.1. Organisms and cultivation methods.....	68
3.2.Quantification of nematodes.....	69
3.3.Quantification of <i>Arthrobotrys conoides</i>	69
3.4.Microcosms experiments.....	70
4. Evaluation of some animal manure for mass production of some nematophagous fungi and related effect on <i>M.</i> <i>incognita</i> infected of sugarbeet	71
4.1. Tested fungi.....	71
4.2. Substrates and their preparation.....	71
4.3. Laboratory test.....	72
4.4. Greenhouse test.....	73
5. Susceptibility of sugarbeet varieties to root-knot nematode, <i>Meloidogyne incognita</i> infection.....	74
5.1. Field experiment.....	74
5.2. Greenhouse experiment.....	77
6. The relationship between initial population of the <i>Meloidogyne incognita</i> and sugarbeet yields	78
RESULTS AND DISCUSSION	82
1. The soil organisms in sugarbeet field at the Ten-	82

thousand faddans area, West Nubaryia	
1.1. Animal fauna.....	82
1.1.1.Arthropod fauna.....	84
1.1.1.1.Acari fauna.....	85
1.1.1.2. Insect fauna.....	89
1.1.1.3. Other arthropod fauna.....	89
1.1.2. Annelida and Mollusca.....	90
1.1.3.Nematode fauna.....	90
1.1.4. Monthly variations of predators (arthropod and nematode fauna) in sugarbeet field at theTen- thousand faddans area, West Nubaryia.....	92
1.1.4.1. Predator arthropod fauna.....	92
1.1.4.1.1. Predator mites.....	92
1.1.4.1.2. Predator insects.....	93
1.1.4.2. Predator nematode fauna.....	96
2. Plant parasitic nematodes associated with sugarbeet in the ten thousand faddans area, West Nubaryia...	97
2.1. species of plant parasitic nematodes.....	97
2.2. Occurrence rate and absolute desity.....	97
2.3.Monthly fluctuation and population densities through the growing season	100
2.4."Importance and Prominence " values	103
3. Field population dynamics of the root-knot nematode, <i>Meloidogyne incognita</i> at different growing stages of sugarbeet plant.....	106

4. Relationship between soil microorganisms and the root- knot nematode, <i>Meloidogyne incognita</i>	109
4.1. Bacterial and fungal isolates.....	109
4.1.1. Fungal isolates	110
4.1.2. Bacterial isolates	110
4.2. Effect of soil fungi and bacteria on the <i>Meloidogyne incognita</i> population and productivity of sugarbeet	110
4.2.1. Soil fungi.....	110
4.2.1.1. Bio-assay test in laboratory.....	110
4.2.1.2. Greenhouse test.....	112
4.2.2. Soil bacteria	120
4.2.2.1. Bioassay test in laboratory.....	120
4.2.2.2. Greenhouse test.....	122
4.3. Effect of soil mites on <i>Meloidogyne incognita</i> population and growth response of sugarbeet root...	129
4.3.1. In laboratory.....	129
4.3.2. In greenhouse.....	131
5. Interrelationship among plant parasitic nematode <i>Meloidogyne incognita</i> , nemtrophagous fungus <i>Arthrobotrys conoides</i> and predator nematode <i>Diplogaster sp.</i> in the rhizosphere soil of sugaebeet Plants	134
5.1. Population dynamic of <i>Arthrobotrys conoides</i> with or without <i>Diplogaster sp.</i> or <i>Meloidogyne incognita</i>	134

5.2. Population dynamic of <i>Diplogaster sp.</i> with or without <i>Arthrobotrys conoides</i> or <i>Meloidogyne incognita</i>	136
5.3. Population dynamic of <i>Meloidogyne incognita</i> with or without <i>Arthrobotrys conoides</i> or <i>Diplogaster sp.</i>	138
6. Evaluation of some animal manures for mass production of six nematode- parasitic fungi and related effects on <i>Meloidogyne incognita</i> infecting sugarbeet plants	142
6.1. Laboratory test.....	144
6.1.1. Fungal growth on substrates.....	144
6.1.1.1. Mycelial growth.....	144
6.1.1.2. Sporulation.....	146
6.2.Greenhouse test.....	148
7. Susceptibility of sugarbeet varieties to the root-knot nematode, <i>Meloidogyne incognita</i> infection.....	153
7.1. Field experiment.....	153
7.2. Greenhouse experiment.....	156
8. Relationship between initial population density (Pi) of the root-knot nematode, <i>Meloidogyne incognita</i> and sugarbeet yields.....	158
SUMMARY AND CONCLUSION	165
REFERENCES	177
ARABIC SUMMARY	

From R to L
side 12 pp.