




REFERENCES

REFERENCES

- Abbassy, M. A.; M. Ashry; F. Adam; F. Khalil and M. A. Abou-Shlou (1984): Toxicological and histopathological studies on the cotton bollworm, *Pectinophora gossypiella* (Saund). Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent., **49** (3):691-698.
- Abd El-Hafez, A.; A. G. Metwalley and M. R. A. Saleh (1982): Rearing of the pink bollworm, *Pectinophora gossypiella* (Saunders) on kidney bean diet (Lepidoptera: Gelechiidae). J. Econ. Entomol., **49**: 559-560.
- Abd El-Geliel, Reda, M.M. Amer (1997): Toxicological studies on the cotton bollworms. M. Sc. Thesis, Faculty of Agric. Cairo Univ.
- Abdel-Megeed, M. I.; W. M. Watson; A. A. Zidan; G. M. Hegazy and N. M. Hussein (1987): The ovicidal efficiencies of synthetic insecticides and insect growth regulators against the spiny and pink bollworms. Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent., **52** (2b) : 495 - 499.
- Abdel-Sattar, M. M. and M.A. El-Guindy (1988a): Effect of the juvenoid triprene on the biological activity of the pink bollworm, *Pectinophora gossypiella* (Saund). Agric. Res. Rev., **66** (1): 19-23.

- Abdel – Sattar, M.M and M.A. Guindy (1988 b):** Joint action of insect growth regulator, pyrethroid mixtures on the biological activity of the pink bollworm, *Pectinophora gossypiella* (Saund.). Agric. Res. Rev., **66**: 33 – 38.
- Aioub, A. A. A.; S. A. A. Raslan; S. S. M. Hassanein and M. E. A. Hegab (2003):** Effect of certain insecticides on the pink bollworm, *Pectinophora gossypiella* (Saund.). Zagazig. J. Agric. Res., **30** (4) :1711-1734.
- Ali, M. I.; A.J. Treen; N. G. Roy and M. A. Rahman (1995):** Comparative performance of different types of sprayers for the control of cotton insect pests in Bangladesh. Bangladesh J. of Entomol., **5** (1-2): 53-55.
- Allen, C. T.; S. Frizzell and A. C. Riddle (1997):** Effect of insecticide, rate, and egg age on ovicidal control. Proceedings Beltwide Cotton Conferences, New Orleans, LA, USA, January 6-10, Volume 2 :1264-1268.
- Amer, A. E. A. (2004):** Ecological and physiological studies on bollworms. Ph.D. Thesis. Fac. of Agric. Zagazig, Univ. Moshtohor.
- Bariola, L.A. (1984):** Pink bollworm (Lepidoptera: Gelechiidae) effects of low concentrations of selected insecticides on mating and fecundity in the laboratory. J. Econ. Entomol., **77** (5): 1278-1282.
- Bruce, E.T.; M.D. Rethwisch and M.W. Johnson (1988):** Variation in adult mortality and knockdown caused by insecticides among populations of diamondback moth

(Lepidoptera:Plutellidae). J. Econ. Entomol., **81** (2):437-441.

Cashad, S. S.; M. V. Malipatil; P. D. Sharma and O. P. Chaudhary, (2001): Relative performance of different sprayers, nozzles and spray volumes in controlling insect-pests of cotton in Haryana. Crop Res. Hisar., **21**(3): 324-331.

Chhuneja, P.K.; H. S. Singh and H. Singh (1991): Economics of spray technology for pest control on cotton. J. of Insect Sci., **4**(2): 200-202.

Cook, D.R.; Leonard, B. R. and J. Gore (2004): Field and laboratory performance of novel insecticides against armyworms (Lepidoptera: Noctuidae). Florida Entomologist, **87**(4): 433-439.

Dabalbaje, B. B. and A. D. Deshpande (1985): Efficacy of different spray volumes of insecticides against cotton bollworms on NHH-1 and SRT-1. Pesticides, **19** (9):47-48.

Dhawan, A. K.; G. S. Simwat and A. S. Sidhu (1992): Note on the possible use of moulting inhibitor in management of bollworms on cotton. Indian J. of Plant Protection, **20** (1): 72-74.

Douglas, W. F. ; R. Cecil and J.R. Smith (1992): Pink bollworm (Lepidoptera: Gelechiidae) penetration into and survival in bolls of resistant and Susceptible cotton lines. J. Agric. Entomol., **9** (3): 165 – 173.

- Edomwande, E.O.; Schoeman, A.S.; Brits, J. A. and M. van-der Merwe (2000):** Effect of the chitin synthesis inhibitor lufenuron on the American bollworm. The BCPC Conference: Pests and Diseases, Volume 1, Proceedings of an International Conference Brighton, UK, 13-16 November; 523-526.
- Ellington, J. (1990):** Proposal of the collaborative research project 'cotton IPM with emphasis on biological control of pink bollworm' USA-NARP Project in Egypt.
- El-Sheikh, Fatma, M.; Abo-Seda, Samia, A. B. and A. M. Al-Beltagy (1994):** Laboratory and field studies on the effect of certain insecticides, IGR's, insecticides mixtures, IGR's and oils on *Pectinophora gossypiella* (Saund.) (Lep. Gelechiidae). Egypt. J. Appl. Sci., **9** (2): 513 – 524.
- Finney, D. I. (1971):** Probit analysis 3rd ed., Cambridge Univ. Press, London.
- Fisher, A.R. (1950):** Statical method for research workers. Oliver and Poyed Edinburgh and London.
- Flint, H.M. and R.L. Smith (1977):** Laboratory evaluation of TH 6040 against the pink bollworm. J. Econ. Entomol., **70**:51-53.
- Graves, J. B.; B. R. Leonard; A. M. Pauloff; G. Burrisik; K. Ratchford and S. Micinski (1988):** Monitoring pyrethroid resistance management implications. J. Agric. Entomol., **5** (2): 109-115.

- Harcharan, S.; P.K. Chhuneja and H. Singh (1987):** Comparative performance of h.v., l.v. and u.l.v. sprays for the control of cotton bollworms. *Tropical Pest Management*, **33** (1):73-80.
- Haynes, K.F. and T. C. Baker (1985):** Sublethal effects of permethrin on the chemical communication system of the pink bollworm moth, *Pectinophora gossypiella*. *Archives of Insect Biochemistry and Physiology*, **2**(3): 283-293.
- Hegab, M. E. A. (2002):** Studies on the bollworms infesting cotton in Sharkia Governorate, Egypt. M. Sc. Thesis, Faculty of Agric. Zagazig Univ.
- Henderson, C. F. and E.W. Telton (1955):** Tests with acaricides against the brown wheat mite. *J. Econ. Entomol.*, **48**: 157-161.
- Horowitz, A. R.; M. Klein; S. Yablonski and I. Ishaaya (1992):** Evaluation of benzoylphenyl ureas for controlling the spiny bollworm, *Earias insulana* (Boisd.), in cotton. *Crop Protection*, **11** (5): 465-469.
- Hussein, N. M.; M. Naguib ; W. M. Watson; M. I. Abdel-Megeed and F. A. Mariy (1985):** Ovicidal and latent biological activities of IGR, insecticides and their binary mixtures on the American bollworm *Heliothis armigera* (Hbn.). *Bull. of the Entomol. Soc. of Egypt, Econ. Ser.*, **14**: 413-421.
- Jai-Singh; H. Sing; J. Singh and H. Singh (1987):** Determination of optimum droplet parameters and insecticidal deposit at

the target sites vis-à-vis spray swaths for the control of bollworm on cotton. *Tropical Pest Management*, **33**(2): 182-187.

Jai-Singh and H.Singh (1989): Optimization of insecticidal deposit for the control of bollworms on cotton. *J. Entomol. Res.*, **13**(1-2): 43-56.

Javaid, I. (1991): Application of insecticides on cotton in Zambia: comparison of application techniques. *Insect Science and its Application*, **12**(1-3): 111-125; In *Proceedings of the Second International Conference on Tropical Entomol.*, held in Nairobi, Kenya, on 31 July-4 August .

Kathuria, V.; Pala Ram and R.K. Saini (2000): Susceptibility of *Heliothis armigera* (Hubner) eggs to different insecticides. *J. of Entomol. Res.*, **24**(3): 213-217.

Khodzhaev, S. H. and O.T. Eshmatov (1983): The ovicidal action of pyrethroids. *Zashchita, Rastenii.*, (7): 24-25.

Khurana , A. D. and A.N. Verma (1990): Comparative damage caused by bollworms and yield of seed cotton. *J. insect Sci.*, **3** (2): 180 – 182

Klein, M.; S. Levski and S. Keren (1982): Comparative toxicity of several pyrethroids, parathion and chlorpyrifos to adults of the spiny bollworm, *Earias insulana* (Boisd.) (Lepidoptera: Noctuidae), in laboratory assays. *Israel J. of Entomol.*, **16** : 99-103.

Kostandy, S. N; M. F. Rofail; S. H. Taher and A. A. El-Hafez (1999): Effect of an insect growth inhibitor on the newly

hatched larvae of *Pectinophora gossypiella* (Saund.); cyanox resistant and susceptible strains. Egyptian J. of Agric. Res., 77(4): 1477-1484 .

Leonova, I.N. and N.M. Slynko (1996):Comparative study of insecticide susceptibility and activity of detoxification enzymes in larvae and adult of cotton bollworm, *Heliothis armigera*. Archives of insect biochemistry and Physiology .32:157-172.

Little, T. M. and F.J. Hills (1975): Statistical methods in agricultural research available from U.C.D. Book store, University of California, Davis:241pp.

Lopez, J.D., Jr and M.A. Latheef, (2000): Effect of hexaflumuron on feeding response and reproduction of adult bollworm. Proceedings Beltwide Cotton Conferences, San Antonio, USA, 4 -8 January, Volume 2; 1316-1321.

Mahar, A. N.; Munir, M.; Darban, D.A.; Mahar, A.Q.and G.S. Markhand (2004): Field evaluation of different insecticides against pink bollworm on four cotton varieties in Sindh, Pakistan. Indus-Cottons, 1(2): 65-72.

Malipatil, M. V.; S. S Dashad; P. D Sharma and O. P. Chaudhary (2002): Studies on the effect of spray volumes on insect-pests and yield of seed cotton in Haryana. Crop Res. Hisar., 24(1): 166-172.

Mambiri,A. M. (1987) : Evaluation of some crop sprayers in the application of insecticides on cotton in Kenya. Tropical Pest Management, 33 (3): 189-191.

- Meisner, J. ;K. R. S Ascher and F. Assi (1987):** Effect of some benzoylphenyl ureas on *Earias insulana* eggs of different age. *Phytoparasitica*, **15** (2):91-96.
- Meisner, J. ; M. Klein and K.R.S. Ascher (1986):** The effect of some benzoyl phenylureas on the larvae of *Earias insulana*. *Phytoparastica*, **14** (1): 3 –9.
- Mohamady, Aziza, H. (2000):** Biochemical and toxicological studies on the cotton leafworm, *Spodoptera littoralis* (Boisd.). M. Sc. Thesis, Fac. Agric., Zagazig Univ.
- Mourad, M.A.; M.E. Omar and A.A. Mahran (1991):** Field potency of different insecticides against cotton bollworms. *Egyptian J. of Agric. Res.*, **69** (1): 57 – 62.
- Muralidharan, C. M. and M. S. Chari (1993):** Effect of spray deposition on bollworm (*Earias vittella* Stoll) damage and yield in hybrid 6 cotton. *Gujarat Agric. Univ. Res. J.*, **18** (2): 65-70.
- Moursy, E. B. and A. C. Bartlett (1991):** Effect of 'pyriproxyfen' as juvenile hormone analogue on the reproductive potentiality of the pink bollworm, *Pectinophora gossypiella* (Saunders). *Bull. of the Entomol. Soc. of Egypt, Econ. Ser.*, **19**, 191-197.
- Nada, M.A. (1996):** Effect of some insecticides on susceptible and field strains of pink bollworm, *Pectinophora gossypiella* (Saund.) at Sharkia and Fayum Governorates. *Egypt. J. Appl. Sci.*, **11** (5): 356-361.

- Natarajan, N.; T. Kumaraswami; M. Balasubramanian; A. Regupathy and S. Jayaraj (1985):** Efficacy of moult inhibitors on cotton pests under field conditions. Behavioural and Physiological Approaches in Pest Management, 167-170.
- Obando, R. A.; S. D. Garcia; J. S. Rea and P. Dugger and D. Richter (1998):** Confirm 2F and tracer as a useful alternative for IPM against boll worm, tobacco budworm and beet armyworm in cotton in northern Mexico. Proceedings Beltwide Cotton Conferences, San Diego, California, USA, 5-9 January. 2: 1228-1230.
- Panickar, B.K.; Bharpoda, T.M.; Patel, J.J. and J.R. Patel (2003)** Ovicidal effect of botanical and synthetic insecticides on boll worms. Indian Journal of Entomology, 65(2): 292-293.
- Perimmer, T. R. (1979):** *Heliothis* spp. control on cotton with pyrethroids, carbamates, organophosphates and biological insecticides. J. Econ. Entomol., 72: 593-598.
- Ramalho, F. S. and F. M. M. Jesus (1989):** Evaluation of electrodynamic and conventional insecticides against cotton boll weevil and pink bollworm. International Pest Control, 31(3): 56-58.
- Ramalho, F.S. and J.V. Gonzaga (1991):** Methodology of the application of pyrethroids against cotton boll weevil and pink bollworm. Tropical Pest Management, 37 (4): 324-328.

- Rashad, Amira, M.; N. M. Abd El-Salam; Nagwa M. Hussein and M. A. Zidan (1993).** IGR,s as bioactive agents against the pink and spiny bollworms. Egypt. J. Appl. Sci., **8** (1): 303-315.
- Salem, M. S.; E. B. Maursy and S. A. Mohamed (1994):** Effect of sublethal doses of Deltamethrin and Pirimiphos-methyl on the histological aspects of production in the cotton bollworms, *Pectinophora gossypiella* (Saund.) and *Earias insulana* (Boisd.). Bull. Ent. Soc. Egypt, Econ. Ser., **21**: 269-276.
- Satyanarayana, C. and B. V. Patil (1999):** Evaluation of different sprayers in the management of insect pests of irrigated cotton. Karnataka J. of Agric. Sci., **12** (1-4): 64-68.
- Satyanarayana, C. and B. V. Patil (2000):** Evaluation of different spray volumes in the management of cotton bollworms under irrigated conditions. Karnataka J. of Agric. Sci., **13** (2): 448-450.
- Shaaban, Z. and S. M. F. Ibrahim (1993):** Inhibition of growth and development of the spiny bollworm by benzoyl phenyl ureas. Alex. Soc. Exch., **14**: 129-142.
- Simwat, G.S. and A.K. Dhawan (1992):** Efficacy of diflubenzuron and in combination with insecticides for control of bollworms on different varieties of upland cotton (*Gossypium hirsutum*). Indian J. of Agric. Sci., **62** (6): 424 -426.

- Snedecor, G.W. (1977):** Statistical method applied to experiments in agriculture and biology. Iowa State Press, U.S.A. 534pp.
- Sun, Y. P. (1950):** Toxicity index: an improved method of comparing the relative toxicity of insecticides. J. Econ. Entomol., **43** :45-53
- Vadivelu. S.; P.V.S.Rao and M. Balasubramanian (1986):** Efficacy of different types of sprays on the control of cotton pests. Pesticides, **20** (10): 33-34.
- Wang, G; K. Wang; X. Jiang; M. Yi; G. Wang; K. Y. Wang; X. Y. Jiang and M. Q. Yi (1999):** Effect of four insecticides of the third instar larvae of cotton bollworms with different resistance. J. Shandong Agric. Univ., **30** (2): 104 -108.
- Watson, W. M.; A. M. Rashad and N. M. Hussein (1986):** Potencies of certain insecticides against the pink bollworm, *Pectinophora gossypiella* (Saund.) as influenced by chemical control program in Egypt. Bull. Of the Entomol. Soc. of Egypt, Econ. Seri., **15**, 79-86.
- Weiland, R. T.; C. K. Witco; P. Dugger and D. Richter (2000):** Scouting for early instar fall armyworms in cotton and their control with Dimilin R (Diflubenzuron). Proceedings Beltwide Cotton Conferences, San Antonio, USA, 4-8 January, (2):996-999.
- Wolfenbarger, D. A.; D. J. Wolfenbarger; P. Dugger and D. Richter (1999):** Contact toxicity of diethylhydrazine and

diphenyl benzoyl urea insect growth regulators against beet armyworm, bollworm and tobacco budworm. roceedings Beltwide Cotton Conferences, Orlando, Florida, USA, 3-7 January, **2**, 1254-1257.

Xia Jing Yuan; Pan DengMing; Cui XueFen and Hu FaQing (1996): Studies on the effect of Larvin on the management of pesticide resistant bollworms. China Cottons, **23** (11): 12-14.

Zidan, Z. H. and M. I. Abdel-Megeed (1988): New approaches in pesticides and insect control. Arabic publishing House and Delivery, Cairo: 605 pp. (In Arabic Language).