







REFERENCES

REFERENCES

- Abbott, W.S. (1925):** A method of computing the effectiveness of an insecticide. J. Econ. Ent. 18 : 265-267.
- Abdallah, M.D; M.A. Kandil ;A.A., Farag(1986):** Isolation and identification of biologically active compounds from extracts of Minteena, Barnoof and Ullaig. Bull of the Ent. Soc. Of Egypt, Econ. Ser.; 15: 191-197.
- Abdel - Aziz, Shadia and E.A.,Omer (1995):** Bio Evaluation of *Dodonaea*, *Dodonaea viscosa* L. Jac 9 . Extracts on the cotton leaf worm, *Spodoptera littoralis* (Boisd.) As indicated by life table parameters. Annals. Agric. Sci. Ain. Shams Univ. ;Cairo , 40 (2), 891- 900.
- Abo -seda, S.A.B. and F.M., El – Sheikn (1994):** Toxicity and antifeeding activity of *Ambrosia maritime* extracts and its synergistic action when applied in combination with certain insecticides on *spodoptera littoralis* (Boisd.) J. Agric. Res. Tanta Univ.; 20 (4) : 776- 780.
- Aboutabl, E.A.; G., Abdelhakim, and F.A., Moharram, (1996):** A study of some pharmacodynamic actions of certain *Melaleuca* species grown in Egypt. Phyotherapy, Res. 10(4): 345-347
- Ahmed, A.A.I. (1995):** studies on the effect and mode of action of *Melia azedarach* L. extraction some lepidopterous insects Ph.D. Thesis, Fac. Agric. Al-Azhar univ.
- Ahmed, S.M.S. (2000):** Laboratory evaluation of some biological activity of *Ricinus communis* seed extracts against the rice weevil , *Sitophilus oryzae* L. Arab Univ. J. Agric. Sci. , Ain Shams Univ., Cairo , 8 (3) , 853- 861 .

- Ahmed, S.M.S. (2001):** Lupin seed *Lupinus termis* extracts as Grain protect ants against the rice weevil *Sitophilus oryzae* L. and the lesser grain borer *Rhizopertha dominica*. Egypt. J. Agric. Res., 79 (1).
- Ahmed, A. Barakat; A., Mohamed Kandil; S.M., Hosni , Fahmy and A., Makram (1985):** Naturally occurring insecticides: III-Isolation, identification and assessment of insecticidal properties of plant extracts. Bull. Fac. Of Agric., Univ. of Cairo, 36 (2).
- Ahmed, A.A.I.; G.H., Schmidt ; M., Breuer (1997):** The chinaberry fruit extract as bio insecticide against the black cut worm larvae . International conference on pests in Agric. at le corum, Montpellier, France. 3: 1163-1170.
- Ajaj, Mishra; N.K., Dubey ; Sajla singh; C.M., Chaturvedi (2002):** Biological activities of essential oil of *chenopodium ambrosiodes* against storage pests and its effect on puberty attainment in Japanese quail. National Academy Sci., Letters, 25 (516), 174-177.
- Al- Moajel, N.H. and S.M., Abd El Baki (2000) :** *Brassica rapa* (Rape) seed extracts as protect ants to wheat grains against the lesser grain borer *Rhizopertha dominica* (F.). Annals Agric. Sci., Ain shams Univ., Cairo, 45 (1) 353-362.
- Aly, M.M; S.M., Radwan ; Z.H., Zidan ; A., El Hammady and S.H., Abdel -All (2000):** Food consumption and antifeeding behavior of the cotton leaf worm larvae as affected by Eucalyptus plant extracts in laboratory.

Annals, Agric Sci., Ain shams Univ., Cairo, 45 (2) , 769-775.

- Amr, H.E.S. and Nagway (1995):** Impact of *Nerium oleander* leaves extracts on some physiological change of the cotton leaf worm *spodoptera littoralis* (Boisd). Bull. Ent. Soc., Egypt, 73 : 159-166.
- Anderson, p. and J., Lofqvist (1996)** Oviposition deterrents from potato wheat gram, larval frass, and artificial diet for *Agrotis segetum* (Lepidoptera : Noctuidae). Environmental, Entomol.; 25 (3): 653-658.
- Anderson, B.A.; R.T., Holman ; L., Lundgren and G., Stenhagen (1980):**Capillary gas chromatography of leaf volatile. A possible aid to breeders for pest and disease resistance. J. Agric . Food Chem. 28: 985-989.
- Anonymous, (1966):** International rules for seed testing proceedings of international seed testing Association, XXXI ,pp 49-9.
- Ansari, S.H.; M., Ali and J.S., Qadry (1993):**Essential oils of *Pistacia integerrima* galls and their effect on the central nervous system. International, J., Pharmacognosy. 31:2 89-95.
- Babu, P.B.S; M.S., Nair ; B., Sumitha (1997):** Antifeedant and insecticidal effects of some plant extracts against castor semilooper *Achaea janata* Linn. J. of Insect Sci., 10(2): 179-180.
- Bai, H. and G., Koshy (1999):** Yellow oleander (*Thevetia neriifolia* Jass.) a bio – antifeedant for epilachna beetle (*Henosepilachna vigintioctopunctata* L.). J., of Tropical, Agric., 37:1-2, 64-67.

- Batanouny, K.H.; M.N., Hussein and M.S.A., Abo El-Kheir (1991):** Responed *Zea mays* to temoral variation of irrigation and salinitg under farm conditions in the Nile, Egypt. Proc. Intern. Conf. Plant Growth. Drought and Solinty in the Arab region.
- Behera, D.D.; K., Baral and S.R., Mitra (2002):** Effect of plant extracts on rice weevil *Sitophilus oryzae* Linn. J. of interacademia , 6 (1) : 59-64 .
- Behera, U.K. and C.R., Satapathy (1996):** Screening indigenous plants for their insecticidal properties against *Spodoptera litura* Fb. Insect . Environment ;2 (2) 43-44.
- Bekele, A.J.; D., Obeng ofori and A., Hassanali (1996) :** Evaluation of *Ocimum suave* (Willd) as a sourace of repellents, Toxicants and protectants in storage against three stored product insect pests. International J. of pest Management, 42 (2) : 139 – 142.
- Bekele, A.J.; D., Obeng – ofri and A., Hassanali (1997):** Evaluation of *ocimum kenyense* (Ayobangira) as source of repellents , toxicants and protectants in storage against three major stored product insect pests .J. of Appl. Ent. .; 121 (3) :169-173.
- Breuer, M. (1993):** Biologische and Chemisch- analytische untersuchungenzur insektiziden wirkung von *Melia azedarach* L. Ph.D. Thesis, Hannover Univ., Germany, pp 129.
- Breuer, M. and G.H., Schmidt (1990):** studies on the effect of *Melia azedarach* extracts on *Spodoptera frugiperda* (

- J.E.Smith*) (Lepidoptera : Noctuidae). Mitt. Dtsch . Ges. Allg. Angew. Ent. 97: 419-429.
- Bhusan,T.K.; S.S., Ghatak (1991) :** Evaluation of some plant powders as protectants against rice weevil *Sitophilus oryzae* Linn. (Curculionidae : Coleoptera). Environment and Ecology; 9(1): 115 -117 .
- Chakraborty, S.K.; and S.K., Ghose (1988):** Efficacy of some plant materials against the rice weevil *Sitophilus oryzae* (L.) Curculionidae : Coleoptera . Environment and Ecology ; 6(4) 833-839.
- Champagne, D., M.B., Isman and G.H., Neil Towerd (1989) :** Insecticidal activity of phytochemical and extracts of the Ueliaceae pp. 95 – 109 In Arnason, J.T., Philogene B.J.R and Uarand P. (eds), insecticides of plant origin American chemical society Washington D.C.
- Chen, W.; M.B., Isman and S.F., Chin (1995) :** Antifeedant and growth inhibitory effects of the *limonoid toosendanin* and *Melia toosendan* extracts on the variegated cutworm, *peridromasaucia* (Lep.:Noctuidae). J. Appl. Ent . 119 : (367-370).
- Chien –Chung chen; Shu –jen Chang; Ling – Ian Cheng and R., Hou (1996):** Deterrent effect of the chinaberry extract on oviposition of the diamondback moth *Plutella xylostella* (L.) (Lep. :Yponomeutidae). J. Appl. Ent 120, 165-169.
- Cozzolino, F.; R., Fellous ; G., Vernin and C., Parkanyi (2000):** GC/MS Analysis of the volatile constituents of *calamintha nepeta* L. Savi ssp. *Nepeta* from Southeastern France. J. Essent. Oil Res., 12, 481-486.

- Ding – Hong Jian; Guo yuyuan; Wu – caihon; Ding –Hj; Guo yy and Wu –ch (1997): Isolation and identification of semi chemicals from carrot and behavioral responses in cotton bollworm moth. *Acta Ent. Sinica*; 40 (73-78).
- Disbrey, B.O. and J.H., Rack (1970): histological laboratory methods. E&S Livingstone Edinbergh and London.
- Dwivedi, S.C.; Seema , Garg and S., Garg (1997) : Screening of plant extracts for ovicidal effect on the rice moth *corcyra cephalonica* (stainton). *Pest, Management and Econ. Zoology*; 5 (1) :53-55 .
- El- Gengaihi, S.E.; I.A., Ismail and G., Atwa (1996): Anti insect properties of the *Antholyza ricngens* Andr. (Family : Iridacea) extract against the American cotton bollworm, *Helliothis armigera* (Hubn). *Lepidoptera – Noctuidae. J. of Egypt Ger. Soc. Of Zoll. 21, Entomology*, 107 -119.
- El-Khayat, E.F. (1985): Pest management agents in the extracts of some plants from family Euphorbiaceae. M.SC. Thesis, Fac. Of Agric. Moshtohor, Zagazig University.
- El-Lakwah, F.A.M.; A.A., Darwish and O.M., Khaled (1992) : Effectiveness of Dill seed powder (*Anethum graveolens*) on some stored product insect. *Annals of Agric. Sc., Moshtohor*, 30 (4).
- El-Lakwah, F.A.M.; O.M., Khaled and A.A., Darwish (1993): Laboratory studies on the toxic effect of some plant seed extracts on some stored product insects. *Annals of Agric Sc., Moshtohor*, 31 (1).

- El- Lakwah, F.A.; A.M., Abdel -latif and Z.A., Halawa**
(1998): Effect of *Conyza dioscoridis* leaf extracts on three major insects of stored products in Egypt. Egypt. J. Agric. Res. ,76 (3).
- El-Sayed, N.H; M.I., Mogahed ; A.A., Haron and T.J., Mabry** (1998) :Flavonoids and other constituents from *Bassia muricata* and their insecticidal activities. Revista . Latino Americana, de. -Quimica ; 26(3): 81-85.
- El -Sayed, N.H.; M.I., Mogahed ; A.A., Haron and T.J., Mabry** (1999): Flavonoids and other constituents of *Vernonia biopontani* and insecticidal and fungicidal activities , Revista – Lantino Americana –de . Quimica ; 27 (1): 5-8.
- El- Shaarawy, M.F.; A.G., Antonious ; G., Hegazy and L.A., Yousse** (1992): Reaction of feeding the cotton leaf worm, *Spodoptera littoralis* (Boisd.) on certain plant leaves. 1: antifeeding effect International symposium on crop protection. Mededeling. Van- deFaculieit – Landbouwwetenschappen – Rijksuiversiteil – Gent. 57(3A): 689-695.
- Essaway, G.S.; H.M., Sobhhy and H.A., El- Banna** (1995): The hypoglycemic effect of volatile oil of some Egyptian plants. Veterinary- Medical, J. Giza 43:2 167-172.
- Farag, R.S.; S.A.S., Hallapo ; F.A., Hawedi and A.I.I., Basyony** (1986) : Chemical evaluation of rape seed oil Fette. Sei Fen. Anstrichemittel ; 88 : 391-397.
- Farag, A.A.; M.K.H., El- shemy ; A.F., Shehata and Y.S., Mohareb** (1988) : Isolation and identification of like

- phermon compounds from Egyptian plants. *Annals of Agric. Sc., Moshtohor*; 26 (1).
- Farnworth, R.; W., Lou; R., Blomster and M., Gortman (1964)** : Pericylivine anew catharanthus alkaloid. *J. Pham. Sci.* 53 : 1558 – 1564.
- Figueriedo, A.C.; L.G., Barrosa, Pedro; S.S., Fontinha;A.J., Ooman and J.J.C., Scheffer (1994)**: Composition of the essential oil of *Artemisia L. Her.*, an endemic species of the *Maderia archipelago* . *Flav . Farg . J.*, 9 (5) :229-232.
- Finney, D.F. (1952)**: Probit Analysis. Cambridge University Press, London, 256pp.
- Gamal, E.S.; Abo El- Ghar; S., Mohamed ; Khalid and M.E., Tahany (1994)** : Effects of plant extracts on development and fecundity of *Agrotis ipsilon* . *Bull, ent Soc. Egypt, Econ. Se .*, 21 (171).
- Gamal, E.S.; E. Abo El- Ghar; Mohamed; Khalid and M.E., Tahany (1995)** : Some bio chemical effect of plant extracts in the black cut worm, *Agrotis ipsilon* (Hufnagel). *Bull. Ent. soc. Egypt. Econ. Ser.* 22, (85).
- Garcia, R.; L., Lemus; P., Rivera and S., Erazo (1997)**: Biological and chemical study of paico *Chenopodium chilense*, *Chenopodiaceae*. *J., Ethno pharmacology* 57:2 85-88.
- Geissman, T.A. (1961)** : Chemistry of flavonoids. 126 pergmon press co. London.
- Ghatak, S.S. and T.K., Bhusan (1995)**: Evaluation on the ovicidal activity of some indigenous plant extracts on bihar hairy caterpillar *Spilosoma oblique* (WK.)

[Arctiidae : Lepidoptera]. Enviroment and Ecology ;
13 (2) : 294- 296 .

- Guo – Xianru; Yuan – Guo Hui ; Fan – Cailing ; Zheng – Qiwei and Ma – Jisheng , (2001) :** Determination of chemical components of the odor with luring activity from withered black poplar leaves by gas chromatography – mass spectrometry. J. of Henan – Agric. Univ; 35 (1): 24-25.
- Guy, I. And G., Vernin (1996):** Minor compounds from *Cistus ladaniferus* L. essential oil from Esterel. 2. Acids and Phenols. J. Essent., Oil Res., 8, 455-462.
- Han, J. ;W.H., Lin ; R.S., Xu ; W.L., Wang and S.H., Zhae (1991):** Chemical constituents of *Melia azedarach* L. *Yaoxue Xuebae*, 26 (6) : 426 – 429 .
- Hanson, J.R. (1972):** Chemistry of terpenes and terpenoids. (Ed. A.A. Newman). Academic Press. New York.pp 155-206.
- Harish – chander; S.G., Kulkarni and S.K., Berry (1990) :** Acorus calamus rhizomes as aprotectant of milled rice against *Sitophilus oryzae* and *Tribolium castaneum*.J. of Food and Technology, India 27 (3) : 171- 174 .
- Harris, H.F. (1989):** Cited in Carleton *et al.* (1967) Carletons Histological Techniques. 4th Ed. Oxford Univ. Press, Ner York, Toronto.
- Hashem, H.O.; A.M., Kheirellah ; M.H., Swidan and W.E., Osman (1994) :** Effect of alcoholic fruit extracts of *Melia azedarach* L. on the ovaries, fecundity and fertility of cotton leaf worm moths J. Agric. Res., 39 (1) : 221-227.

- Hegazy, G.; A.G., Antonious ;M.F., ElShaarawy and L.A., Youssef (1992) :** Reaction of feeding the cotton leaf worm, *Spodoptera littoralis* (Boisd.) on certain plant leaves and the effect of the plant extract on 4th larval instar Mededeligen Vande Faculties Land louwwetenss chappen, Rijksuniver –siteit. Gent., 57 (3A): 697 -705.
- Helen, C.F. and Su, (1989) :** Laboratory evaluation of dill seed extract in reducing infestation of rice weevil in stored wheat. J. Ent. Sci 24 : 317- 32 .
- Hiroi, T.; Y., Miyazaki; S., Imaoka ; Y., Funae (1995):** Induction of hepatic P₄₅₀ in rat by essential wood and leaf oils. Xenobiotica 25(5): 457-567
- Jain, N.C. (1986):** Schalms veterinary Hematology. 4th Ed. Lea & Febiger, Philadelphia, USA.
- Kady ,M.M.; R.A., Hassan and A.A., Said (1982) :** Chemical and toxicological studies on saponins isolated from saponaria roots. J.Agric. Sci., Mansoura Univ., 7, 37-379 .
- Kavimani, S.; R.I., Karpagam ; B., Jaykar ; A.R., Ilango-Karpagam, (1997):** Anti inflammatory activity of volatile oil of *Posidiumguajava*. Indian, J. Pharmaceutical, Sci. 59:3 (142-144).
- Khan , M., and R.M., Shahjahan (1998) :** Evaluation of eucalyptus leaf extracts as repellent for adult *Sitophilus oryzae* (L.) [Col : Curculionidae] and attractant for *Callasabruchus chinensis* (L.) [Col: Bruchidae]. Bangladesh. J. of Ent.; 8 (1/2) 81 -88.
- Khattab ,M.M.; S.M., Ahmed ; E.H., Abdel – karim and E.F., El- Khaiat, (1988):** Effect of some Euphorbiaceae

- plants as antifeedants against *Spodoptera littoralis* (Boisd). *Annals of Agric Sc., Moshtohor* , 26 (1) .
- Khodair, A.I. ; F.M., Hammouda ; S.I., Ismail ; M.M., El-Missiry and Abdel – Azim (1993) :** Phytochemical investigation of *Thymus decussates* . L. 1- Flavonoides and Volatile oil *Univ . Sci .J.*, 13 (2) .
- Kinsella, J.E., (1966) :** Metabolic patterus of the fatty acids of *periplaneta Americana* during the eniberyonic development . *Can J. Biochemistry* 44 : 247 – 258 .
- Lermioglu, F.; S., Bagci ; S., Onderoglu ; R., Ortac and L., Tugrul (1997):** Evaluation of the long term effects of *Oleum origami* on the toxicity induced by administration of streptozotocin in rats. *J., Pharmacy and Pharmacology* 49:11 (1157-1161).
- Lewis, H. and C., Smith (1967):** Sugaralcohls in fungi and green plants. *Methods of detection and estimation* . *New phtol .* ; 66 , 185- 204. Referred to Harbrone, J.B. (1998). *Photochemical methods, A guide to modern techniques of plant analysis*, p 54 , New York , U.S.A.
- Lgnatowicz, S. (1997a) :** Screening of plant powders for repellency to the grain weevil, *Sitophilus granaries* L. [Coleoptera : Curculionidae] *Annals. Of Warsaw Agric. Univ., SGGW, Horticulture*; 18 pp: (13-19).
- Lgnatowicz, S., (1997b) :** Repellency of powdered plant materials of the ribbed melilot and the white melilot to the grain weevil and the rice weevil. *Progress in plant Protection*; 37 (2pp) : 46-49.
- Li yunshou ; Zou Hua ying ; Wang Luxiang ; Nai Zhu ; Li Wan Yi ; Na xiao Yan; Tnagshaozong and Yang**

- Yizhang, (2001):** Insecticidal activity of extracts from *Eupatorium adenophorum* against four stored grain insects. Ent., Knowledge ; 38 (3) ,pp 214-216.
- Lieken Jie, M.S.F. and C.Y.C., Chol (1991) :** Re – examination of the fatty acid composition of *Biota orientalis* seed oil by gas chromatography – mass spectrometry of the picoliny ester derivatives, J. of Chromatography , 543: (257- 261).
- Lorente, I.; M.A., Ocete ; A., Zarzuelo ; M.M.,Cabo and J., Jimenez (1989):** Bioactivity of the essential oil of *Bupleurum fruticosum*. J., Natural Products 52:2 (267- 272).
- Mahgoub, S.M. and S.M.S., Ahmed (1996) :** *Ricinus communis* seed extract as protectants of wheat grain against the rice weevil *Sitophilus oryzae*. Annals. Agric. Sci., Ain Shams Univ ., Cairo , 41 (1) , 483- 491
- Makanjuola, W.A., (1989):** Evaluation of extracts of neem (*Azadirachta indica* A. Juss) for the control of some stored product pests. J. of stored product Res.; 25 (4) : 231-237.
- Malek, M.A.; B., Parveen and D., Talukder (1996) :** Insecticidal properties of four indigenous plant extracts against adults of cr-1 strain of *Triblium castaneum* Herbst . Bangladesh , J. of Ent.; 6 (1/2) 7-11.
- Melkani, A.B.; G.C., Shan ; Rashmi –parihar; R., Parihar ;S.C., Bhattcharyya ; N., Sen and K.L., Sethi (1990):** *Mentha longifolia* subsp. himalaiensis a new species as a source of aroma chemicals. Proceedings of the 11th international congress of essential oils, fragrances

- and flavors. New Delhi, India . 12-16 November , vol, 4
chemistry – analysis and structure , 177-180.
- Michigan State University , (1983).** MSTAT-C: Micro-
computer Statistical Program, Version 2.0.
Michigan State University, East Lansing.
- Mogahed, M.I. A.H., and Mohanna (1992) :** Toxic effect of
Barnoof crude extract on developmental stages of
Spodoptera littoralis (Boisd.) Zagazig J. Agric .Res . vol
.19 (3) 1437- 1445.
- Mohamed, S. Fouad and A., Alaa ,Abdel – Alem, (1994) :**
Comparative efficacy of some plant oils against *Sitophilus*
granaries (L.) infesting wheat grain, Bull. Ent Soc.
Egypt. Econ. Ser. 21 (65).
- Mohamed, E. Khalil; E.S., Gamal ; Abo El- Ghar and M.E.,
Tahany (1994) :** The antifeedant and growth- disrupting
activity of plant extracts on the black cutworm.
Agrotis ipsilon (Hufn) [Lepidoptera : Noctuidae], Bull.
Ent. Soc. Egypt. Econ. Ser., 21 (135).
- Mohapatra, S.; S.K., Sawarkar and H.P., Patnaik (1996) :**
Comparative bioassay of different solvent extracts of
neem seed kernels against *Sitophilus oryzae*, L. in rice.
current. Agric. Res. , 9(½) 89-93.
- Nagwa, Y.; A., Salem, Hoda; Ramadan and A., Elham
Sammour, (2003):** Physiological and histopathological
effect of some wild plant extracts on the cotton leaf
worm *Spodoptera littoralis* (Boisd), (Lepidoptera :
Noctuidae), Bull. Ent. Soc. Egypt, Econ. Ser., 29 (113).

- Niber , B.T.; J., Helenius and A.L., Varis (1992) : Toxicity of plant extracts to three storage beetles (Coleoptera).J. of Appl. Ent.; 113 (2) 202- 208 .
- Obeng –Ofori, D. and C., Reichmuth (1997) : Bioactivity of eugenol a major component of essential oil of *Ocimum suave* (Wild) against four species of stored – product . International, J. of pest management vol. 43 (1) 89- 94.
- Peterson, G.S.; M.A., Kandil ; M.D., Aballah and A.A.A, Farag (1989) : Isolation and characterization of biologically. active compounds from some plant extracts. Pesticide , Sci .; 25 (4) :337-342.
- Peterson, C.; A., Fristad ; Tsao Rong and J.R., Coats (2000) : Osajin and comfier two isoflavones purified from *osage orange* fruits tested for repellency to the maize weevil [Coleoptera : Curculionidae] . Environmental Ent.; 29 (6) : 1133- 1137.
- Parakash, A.; L.C., Pasalu and K.C., Mathur (1982): Evaluation of plant products as grain protection in paddy stroge. Int. J. Ent., India, 1 (1): 75-77.
- Prakash, A.; J., Rao ; S.P., Gupta and J., Behra (1993) : Evaluation of botanical pesticides as grain protectants against rice weevil, *Sitophilus oryzae* Linn. Botanical, pesticides. In, integrated pest management ; (360- 365).
- Rahman, M.L. ; M., Hosain and M., Ahmed (1999) : Repellent effects of urmoi , neem and turmeric extracts against rice weevil and granary weevil . Bangladesh J. of Ent.; 9 (12) : 9-16 .

- Roshdy, Tahany, A.Z. (1998):** Using some plant extracts and their mixtures with pesticides to control cotton pests. M. Sc. Thesis, Fac. Agric. Moshtohor, Zagazig, Univ.
- Roychoudhury, N. (1993) :** Antifeedant and insecticidal properties of some plant extracts for the adults of *Sitophilus oryzae* Linn. Curculionidae . Uttar , Pradesh, J. of Zool.; 13 (2) 97- 100.
- Ruberto, G. ; Biondi and M., Piattelli (1992):** The essential oil of Sicilian *Thymus capitatus* (L.) Hoffmanns J. Essential oil . Res., 4 (4) : 417 – 421.
- Rustaiyan, A.; A., Monfared and S., Masoudi (2001):** The essential oil of *Ferula flabelliloba* Rech. F. et Aell. J. Essent. Oil Res., 13, 403-404.
- Saleh, M.A.; M.M, El- Bolok ; K.A., Abdel salam and N.A., Ibrahim (1986) :** Plant extracts affecting insect feeding growth and metamorphosis. Bull. Fac. Agric. Cairo . Uni., 37 (1) : 526- 539
- Salem, S.A. (1991) :** Evaluation of neem seed oils Tuber protectant against *Phthorimaea operculella* Zell. [Lepidoptera : Gelechiidae]. Annals of Agric. Sc., Moshtohor, 29 (1).
- Salem, S.A. (1994) :** Physiological effects of *Melia azedarach* on the black cut worm *Agrotis ipsilon* (Huf.) Bull. Ent., Soc., Egypt ,27, (25).
- Sarac, A. and I., Tunc (1995) :** Residual toxicity and repellency of essential oils to stored – product insects . Rucktandstoxizital and abstossende wirkung von atherischen ole auf vorratsschadliche nsekten.Zeitschrift–

fur, Pflanzenkrankheiten und Pflanzenschutz;102(4):429-434.

Sattar, A.; M.S., Malik and S., Khan (1991) : Essential oils of the species of labiatae . Part IV . Composition of the essential oils of *Thymus serpyllum*. Pakistan J., Sci. Industr . Res., 24 (4) :119-120

Schafer, M. and O., Schimmer (2000): Composition of the essential oils from flowers, leaves and stems of *Grindelia robusta* and *G. squarrosa*. J. Essent. Oil Res., 12 553-555.

Schmidt, G.H.; A.A.I., Ahmed and M., Breuer (1997) : Effect of *Melia azedarach* extract on larval development and reproduction parameters of *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (Hufn.) [Lep. : Noctuidae]. Anzeiger. Fur, schadlingskunde, Pflanzenschutz , umweltschutz ; 70 (1) : 4-12.

Schmidt, G.A.; H., Rembold ; A.A.I., Ahmed and M., Breuer (1998): Effect of *Melia azedarach* fruit extract on juvenile hormone titer and protein content in the hem lymph of two species os noctuidae Lepidoptera larvae (Insecta : Lepidoptera : Noctuidae). Phytoparasitica ; 26 (4) :283-292.

Sharaby, Azeza (1988) : Evaluation of some Myrtaceae plant leaves as protectants against the infestation by *Sitophilus oryzae* L. and *Sitophilus granaries* L. Insect – Sci . and its Application; 9 (4) 465- 468.

Sharaby, Azeza (1989) : Some Myrtaceae leaves as protectants of rice against the infestation of *Sitophilus oryzae* L. and

- S. granaries* L. (Coleoptera). Polskie, Pismo, Ent.; 59 (2) : 377- 382.
- Sharaby, Azeza and N., Ammar (1997) :** Biological activity of extracts of *Tephrosia nubica* (Boiss) Baker against *Spodoptera littoralis* (Boisd.) and *Agrotis ipsilon* (Hufn.) .Tropenlandwirt ; 98 (October) : 143- 150.
- Sharma, R.K. (1999) :** Efficacy of neem products against rage pests in maize. Annals of Agric., Res .; 20 (2) :198-201.
- Shellard, E.J. (1957):** Practical plant chemistry. Londen Ritman Medical Publishing Co. Ltd.
- Shukla, R.M.; G., Chand and M.L., Saini (1992):** Laboratory evaluation of effectiveness of edible oils against three species of stored grain insects. Plant, Protection, Bull., Faridabad ; 44 (1/2) 14-15.
- Singh,S.; S., Puri ; H.M., Dani and R., Sharma (1997) :** Studies on the protective properties of Garlic oil against acetaminophen- induced hepatotoxicity in the rat, Indian, J. Pharmaceutical, Sci. 59:5 (259-262).
- Snodgrass, R.E. (1935):** The abdominal mechanisms of a grass hopper, Ibid., 94.
- Stahle, E. (1969) :** Thin -l ayer chromatography, springer, verlag . Berlin, Heidelberg, New York, P. 439, 441, 856 and 873.
- Steinmetz, M.D.; P., Joanny; Y., Millet and F., Giannallini (1985) :** The effect of essential oils of sage *Thuja* and *Hyssop* and some their constituents on the respiration of slices of cerebral cortex in vitroplants. Medicinalset, Phytotherapie, 19:1 (35-47).

- Su, H.C.F. (1990):** Biological activities of hexane extract of piper cubeba against rice weevils and cowpea weevils [Coleoptera : curculionidae] J., of Ent., Sci., 25 (1) : 16-20.
- Su, H.C.F. (1991):** Toxicity and repellency of Chenopodium oil to four species of stored -product insects . J., of Ent., Sci , 26 (1) : 178-182.
- Talukder, F.A. and P.E., Howse (1994) :** Laboratory evaluation of toxic and repellent properties of the pithraj tree, *Aphanamixis polystachya* walloparker, against *Sitophilus oryzae* L. International. J., of pest management, 40 (3) : 274- 279.
- Talukder, D.; M.A., Malek; L.A.M., Khanam and K.C., Dey (1998) :** Toxicity of some indigenous plant seed oil against *Tribolium confusum* Duval . [Coleoptera : Tenebrionidae] adult. Pakistan, J. Zool., 30 (4): 331-334.
- Thakur, .S.S (1997) :** *Rumex nepalensis* , a biopesticide against black cut worm , *Agrotis ypsilon* (Rottenburg). Insect . Environment ; 3 (2) 34- 35.
- Trematerra, P. and V., Lanzotti (1999) :** The activity of some compounds extracts by Allium on stored product insects *Oryzaephilus surinamensis* L. , *Sitophilus oryzae* L. and *Tribolium castaneum* (Herbst) Anzeiger. Fur, Schadlingskunde ; 72 (5) : 122- 125 .
- Tulyabaev, E.; K.H., Kushiev ; L.K., Abdullaeva and D.N., Dalimov (1995):** Alkaloids and thrieposphorylated derivatives as regulators of the activity of catalytic

- cholinesterase insect. Chem. Of Natural, Compound ; 31
(2) : 183- 187.
- Van Kampen, E.J. and W.G., Zijlstra (1961):** colorimetric
determination of haeglobin. Clin. Chem. Acta, 6 :538.
- Vilaseca, A.; H., Guinaudeau; B., Charles; I., Guy; A.R.D.,
Arias and A., Fournet (2004):** Chemical composition
and insecticidal activity of *Hedeoma mandoniana*
essential oils. J. Essent. Oil Res., 16, 380-383.
- Wiseman, B.R.; R.C., Gueldner; R.E., Lyuch ; and R.F.,
Serverson (1990):**biochemical activity of centipede
grass against fall army worm larvae . J. Chem. Ecol., 16
(9): 2677-2690.
- Wyatt, G.T. and M.L., Pan (1978):** insect plasma Proteins,
Ann. Rev. Biochem. 47-779.
- Xie, Y.S.; P.G., Fields and M.B., Isman (1995a) :** repellency
and toxicity of azadirachtin and neem concentrate to
three stroed product beetle. J. of Econ. Entomology.
88:4, 1024-1031.
- Xie, Y.S.; P.G., Fields; M.B., Isman; W.K., Chen and X.,
Zhang (1995b):** Insecticidal activity of *Melia toosendan*
and toosendanin against three stored product insects. J.,
of stored products, Res. 31:3, 259-265.
- Yang, S.L.; M.F., Roberto; M.J., O Neill ; F., Bucar and
Phillipson (1995) :**Flavonoids and chromenes from
Artemisia annua . Phytochemistray, 38 (1): 225-257.
- Yogita- Lohra; P.M., Singhvi ; Mamta- Panwar; Y., Lohra ;
M., Panwar (2001):** Effect of certain plant extracts on
oviposition of rust- red flour beetle, *Tribolium*
castaneum. J. of Applied Zoo, Res., 12(1): 67-70.

Zidan, Z.H.; A., El-Hammady; S.M., Radwan ; S.M., Abdelp-All and M.M., Aly (2000): enzyme and biochemical responses of the cotton leafworm larvae to the tested Eucalyptus plant extracts in laboratory. Annals Agric. Sci., Ain Shams Univ., Cairo, 45(2),