

VI- REFERENCES

- Abbott, W. S. (1925):** A method of computing the effectiveness of an insecticid.
J. Econ. Ent., 265-267.
- Abd-Elhafiez, A. M. (1974):** survey and biological studies on the mite *Tenuipalpus granati* Sayed infesting grapevine in Giza.
M. Sc. Thesis, Fac. Agric., Cairo Univ., 83pp.
- Abo-Taka, S. M. (1996):** Studies on some ecological factors affecting phytophagous and predacious mites in grapeyards.
Control Straegies for plant-feeding mites, Ohio, U.S.A. 4:177-180.
- Abou-Awad, B. A. (1976):** Incidence of eriophyid mites in Egypt.
Ph. D. Thesis, Fac. Agric., Cairo Univ., 196pp.
- Abou-Awad, B. A. ; El-Sawaf, B. M.; Reda, A. S. and Abdel-Khalek, A. A. (2000):** Environmental management and biological aspects of the two-eriophyoid fig mites *Aceria ficus* (Cotté) and *Rhyncaphytoptus ficifoliae* Keifer in Egypt.
Anzeiger- Fur-Schadlingskunde, 73(1), 5-12.
- Abou-Donia, M. G. (2002):** Ecological and Biological studies on some mites associated with some fruit trees.
Ph.D. Thesis Fac. Agric., Al Azahar. Univ., 186pp.
- Al-Mallah, N. M. and Mohammed, M. A. (1989):** Biological of the fig leaf mite *Rhyncaphytoptus ficifoliae* Keifer (Acarifomes: Rhyncaphytoptidae).

- Arab, Journal of plant protection, 7(1), 23-29.
- Altincag, R. and Akten, T. (1993):** Insect pests in grapevine nurseries and remedies in Aegean region: problem and their Solution. Bitk-Koruma. Bulteni. Turkey, 33, 3-4.
- Amos, J.; Hatton, R.G. and Massee, A.M. (1927):** Experiments in the transmission of reversion black currents. Ann. Rept. East, Malling Research Sta., Kent, 126-150.
- Arias-Giralda, A. and Nieto-Calderon, J. (1983):** Estimate of the losses caused by the common yellow mite *Tetranychus urticae* Koch in Tiertade Barros. Boletin- del-Servicio-de-Defensa- contra-plagas -e- Inspeccion- Fitopatologica, 9(3), 227-252.
- Arias-Giralda, A. and Nieto-Calderon, J. (1985):** The scab mite (*Brevipalpus lewisi* Mc Gregor), a new pest of grapevine in Spain. Fitopatologica, Spain, 11(2), 193-203.
- Arias-Giralda, A.; Baillod, N; Corino, L. Kreiter, S.; Lauren, J. C.; Schruft, G; Valentin, G. and Villa, Y. (1988):** The mite pests of grapevine. Servicio, Spain, 111: 25pp.
- Attiah, H.H. (1967a):** Two new species of mites on figs from Egypt (Acarina). Bull. Soc. Ent. Egypt, 51: 1-5.
- Attiah, H.H. (1967b):** The genus *Eutetranychus* in U. A. R. with description of three new species (Acarina: Tetranychidae).

Bull. Soc. Ent. Egypt, 51: 11-16.

Bawden, F.C. (1950): Plant viruses and virus disease,
3rd, Chronica Botanica Co., Waltham, Mass.

Boller, E. F.; Remund, U. and Candolfi, M. P. (1988): Hedges
as potential sources of *Typhlodromus pyri* Scheuten,
the most important predatory mite in vineyards of
Switzerland.

Entomophaga, 33 (2) 240-255.

Bradley, R. H. E. (1964): Aphid transmission of stylet borne
viruses:

148-174 In plant virology.

Buchanan, G. A; Bengston, M. and Exley, E. M. (1980):
Population growth of *Brevipalpus lewisi* McG.
(Acarina: Tenuipalpidae) on grapevines.

Aust. J. Agric. Res., 31 (5): 957-965.

Castagnoli, M. and Liguori, M. (1985): First observations on
the behavior of *Kampimardomus oberrans* (Oud.),
Typhlodromus exhilaratus Ragusa an *Phytoseius*
plumifer (Cnn. &Fanz.) (Acarina: Phytoseiidae) on
grapevine in Tuscany.

Redia, Italy, 68: 323-337. Advances in Acarology, 11:
95-113.

Castagnoli, M. and Liguori, M. (1986): Further investigations
on the mite fauna of the vine in Tuscany.

Redia, Italy, 69: 257-265.

Conner, R. L.; Thomas, J. B. and Whelan, E. D. (1991):
Comparison of mite resistance for control of wheat
streak mosaic.

Crop. Science (USA). (May-Apr 1991). 31(2), 315-318.

Corino, L. and Duverney, C. (1989): Geographic distribution of different species of phytoseiids (Acarina: Phytoseiidae) and Tydeids (Acarina: Tydeidae) in vineyards of the Aosta Valley.

Rivista di Viticoltura e di Enologia, 42 (3): 13-17.

Credi, R. (1998): Mite transmission of fig mosaic disease agent to periwinkle.

Journal of Submicroscopic Cytology and Pathology 28: 345-355.

Dellei, A. and Szendrey, L. (1988): The occurrence of phytophagous and predatory mite species in the vineyards of the country Heaves.

Novenyvedelem, Hungary. 24(3): 112-116.

Dennil, G.B. (1986): An ecological basis for timing control measures against the grapevine bud mite *Eriophyes vitis* Pgst.

Crop Protection, South Africa, 5(1): 12-14.

Dhar, V. and Rathore, Y.S. (1995): Pattern of population distribution of *Aceria cajani* on pigeonpea plant.

Indian-Journal of Pulses Research, 7: 2, 137-143.

Duso, C. and Liguori, M. (1984): Research on grapevine mites in the Veneto: Faunistic aspects and impact of pesticide treatments on the population of phytophagous and predacious mites.

Redia, Italy, 47: 337-353.

- Duso, C. and Ren, L. (1997):** Further investigations on the grapevine mite found in the Veneto region, North-Eastern, Italy.
Rivista-di-Viticoltura-e-di-Entologia, Italy, 50(2): 11-28.
- Ebrahim, H.M.; Abd El-Samed, M. A. and Fawzy, M.H. (2001):** Occurrence and seasonal distribution of mites associated with grapevine in Egypt.
Al-Azhar J. Agric. Res., 33: 341-350.
- El-Adel, F. E. R. (1982):** Ecological and biological studies on certain Eriophyoid mites in Lower Egypt.
M. Sc. Thesis, Fac. Agric. Ain Shams Univ., 111pp.
- El-Benhawy, E.M.Z. (1967):** Biological studies on predacious mites.
M. Sc. Thesis. Fac. Agric., Ain Shams Univ., 94pp.
- El-Duweini, F.K.; Sawires, Z.R. and Ibrahim, S.M. (1989):** Biological studies on *Tydeus californicus* (Banks) to evaluate its feeding habit.
3rd Nat. Conf. Of Pests & Ismailia, Egypt.
- El-Halawany, A. S. H. (2001):** Studies on some mite species infesting some fruit trees.
M. Sc. Thesis Fac. Agric., Zigzag Univ. 165pp.
- El-Halawany, M. E. and Abdel –Samad, M. A. (1990a):** Population dynamics of the spider mite *Tetranychus arabicus* Attiah and its predatory mite *Phytoseius finitimus* Ribaga in Fig orchard.
Agric. Res. Rev., 68 :25- 29.
- El-Halawany, M. E. and Abdel –Samad, M. A. (1990b):** Three new phytoseiid species.

Agric. Res. Rev., 68 :87- 96 .

El-Halawany, M. E.; Ibrahim, G. A. and Abdel –Samad M. A. (1990): Ecological aspects of *Eriophyes ficus* Cotté and its mite *Phytoseius finitimus* Ribaga in Fig orchards.

Agric. Res. Rev., 68: 49 – 57.

El-Halawany, M. E.; Kandeel, M. H. and Rakha, M.A. (1986): Mites inhabiting deciduous fruit trees.

Agric. Res. Rev., 64 (1) : 115 –122.

El-Komi, G. H. A. (2000): A study of the effect of some pesticides and their combination the susceptibility of *T. urticae* Koch in Gharbia and Kafr El Sheikh Governorates.

M. Sc. Thesis, Fac. Al-Azhar Univ., 126pp.

El-Sebae, A. H.; El-Doksh, A. A.; Marei, A.S. and Mostafa, F. I. (1978): Toxicity of certain pesticides and their combination against the red mite species *Tetranychus cinnabarinus* Alex.

J. Agric., Res., 26(2): 453-460.

English-Loeb, G. M.; Flaherty, D. L.; Wilson, L. T.; Barnet, W.W.; Leavit, G. M. and Settle, W. E. (1986): Pest management affects spider mites in vineyards.

Calif. Agric., U.S.A., 40(3-4): 28-30.

Flechtmann, C.H.W. and Moraes, GJ. De (1991): Two new species of spider mites (Acarina: Tetranychidae) from Brazil.

Enternational- Jouranal of Acarology, 17 (3): 181-185.

- Frost, W. E.; Eagling, D. R. and Manson, D. C. M.(1990):**
Abacarus hystrix (Nalepa) (Acarina: Eriophidae)
 Newly recorded in Australia.
 Journal of the Australian Entomology Society. 182: 3-29.
- Gall, D. L. E. (1992):** The mites at Charentes.
 Paysan Francis, 875: 16-22.
- Gonzalez, R. H. (1983):** The false grape mite *Brevipalpus chilensis* Baker (Acarina: Tenupalpidae).
 Revista Fruticola Chile, 4(2): 61-65.
- Gonzalez, M. Q. D. E. and Vilorio, Z. (1991):** *Tetranychus urticae* Koch and *Oligonicus bagdasariani* Baker & Pritchard (Acarina: Tetranychidae) phytophagous mites of importance on grapes (*Vitis vinifera* L.)
 in Revista de la Facultad de Agronomia, Univ. de Zullia, 8 (1) : 1-14.
- Gutierrez, J; Etienne, J. (1986):** Tetranychidae of Reunion Island and some of their predators.
 Agronomie- Tropicale, 41(1): 84-91.
- Gyorffy, J. M. (1994):** New acaride in grapevine protection,
 Novenyvedel. 30 (9): 428-431.
- Haji, F. N. P.; Moraes, G. J. Lacerrda, C. A. and Neto, R. S. (1988):** Chemical control of tomato russet mite, *Aculops lycopersici* (Massee, 1937).
 Anais -de- Sociedade- Entomologica- de-Barazil. 17(2): 437-442.
- Halawa, A. M. (1998):** Studies on some mites associated with fruit trees,
 M. Sc. Thesis, Fac. Agric., Zagazig Univ., 113pp.

- Hanna, R; Wlison, L. T.; Zalom, F. G.; Flaherty, D. L. and Leavitt, G. M. (1996):** Spatial and temporal dynamics of spider mites (Acari: Tetranychidae) in Thompson seedless vineyards.
Environ. Entomol. U.S.A. 25(2): 370-382.
- Hardman, J. M.; Rogers, R. E. and Maclellan, C. R. (1988):** Advantages and disadvantages of using pyrethroids in Nova Scotia apple orchards.
J. Econ. Entomol. 8(6): 1737-1749.
- Hassan, H. A.; El-Kady, M. H. and Kodirah, S. M. (1967):** On the control of citrus mite, *Phyllocoptruta oleivora* on citrus.
Agric. Res. Rev. 45 (2): 181-184.
- Hatzinikolis, E. N. (1986):** The genus *Tenuipalpus* (Acari: Tenuipalpidae) in Greece Hellenica, 4(1): 19-22.
- Heikal, I. H. (1977):** Ecological and Biological studies on mite occurring on some deciduous fruit trees.
Ph.D. Thesis, Fac. Agric., Al-Azhar Univ., 109 pp.
- Heikal, I. H.; Ibrahim, S. M. and Iskander, N. G. (1996):** Ecological and Biological studies on mite occurring on Fig trees at Brrage District (Qalubiya Governorate).
Egypt. J. Agric. Res., 74 (4): 919- 927 .
- Huchy, M. and Posisil, Z. (1992):** Damage and economic injury level of eriophyid and tetranychid mites on grapes in Czechoslovakia.
Exp. & App. Acarol., 14(2): 95-106.
- Huchy, M.; Pospisil, Z. and Zacharda, M. (1991):** Phytophagous and predatory mites (Acarina:

Tetranychidae, Eriophyidae Phytoseiidae and Stigmaeidae) in South Moravian vineyards, Czechoslovakia, treated with various types of chemical.

Exp. & App. Acarol., 13(1): 41-52.

Inoue, K.; Osakabe, M.; Ashihara, W. and Hamamura, T.

(1986): Factors effecting on abundance of the Kanzawa spider mite, *Tetranychus kanzawi* Kishida in glasshous and influence of pesticidal application on occurrence of the Kanzawa spider mite and its predator.

Bulletin fruit trees Res. Station Japan. 6:103-116.

James, D. G. and Whitney, J. (1991): Biological control of grapevine mites in Inland, Australia and New Zealand wine Industry.

Journal, 6 (3): 210-214.

Jeppson, L. R.; Keifer, H.H. and Baker, E. W. (1975): Mites injuring to economic plants. Barkeley, California, U.S.A., Univ. of California, 96, 437, 438, 517 and 521 pp.

Jubb, G. L.; Masteller, E. C. and Lehman, R. D. (1985): Survey of arthropods in vinyards of Erie country, Pennsylvina.

Acari. Inter. Jour. Of Acarolo. 11(3): 201-207.

Koleva, R; Ferenczy, A. and Jenser, G. (1996): Effect of various plant protection programs on mite poulations in vinyards.

Hortic. Sci, Hungary, 1(2): 79-82.

- Kozar, F. (1989):** Microhabitat specialization and similary of scale insects assemblages on different fruit trees and in different countries.
Ecological Ento., 14 (2): 175-180.
- Laffi, F.; Raboni, F. (1994):** *Aceria tulipae* (Keifer) Eriophyid mite dangerous on garlic plants.
Informatore Fitopatologica. Italy. 44(9): 38-40.
- Liguori, M. (1987):** Population trends of phytophagous and predatory mites in two vinyards of Chiavti area.
Redia, Italy, 70: 141-149.
- Lozzia, G. C.; Nepomuceno, R. and Rancati, M. A. (1984):** Presence and distribution of Acari; Phytoseiidae in some Lombardy vinyards.
Vignevisini, Italy. 11(11): 31-35.
- Lozzia, G.; Rigamonti, I. E. and Blaise, P. (1998):** The effect of vegetation mangement on the populations of phytoseiidae in vinyards of Lombardie. Intergr. Cont. in Viticul.
Proceedings of the Meeting at Goldollo, Hungary, 21(2): 73-74.
- Mali, A. R.; Gandhale, D. N. and Path, A. (1983):** Heavey incidence of two spotted spider mite *Tetranychus urticae* Koch (Acarina: Tetranychidae).
Revista Fruticola Chile, 4(2): 61-65.
- Malison, M. (1996):** Influence of the measurement unit of populations in density dependent fecundity studies of the predatory mite *Amblyseius aberrans* (Oud.) (Acarina: Phytoseiidae).
J. of Appl. Entomol., Italy. 120(1): 1-6.

- Maramorosch, K. (1964):** Aphid transmission of stylet borne viruses:
148-164 In plant virology.
- Mattews, R.E.F. (1991):** "Plant virology". 3rd Ed.
Academic Press Inc. 855 pp.
- Maxiner, M. (1990):** Investigations on insecticide resistance in the predatory mite, *Typhlodromus pyri* Schuten (Acari: Phytoseiidae) on grapevines in the vine region Mosel-Saar-Ruwr. Mitteilungen aus der Biologischen Bundesanstalt für Land-und Forstwirtschaft Berlin-Dahlem, Germany, 257,118pp.
- Mohd, Y.; Sahader, C. and Yousuf, M. (1998):** Infection of four species of Eutetranychus on forest tree in Rajasthan.
Annals of Forestry 6: 1, 79-83.
- Moraes, G. J.; McMurtry, J. A. (1983):** Phytoseiid mite (Acarina) of northeastern Brazil with description of four new species.
International Journal of Acarology. 9 (3): 131-148.
- Nachev, P. (1976):** Studies on eriophyid mites in Bulgaria IX. Eriophyid mites on pear trees (Acarina : Eriophyidae)
Rastit. Nauka, 3: 70-76.
- Naeem, A. and Akhyani, A. (1988):** The fig souring disease in kashan and its transmission factors. Entomology- et-phytopathologia- Appliquees. 55. 1-2, 7-8, 31-42.

- Nemoto, H.; Kobayashi, M.; Osawa, T.; Yamashita, S. and Doi, Y. (1980):** Two eriophyid mites injurious to fig and chrysanthemum in Japan.
Japanese J. Appl. Ent. & Zool., 24 (2): 49-53.
- Nicotina, M. and Cioffi, E. A. (1997):** Population densities of phytoseiid mites in the vineyardes of Lazio, Campania and Basilicata.
Informatore Agrario, Italy, 53 (19): 59-62.
- Nitta, H.; Imada, J.; Kano, T. and Ogasawara, S. (1995):** Occurrence and causes of fig mosaic symptoms in Hirshima prefecture. Bulletin of the Hiroshima prefectural.
Agriculture- Res. Center. 62, 53-65.
- Ocete, R. and Shukhrava, M. (1996):** Note on the presence of *Arthrocnodax Rubsaaman* (Diptera) in wild grapevine.
Boletin- de- sanidad- vegetal, Plages. Spain, 22(2): 465-468.
- Orlob, G.B. (1968):** Relationships between *Tetranychus urticae* Koch and some plant viruses.
Virology 35: 121-133.
- Ozar, A. I. Onder, P.; Saribay, A.; Ozkut, S. Gundodu, M.; Azeri, T.; Arinc, Y. and Genc, H. (1986):** Investigation on improving the measures for controlling pests and diseases occurring on fig Agean region.
Doga, 10 (2): 263-277.
- Papaioannou, S. P.; Markoyiannaki, P. D.; Rumbos, I. and Adamopoulos, I. (1999):** Phytoseiid mites associated

- with vine in various province of Greece : a contribution to faunistics and biogeography, with reference to eco-ethological aspects of *Phytoseius finitimus* (Ribaga) (Acari: Phytoseiidae).
Acarologia, Greece, 40(2): 113-125.
- Peralta, A.L. (1993):** New records of eriophyid species (Acarina, Eriophyoidea) on Fig (*Ficus carica*):
Agricultura- Tecnica- Santiago, 53 (1): 101-103.
- Perez, M. L. and Moraza, M. L. (1997):** Study of *Typhlodromus pyri* Scheuten and its relationship with *Calepitrimerus vitis* (Nalepa) in Rioja vineyards.
Bulletin I ovi. Spain, 70, 801-802, 832-845.
- Plavisić, B. and Milicic, D. (1980):** Intracellular changes in trees infected with fig mosaic virus transmissible by mite *Eriophyes ficus* Cotté.
Acta Hort., 110 : 281 - 286.
- Pritchard, A. E. and Baker, E. W. (1958):** The false spider mires (Acarina: tenuipalpidae).
Univ. Calif. Publ. Ent., 14(3): 175-274.
- Rasmy, A. H. (1966):** Integrated control of citrus mite.
Ph. D. Thesis, Fac. Agric. Cairo. Univ. 128pp.
- Rasmy, A. H. and Abou-awad. B. A. (1972):** Mites inhabiting fig trees in Egypt.
Z. Ang. Ent. 70(3): 314-316.
- Rasmy, A. H. and El-Banhawy, E. M. (1974):** Behavior and bionomics of the predatory mite, *Phytoseius plumifer* (Acarna : Phytoseiidae) as affected by physical surface features of host plants.
Entomophaga, 19 (3): 255-257.

- Rasmy, A. H.; Zaher, M. A. and Abou-Awad, B. A. (1972):**
Mites associated with deciduous fruit trees in U. A. R.
Z.
Ang. Ent., 70: 179-183.
- Rather, A. Q. (1986):** New species and a new record genus
Amblydromella Muma (Acarina: Phytoseiidae) in
India.
Monthly Magazine. (1440/1443) 103-107.
- Rilling, G. (1989):** Differential response of grapevine cultivars
to European red mite, *Panonychus ulmi* Koch
elaboration of a screening method.
Vitis, Germany, 28 (2): 97-110.
- Rizk, G. A.; Karaman, G. A. and Ali, M. A. (1978):** Mite
populations on grapevine in middle Egypt.
Bull. Soc. Ent. Egypt, 62:1.
- Sanap, M. M.; Bendre, N. J. and Deshukh, R. B. (1989):**
Reaction of sterility mosaic resistant pigeonpea lines
to insect to pests.
International- Pigeonpea- Newsletter, 10: 29-30.
- Seiferas, D. L.; Harvey, T. L.; Kofoed, K. D. and Stegmeier,
D. (1996):** Natural infection of pearl millet and
sorghum by wheat streak mosaic virus in Kansas.
Plant disease (U.S.A.). 80(2): 179-185.
- Shehata, K.K. (1967):** Survey and biological studies of mites of
family: Phytoseiidae and its economic importance.
Ph. D. Thesis, Fac. Agric. Cairo. Univ., 371pp.
- Shibao, M. and Tanaka, H. (1997):** Overwintering sites and
stages of the fig bud mite, *Eriophyes ficus* Cotté.

Proceeding of the Kansai plant protection- Society.
Japan, 3(9), 1-3.

Silberschmidt, K. (1960): Types of photo virus Y cause
necrosis to tobacco, history and recent observation.
Ann. Potato. J. 37: 151-159.

Smith, K.M. (1951): "Alatant virus in sugar-beets and
mangolds".
Natura, 167, 1061.

Smith, K.M. (1972): Plant virus disease. 3rd Ed., Longman.
Group Limited, London, 684 pp.

Smith, D. and Papacek. D. R. (1991): Studies of predator
mites, *Tegolophus oustralis* and *Phyllocoptruta*
oleivora . Effect of pesticides. Alternative host plant
and augmenative relas. Experimental and Applied.
Acarology. 12 (3/4): 195-217.

Soliman, Z.R. (1966): A general survey of mites associated with
fruit trees in the U. A. R. with special refrence to
family Cheyletidae and its economic importance.
Ph. D. Thesis, Fac. Agric., Cairo Univ. 253pp.

Solva, J.; Zoschg, M.; Hluchy, M. and Zacharda, M. (1997):
Predatory phytoseiidae mite (Acari: Mesostigmata) in
vinyards and fruit orchards in southern Tyrol.
Anzeiger fur Schadlingsskunde, Pflanzenschutz,
Umweltschutz., Italy, 700 (1): 17-19.

Szendrey, L. and Bathory, G. (1995): Studies of over wintering
of mites in grapevine stands in the Eger and Martalji
wine growing regios.
Noven vedelem, Hungary, 31 (12): 595-599.

- Thomas, J. B. and Whelan, E. D. P. (1991):** Genetics of wheat curl mite resistance in wheat: recombination of Cmc1 with the 6D centromere.
Gron-viden- Havebrug. Denmark 65(6): 8pp.
- Thomsen, A.; Nielson, S. L. and Ramssusen, A. N. (1991):** Stem curl and black currant bud gall mite.
Gron-viden- Havebrug. Denmark 65(6): 8pp.
- Toros, S. (1983):** Mites transmitting plant Pathogenic Viruses.
Bitki, Koruma, Bulteni, 32(2): 74-91.
- Trapman, M. (1989):** Integrated control of red spider mite in fruit culture- Dutch experiences.
Besseres. Obst. 34(1): 8-10.
- Ueckermann, E. A. (1991):** African Aceria (Acari: Eriophyidae): species associated with plant of the family Moraceae
phytophactica, 23, (1): 1-7. 23
- Vila, Y. (1988):** The spider mite on the increase in the vineyards of Champagne and Burgundy.
Phytoma, France, 398: 34-35, 38, 40.
- Villabobos, S.G. and Espinosa, C.L. (1990):** Mites (Eriophyoidea) associated with the fig tree (*Ficus carica* L.).
Revista, Chapingo. 15: 67- 68, 142 -144; 4 ref.
- Villaronga, P.; Marques, J.; Casanovas, S. and Ferragut, F. (1991):** The phytophagous and predatory mites in vineyards of the Alt-Emporda (Gerona, Spain).
Progress Agricole et Viticil, 108(23): 519-523.
- Vogt, H.; Dickler, E. and Grauhan, H. (1990):** Effect of an unique application of Acaricides in apple orchards on

the population dynamics of *Panonychus ulmi* and *Aculus schlechtendali* with special regard to beneficials.

J. of Appl. Ent. 110(1): 35-54.

Wafa, A. K.; Zaher, M.A. and Yousef, A. A. (1969): Survey of the tenuipalpid mites in U.A.R. (Acarina: Tenuipalpidae).

Bull. Zool. Soc. Egypt, 22: 52-59.

Wahba, M. A. (1976): Studies on Tydeid mites (Tydeidae), especially as related to mango and citrus trees.

Ph.D. Thesis, Fac. Sci., Ain Shams Univ.

Watson, M.A. (1936): Factors affecting the amount of infection obtained by aphid transmission of the virus Hy.

III. Roy. Soc. London Phil. Trans. Ser. B. 226: 467-489.

Wermeliner, B.; Candolfi, M. P. and Boumgartner, J. (1992): A model of the European red mite (Acar: Tetranychidae) population dynamics and its linkage to grapevine growth and development.

J. App. Ent. 114 (2): 155-166.

Wildbolz, T.; Staub, A. (1986): Predatory mite settlement in fruit cultivation. Schweizerische Zeitschrift Fur Obst- und Weinbau,

Switzerland. 122: 483-488.

Wood, E.A.; Sebesta, E.E.; Webster, J.A. and Porter, D. (1995): Resistance to wheat curl mite (Acarina: Eriophyidae) in greenbug-resistant (Gaucha) triticale and (Gauch)X wheat crosses.

J. of Eco. Ent. (USA). 88(4): 1032-1036.

- Yousef, A.A. (1970):** Mites associated with vine trees in the U.A.R. (Acarina).
Z. ang. Ent., 65: 1-6.
- Zaher, M.A.; Wafa, A. K. and Shehata, K. K. (1969):** Life history of the predatory mite, *Phytoseius plumifer* and effect of nutrition on its biology (Acarina: Phytoseiidae).
Ent. Exp. Appl., 12:383-388.
- Zandigiacomo, P. and Coiutti, C. (1997):** Survey on the grapevine mite fauna in the area of Trieste. Revista di viticoltura e di Enologia,
Italy. 50 (1): 17-25.
- Zandigiacomo, P. and Frausin, C. (1998):** Entomological problem in grapevine rearing in Friuli- Venezia. Informatore- Fitopatologico,
Italy. 48: 7-8.
- Zhao-Jian; Kuag-Haiyuan (2000):** Three new species of the family Eriophyidae from China (Acari: Eriophyidae).
Acta- Zootaxonomica- Sinica, 25 (1) : 37-40.