




# REFERENCES

## REFERENCES

- Abdel-Tawab, F. M.; M. A. Rashed; S. A. Azer; G. M. I. El-Metabteb and A. F. Khafaga (2008):** Molecular fingerprinting and marker-assisted selection in citrus species. Egypt. J. Genet. Cytol., 37: 335-358.
- Adeishvili, N. I. and L. G. Kharebava (1987):** Volatile compounds of lemon leaves. Subtropicheskie Kultury, (6): 3-66.
- Agrarwal, s. G. Lal; R. K. Thappa ; B. K. Kapahi, and Y. K. Sarin(1989:** Seasonal Studies on Indian Citrus jambhiri Lush leaf oil-a new chemotype Flavour-and-Fragrance-Journal. 4:1,33-36:7 ref.
- Ahmed, E. M. (2004):** Biochemical studies on some Citrus varieties and the use of their volatile oils as antifungi. M. Sc. Thesis, Department of Agriculture Science, Institute of Environmental studies and research, Ain Shams University, 190 pp.
- Akkaya, M. S.; A. A. Bhagwat and P. B. Cregan (1992):** Length polymorphisms of simple sequence repeat DNA in soybean. Genetics; 132: 1131-1139.
- Attaway, J. A.; A. P. Pieringer and B. S. busliq (1968):** The origin of orange citrus flavor components. T.V. The Terpenes of Valencia orange leaf, peel and blossom oils. Phytochemistry. 7: 1695-8.
- Baaliouamer, A. and B. Y. Meklati (1986):** Analysis of bitter orange petit grain essential oil by combined gas chromatography-

mass spectrometry, Agricultural and Biological chemistry,  
50. (8): 2111- 2114.

-----; -----; **D. Fraisse and C. Scharff**  
(1988): Analysis of leaf oils from four varieties of Sweet  
orange by combined GC-MS. Flavor and Fragrance Journal.,  
3 (1): 47-52.

**Blanco T. C.; E. E. Stashenko ; M. Y. Combariza and I. R. Martinez**  
(1995): Comparative study of Colombian citrus oils by  
high-resolution gas chromatography and gas  
chromatography-mass spectrometry. Journal of  
Chromatography, Vol 697 (1/2): 501-513.

**Bukiya, T .G.; I. P. Mudzhiri and N. Z. Yakobashvili (1984):**Petit  
grain essential oil from industrial wastes. Maslo-Zhir Prom-  
st (5) 22. (C.F. Chem. Abst. Vol. 101, No. 136789 a).

**Brophy, J. J. ; R. J. Goldsack and P. I. Forster(2001):**The leaf oils of  
the Australian species Citrus (Rutaceae). Jouranal of  
Essential oil Reasearch,2001,Vol.13,No.4,PP.264-268,33 ref

**Cabrita, P. Elisiario, J. Leitao, A. Guerrerio, C. Dore, F. Dosba and**  
**C. Baril (2001):** Assessment of genetic relationships  
among citrus species and varieties by isozyme and RAPD  
markers. Theor. Appl. Genet., 101:860-864

**Calvarano, M.; F. Salntiro and T. Sacco (1982):**Essential oil from the  
leaves of the Tahiti lime (*Citrus Latifolia* Tanaka S. W.)  
from Brazilin. Essenze Deriv. Agrum. 52 (I): 52-8.

**Cao W; P. Hucl; G. Scoles; R.N. Chibber; P.N. Fox and B.**  
**Skovmand, (2002):** Cultivar identification and pedigree

assessment of common wheat based on RAPD analysis  
wheat Inf. Serv., 95:29-35.

**Charters, Y. M.; A. Robertson; M. J. Wilkinson and G. Ramsay**

**(1996):**PCR analysis of oilseed rape cultivars (*Brassica napus* L. ssp. *Oleifera*) using 5'-anchored simple sequence repeat (SSR) primers. Theoretical and Applied Genetics, 92: 442-447.

**Chu, J.; Z. Zheugju and P. F. Zhizwu (1988):**Chemical constituents of leaf essential oils in *Citrus limonia* Osbeck. Acta Botanica Sinica, Vol. 30 (2): 226-228.

**Dela Carmen, P. and J. C. Sardi (1977):**Study on bitter orange (*C. aurantium* L.) essential oil Arch, Bioquim, Quim Farm. 20, 69-72. C. F. Chem. Abst. Vol. 91 No. 181256 K.

**Della paporta, S. L.; Wood, J. and Hicks, J. B. (1983):**A plant DNA miniprep: Version II. Plant Mol. Biol. Rep. 1:19-21.

**De Pasquale, F. ; M. Siragusa; L. Abbate; N. Tusa; C. De Pasquale**

**and G. Alonzo (2006):**Characterization of five sour orange clones through molecular markers and leaf essential oils analysis. Scientia Horticulture, 109 : 54-59.

**El-Deeb, M. A. S.; S. M. Mohamed; A. M. H. El-Zahawy and E. A. El-Gamal (1993):** Effect of nitrogen sources and levels on the growth, seed yield and oil content of *Negilla sativa* L. plant. Egypt. J. Appl. Sci. 8 (6): 387-409.

**El-Samahy, S. K. A. Ashour and M. G. Abdel-Fadeel.(1982):** Quantitative analysis of some citrus peel oils. R. A. K.,

Riechst, Aromen, Kosmet, 32(2), 68-70. C.F. (chem.. Abst. Vol. 26, 1982. No. 205205.

**Fadel, H. H. M. (1991) :** comparison studies on leaf oils of egyption citrus varieties – journal of isamic Academy of Sciences 4-3: 196-199.

**Fang, D. Q.; M. I. Roose; R. R. Krueger and C. T. Federici (1997):** Fingerprinting trifoliate orange germplasm accessions with isozymes, RFLPs and inter-simple sequence repeat markers. Theoretical and Applied Genetics, 95:211-219.

**Fischer, A.; W. Grab and P. Schieberle (2008):**Characterization of the most odour-active compounds in a peel oil extract from Pontianak oranges (*Citrus nobilis* var. *Lour. microcarpa* Hassk.) Vol 227 (3):735-744.

**Guenther, E. (1960):**The essential oils 4th ed, Vol. I, II. D. Van Nostrand Co., INC., New York.

**Gurib-Fakim, A. and F. Demarne (1995):**Aromatic plants of Mauritius volatile constituents of the leaf oil of *Citrus aurantium* L., *Citrus paradise* Macfad and *Citrus sinensis* (L.) Osbeck. Journal of essential oil Research, Vol. 7 (1): 105-109.

**Hognadottir, A. and R. L. Rouseff, (2003):**Identification of aroma action compounds in orange essence oil using gas chromatography – olfactometry and gas chromatography – mass spectrometry. Jouranal of chromatography A, 998 ;201-211.

**Jaccard, P. (1980):**Nouvelles rechearches sur la distribution locale. Bull. Soc. Vaud. Sci. Nat. 44: 223-270.

- Kantety, R. V.; X. Zhang; J. L. Bennetzen and R. Z., Zehr (1995):**  
Assessment of genetic diversity in dent and popcorn (*Zea mays L.*) inbred lines using inter-simple sequence repeat (ISSR) amplification. Mol. Breed. 1: 365-373.
- Karawya, M.S. ; S. I. bulbaa and M.S. Hifnawy (1971):**Study of stem distilled peel and leaf oils of certain citrus species growing in Egypt. U.A.R. J. pharm. Sci., 12 (2) 361-70.
- **and M.S. Hifnawy (1977):**Leaf essential oils three different varieties of *Citrus reticulata* Blanco growing in Egypt. Perfumer and Flavorist, 2 (5), 60.
- Kekelidze, N.A.; M.I. Dzhanikashvili and V. V. Kutateladze (1981-a):**  
Petitgrain oil from the mandarin variety unshu. Maslo-zhir. Prom-st. (2) 35-6. C.F. Chem. Abst. Vol 94, No 162583.
- ; ----- **and ----- (1981-b):**  
Essential oil of the lemon tree leaves subtrop Kul't (5) 46-8 C.F. Chem. Abst. Vol. 96, No 74486.
- ; ----- **and ----- (1982-a):**  
Petitgrain oil from the Washington navel orange. Maslo-zhir. Prom-st. (6), 14. C.F. Chem. Abst. Vol 97, No 6087.
- ; ----- **and ----- (1982-b):**  
Essential oil of *Citrus wilsonii* leaves. Khimiya prirodnykh soedimenii No (6),785-786. C.P. (Hort. Abst. Vol 53, No 3718).
- **and ----- and G.M. Fishman (1985):**  
Production of absolute essential oil from flower of the mandarin (*Citrus Unhsiu*) Maslo-zhir. Prom-st. 18-19 C.F. Chem. Abst.Vol. 104, No 95242).

- Khaleel, A. E. ; S. H. El—Gayed; J. Nolte and R. Fobbe (2001):** comparative study of the essential oils of the leaves and fruits of *Fortunella margarita* (lour.) Swingle grown in Egypt. Az. J, pharm . Sci. 28 :258-265.
- Kijas, J. M. H.; Fowler, J. C. S. and Thomas, M. R. (1995):**An evaluation of sequence-tagged microsatellite site markers for genetic analysis within *Citrus* and related species. Genome. 38: 349-355.
- Lin, Z.; Y. Hua and G. Yuhong, (1986):** The chemical constituents of The essential oil from the flower, leaves and peels of citrus *Aurantium zhiwn*, xuebac, 28(6), 635-40. C.F.(Chem.Abst. Vol.106,1987, 182421 a).
- Lota, M. L.; D. De RoccaSerra; F. Tomi; C. Jacquemond and J. Casano (2001):**Chemical variability of peel and leaf essential oils of sour orange. Flavor and Fragrance Journal, Vol. 16 (2): 89-96.
- Lund, E. D.; P. E. Shaw and C. L. Kirkland (1982-a):** Components of key lime leaf oil. J. of Agricultural and Food Chemistry, 30 (1): 94-95.
- ; ----- and ----- (1982-b): Components of Meyer lemon leaf oil. J. of Agricultural and Food Chemistry, 30 (1): 95-97.
- Luro, F.; F. Laiget; J. M. Bove and P. Ollitrault (1995):** DNA amplified fingerprinting, a useful tool for determination of genetic origin and diversity analysis in *Citrus*. Hort. Science. 30 (5): 1063-1067.

- Martins, L.P. cavagnaro; R. Masuelli and J. Rodrigues (2003):**  
Evaluation of diversity among Argentine gropevine (*Vitis vinifera* L.) varieties using morphological data and ALFP markers . Elctronic Journal Botechnology, 6(3):241-253.
- Masada, Y. (1976):** Analysis of essential oil, by gas chromatography and mass sepectro scop.P. 142-158.John Wiley and Sons, Inc. New Yourk.
- Mehana, S.A. (1983):** Taxonomic and physiological studies on Egyptian sweet lime. Ph. D. Thesis Cairo, Univ.
- Melendreras, F. A.; J. Laencine; J. Flores and G. Guzman (1984):**  
Leaf essential oils from lemon tree cultivars (*Citrus limon* L. Burn. F.). Am. Fdafol. Agrobiol. 43, (7-8): 1161-1180.
- Mukhtar A.; M. L. Arpaia and R. W. Scora (2001):**Seasonal variation in lemon (*Citrus limon* L. Burm. t) leaf and rind oil composition. Journal of Essential Oil Research, Vol.13, (3):149-153.
- Naik, N.V. (1984):** Taxonomy of angiousperms. Marathmada university aurangabad Tata McGraw-Hill publishing company limited 1984, New Delhi.
- Nicolosi, E.; Z. N., Deng; A., Gentile; S., La-Malfa; G., Continella and E., Tribulato (2000):***Citrus* phylogeny and genetic origin of important species as investigated by molecular markers. Theoritical and Applied Genetics, 100:1155-1166.
- Osman, A. M; M. E. Younes and M. F. M. Ata,(1970):** Chemical examination of local plants . 1(constituents of the peel of Egyptian orange ( *citrus sinensis* ) J. chem. UAR, 13 (3), 337-45 En.



- Pasquale, M. R. Garcia, M. J. Asins, J. Forner, E. A. Carbonell (2006):** Genetic analysis of apomixes in *Citrus* and *Poncirus* by Molecular markers. Theor. Appl. Genet., 99:511-518.
- Pieringer, A. P.; G. J. Edwards and R.W., Wolford (1964):** The identification of citrus species and varieties by instrumental analysis of citrus leaf oils. Prop. Amer. Soc. Hort. Sci., 84:204-212.
- Rohlf, F. J. (2000):** NTSYS-pc: Numerical taxonomy and multivariate Analysis system Version 2.02 Exeter Software, Setauker, N. Y.
- Rodriguez, G.O.de; M. de; Godoy, N. G. de Colmenres and C.S. Ferrer (1998)** composition of Venezuela Lemon essential oil *Citrus limon*( L.) Burm. F. J. Agric. Food chem.. 27 (2):17-24.
- Rojas, E. and Scorza, J. V.(1991):** The use of essential lemon oil as a sand fly repellent. Transactions of the Royal Society of Tropical Medicine and Hygiene, No.6, P803, 1 ref.
- Saleh, M. M.; F. A. Hashem and K. W. Glombitzab (1998):** Study of *Citrus sinensis* and radical scavenger activity of the flavonoids isolated Food Chemistry, Vol. 63, No. 3, pp. 397-400.
- Salib, A. G.; M. A. Saleh and G. S. Abdel Malik, (1978):** Chemical and physical studies on peel essential oils of some Egyptian citrus fruits. Annals of Agricultural Science Moshtohor , 2, 65-72.

- Salimath, S. S.; A. C., Olivera, I. D. Godwin and J. L. Bennetzen (1995):** Assessment of genome origins and genetic diversity in the genus *Eleusine* with DNA markers. *Genome*. 38: 757-763.
- Santos, P. K.; Dornelles, A. L. D. and de Freitas, L. B. (2003):** Characterization of mandarin citrus germplasm from Southern Brazil by morphological and molecular analyses. *Pesq. agropec. bras.*, Brasília, v. 38 (7): 797-806.
- Sawamura, M.; S. Klwahra; K. Shicihri and T. Aoki. (1990):** Volatile constituents of several of pummelos and a comparison of the nootktone levels in pummelos and other citrus , fruits. *Agricultural and Biological chemistry* (1990), 54(3): 803-805(En-10 ref.)
- Scora. R and M. Ahmed (1994):** The leaf oils of *Severinia buxifolia* (Poir.) Tenore. *Journal of Essential Oil Research*. 6 (4): 363-367.
- **C. W. Cameron and A.B. Engand (1968):** Rind oil components of intergeneric citrus poncirus hybrids and their parents . *Proc.Amer. Soc. Hort. Sci.*,92:221-6.
- Scora, R.W. and S. Torrisi (1965):** Reation of taxonomic, climatic and tissue maturity factors of the essential oil constituents in leaves and fruits in the Aurantioideae, citrus research center and fruits in experiment station, univ. of california, November 2. 1965,paper No. 1684.
- Scora, R. W., S. Torrisi and C. W. Cameron and A. B. England (1968):** Rind oil components of intergeneric citrus poncirus hybrids and their parents. *Proc. Amer. Soc. Hart. Sci* , 92: 221-261.

- Sergio, L. and Gianni, (2005):** Molecular markers based analysis for crop germoplasm preservation. The role of Biotechnology, Villa Gualino, Turine, Italy, 5-7 March. [www.fao.org/biotech/does/ianteri.pdf](http://www.fao.org/biotech/does/ianteri.pdf).
- Shams El-Dein, M.M. (1980):** Storage of orange peel oil . Ph. D. Thesis (Agric. Chem.)Fac. Of Agric.Ain Shams Univ.
- Shaw, P. E. and C.W. Wilson (1976):** 111- comparison of extracted peel oil composition and Juice flavor for rough lemon , Persian lime and lemon –lime crose. J. of Agric. And food xhemistry 24 (3): 664-66.
- Shaw, P. E.; M. G. Moshonasa and K. D. Bowmanb (1999):** Volatile constituents in juice and oil of Australian wild lime (*Microcitrus inodora*).
- Smith, D. C.; S. Forland; E. Bachanos; M. Matejka and V. Barrett. (2000):** Qualitative Analysis of Citrus Fruit Extracts by GC/MS: An Undergraduate Experiment.
- Sun, H.; D. Lishong; W. Yu and M. Zhonghui (1984):** Chemical constituents of essential oil from *Citrus madica* leaves. Yunnam Zhiwu Yanjiu, 6 (4): 457-460. (C.F. Chem. Abst. Vol. 102,).
- Teranishi, R.; T. H. Schlitz; W. H. Mcfodden, R. E. Lundin and D. R. Blach, (1963):** Volatiles from oranges 1. J. Food. Sci. 28,541-8.
- Tsumura, Y.; K. Ohba and S. H. Strauss (1996):** Diversity and inheritance of inter-simple sequence repeat polymorphisms in douglas-fir (*Pseudotsuga menziesii*) and sugi (*Cryptomeria japonica*). Theor. Appl. Genet. 92: 40-45.

- Vannier, H. (1962):** Monoterpene hydrocarbon composition of citrus oils.  
J. Food. Sci. 27, 593-6.
- Vekiari, S. A.; E. E. Protopapadakis ; P. papadopoulou; I. D. Papanicolaou; C. Panou and M. Vamvakias (2002):**  
Composition and seasonal of leaves and peel of a Cretan lemon variety. Journal of Agricultural and Food Chemistry, Vol. 50 (1):147-153.
- Wolff, K.; I. Zietkiewicz and H. Holstra (1995):** Identification of chrysanthemum cultivars and stability of DNA fingerprint patterns. Theoretical and Applied Genetics, 91:439-447.
- Wu, K. S. and S. D. Tanksley (1993):** Abundance polymorphism and genetic mapping of microsatellites in rice. Mol. Gen. Genet. 241: 225-235.
- Youssef, A. A.; A. L. Eissa ;M.A. Habib and E. N. Abou-Zied (1994):**  
Biochemical studies on some Citrus species. Ph.D. thesis Faculty of Agriculture. Minufiya University.
- Zayas, J. P.; R. Baluja; R. Tapanes; W. Schmidt and A. Rosado (1980):**On the chemical composition of Cuban petitgrain oils. III. Analytical study of Shaddock petitgrain oils. Rev. Cenic, Cienc. Fis. 11, (1-2), 9-16 (C.F. Chem. Abst. Vol. 95).
- Zielinski, Q. B. (1955):** Modern systematic Pomology Wu. C Brown Co. Iwa.; 186-194.
- Zietkiewicz, I.; N. Rafalski and D. Labuda (1994):** Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. Genome. 20: 176-183.