

VI- REFERENCES

- Abbott, W.S. (1925):** A method of computing the effectiveness of an insecticides. J. Econ. Ent., 18: 265-269.
- Abdallah, M.D.; Kandil, M.A. and Farag, A.A. (1986):** Bioactivity of plant extracts against *Sitophilus granarius* (L.) and *Tribolium castaneum* (Hbst.). Bull. Entomol. Soc. Egypt. Econ. Sci., 15: 199-205.
- Abd El-Aziz, A.E. (2002):** Effectiveness of petroleum ether extract of some plants alone and under controlled atmospheres of carbon dioxide against the cowpea beetle *Callosobruchus maculatus* (F.). Annals of Agric. Sci., Moshtohor, 40 (2): 1309-1319.
- Abo El-Ghar, G.E.S. and El-Sheikh, A.E. (1987):** Effectiveness of some plant extracts as surface protectants of cowpea seeds against the pulse beetle *Callosobruchus chinensis*. Phytoparsitica, 15 (2): 109-113.
- Abo El-Ghar, M.R. and Badawy, A. (1961):** A study on the effect of malathion and katelsous on stored grain pests. Bull. Soc. Ent. Egypt, 45: 445-452.
- Afifi, F.A.; Hekal, A.M. and Salem, M. (1988):** Fenugreek seed extracts as protectants of wheat grains against certain stored product insects. Annals Agric. Sci., Cairo, 33 (2): 1331-1341.

- Aguiar, J.B.W.; Monteiro, M.D. and Calafiori, M.H. (1994):** Alternative control of *Sitophilus sp.* on stored rice (*Oryza sativa* L.). E cossistema, 19: 67-74.
- Aku, A.A.; Ogunwolu, E.O. and Attah, J.A. (1988):** *Annona senegalensis* L. (Annonaceae): performance as a botanical insecticide for controlling cowpea seed bruchid, *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) in Nigeria. Zeitschrift-fur-pflanzenkrankeiten-und-pflanzenschutz, 105 (5): 513-519.
- Arshad, J.H. and Salehuddin, A.M. (1988):** Persistence of ¹⁴C-malathion residues in stored milled rice. Dept. Biochem., Fac. Sci. 8 Environmental studies, Univ. Pretanian Malaysia, Pertanika, 1988, 11 (1): 73-78.
- Babu, A.; Raja, N.; Albert, S.; Ignacimuthu, S. and Dorn, S. (1999):** Comparative efficacy of some indigenous plant extracts against the pulse beetle *Callosobruchus maculatus* F. (Coleoptera: bruchidae). Biological Agric. and Hortic., 17 (2): 145-150.
- Bandyopadhyay, B.; Paul, S.K. and Ghosh, M.R. (1996):** Effect of sublethal dose of some insecticides on the biology of *Sitophilus oryzae* (L.) (Coleoptera: curculionidae). India, Bidhan, Chandra, Krishi, Viswavid, Yalaya, Mohanpur 741252, Enviroment and Ecology, 14 (2): 334-338.
- Barbara, N.A. and Linda, J.M. (1983):** Degradation of ¹⁴C-malathion in stored corn and wheat inoculated with *Spergillus flavus*. J. Econ. Ent., 76: 733-736.

- Bareth, S.S. and Gupta, H.C. (1989a):** Efficacy of six insecticides for the protection of stored wheat seeds against *Rhizopertha dominica* (Fab.). Seed Res., 17 (1): 43-46.
- Bareth, S.S. and Gupta, H.C. (1989b):** Efficacy of gunny bag and seed treatment on the natural infestation of *Rhizopertha dominica* on stored wheat. Seed res., 17 (2): 178-181.
- Behera, D.K.; baral, K. and Mitra, S.R. (2002):** Effect of plant extracts on rice weevil *Sitophilus oryzae* L. J. of intracademia, 6 (1): 59-64.
- Bell, C.H. (1984):** Effect of oxygen on the toxicity of carbon dioxide to stored insects. Developments in Agric. Engineering: controlled atmosphere storage of grains, edited by Ripp et al., B.E., 67-74.
- Bellal, M.H.; Abd El-Rahman, A.G.; Ibrahim, N.M. and Ali, E.A. (2001):** Observations on the repellent effect of some desert plant extracts on the red beetle *Tribolium confusum*. Integrtaed pest management proceedings of the 1st Congress, cairo Univ. Fac. of Agric. Dept. of Econ. Entomol. And Pesticides, 22-23 April 2001.
- Bhuiyah, M.I.M. (1993):** Evaluation of insecticides in sacks for the control of *Sitophilus oryzae* L. in stored wheat. Bangladesh J..of Zoology, 21 (2): 67-75.
- Bhuiyah, M.I.M.; Alam, S. and Karim, A.N.M.R. (1988):** Studies on the nature and extent of damage to stored rice caused by rice weevil *Sitophilus oryzae* (Linnaeus)

and angoumois grain moth *Sitotroga cerealella* (Olivier) and their control in Bangladesh. Bangladesh J. of Zoology, 16 (2): 99-109.

Bitran, E.A.; Campos, T.B.; Brisolla, A.D. and Chiba, S. (1986): Evaluation of the residual persistence of insecticides on the protection of barley grains. Biologico. 1986 Publ. 1988, 52: (4-6), 53-55.

Chander, H. and Ahmed, S.M. (1987): Laboratory evaluation of natural embelin as a grain protectant against some insect pests of wheat in storage. J. of Stored Products res., 23 (1): 41-46.

Chimbe, G.M. and Galley, D.J. (1996): Evaluation of material from plants of medicinal importance in Malawi as protectants of stored grains against insects. Crop-protection, 15 (3): 289-294.

Dakshinamurthy, A. and Goel, S.C. (1992): Insect management in grain and seed storage of wheat using non-hazardous materials. Bioecology and control of insect pests: proceeding of the national Symposium on growth, development & control Tech. of Insect Pests: 265-268.

Daoud, A.S. and Al-Muffti, A.S. (1987): Effect of certain pesticides on germination of wheat seeds as influenced by storage temperature and type of sacks. Arab. J. of Plant Protection, 5 (2): 63-66.

Darwish, A.A. (1997): Effect of neem and datura plant extracts under modified atmospheres on certain stored products

insects. Annals of Agric. Sci., Moshtohor, 35 (4): 2529-2544.

Deb-Kirtaniya, S.; Ghosh, M.R.; Mitra, S.A.; Adityachau, D.N. and Chatterjee, A. (1980): Extract of chilli as possible source of insecticides. Indian J. Agric. Sci., 50 (6): 507-510.

Deighton, J.C.; Donahaye, E. and Navarro, S. (1987): Actellic advantages for grains storage. Proceedings of the fourth International working conference on stored product Protection, Tel Aviv, Israel, 21-26 September 1986, 528-539.

El-Lakwah, F.A.M. and Abd El-Latif, A.M. (1998): Effect of surface treatment of grain bags with the botanical insecticide neemazal-T and certain plant extracts on weight loss of stored wheat and maize grains due to infestation with *Sitophilus oryzae* (L.), Egypt. J. Agric. Res., 76 (3): 1003-1016.

El-Lakwah, F.A.M.; Abd El-Latif, A.M. and El-Kashlan, H.I. (1999): Influence of diet treatment with black pepper (*Piper nigrum* L.) powder on the populations of *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst.). Annals of Agric. Sci., Moshtohor, 37 (1): 639-650.

El-Lakwah, F.A.M.; Abd El-Latif, A.M. and Halawa, Z.A. (1998): Effect of *Conyza dioscoridis* leaf extracts on three major insects of stored products in Egypt. Egypt. J. Agric. Res., 76 (3): 971-982.

- El-Lakwah, F.A.M.; Azab, M.M. and Abd El-Latif, A.M. (2001):** Efficiency of lupine seed extracts with and without controlled atmospheres against some stored product insects. *Annals of Agric. Sci., Moshtohor*, 39 (4): 2499-2512.
- El-Lakwah, F.A.M.; Darwish, A.A. and Halawa, Z.A. (1996a):** Toxic effect pf extracts and powders of some plants against the cowpea beetle (*Callosobruchus macalutus*). *Annals of Agric. Sci., Moshtohor*, 39 (4): 1849-1859.
- El-Lakwah, F.A.M.; Darwish, A.A. and Omnia, M. Khaled (1992a):** Effectiveness of dill seed powder (*Anethum graveolens*) on some stored product insects. *Annals of Agric. Sci. Moshtohor*, 30 (4): 2031-2037.
- El-Lakwah, F.A.M.; Darwish, A.A. and Omnia, M. Khaled (1992b):** Toxic effects of pulverized black pepper (*Piper nigrum* L.) on some stored product insects. *Annals of Agric. Sci. Moshtohor*, 30 (4): 2049-2057.
- El-Lakwah, F.A.M.; Darwish, A.A.; Khattab, M.M. and Abd El-Aziz, A.E. (1993a):** Effect of some insecticides on some stored product insects. *Annals of Agric. Sci. Moshtohor*, 31 (1): 529-547.
- El-Lakwah, F.A.M.; El-Kashlan, H.I. and Abd El-Aziz, E.A. (2000 a):** Effectiveness of Dill (*Anethum graveolens* L.) seed extract under modified atmospheres against some stored product insect pests. *Adv. Agric. Res.*, 5 (3): 1589-1604.

- El-Lakwah, F.A.M.; Halawa, Z.A. and Abd El-Latif, A.M.**
(1997): Effect of the extracts of *Withania somnifera* leaves and fruits on some stored product insects. Annals of Agric. Sci., Moshtohor, 35 (1): 537-552.
- El-Lakwah, F.A.M.; Hamed, M.S. and Abd El-Latif, A.M.**
(1996c): Effectiveness of *lantana Camara* and *Nerium oleander* extracts alone and mixtures with two insecticides against *Rhizopertha dominica*. Annals of Agric. Sci., Moshtohor, 34 (4): 1879-1905.
- El-Lakwah, F.A.M.; Khattab, M.M. and Darwish, A.A.**
(1989): Effect of fenugreek addition to the diet on the populations of *T. castaneum* and *S. oryzae*. Egypt J. Appl. Sci., 4 (3): 665-672.
- El-Lakwah, F.A.M.; Mahgoub, S.M. and Mohamed, S.M.**
(2000 b): Effect of maize husk ash and mustard seeds powder (*Brassica arvensis*) as grains protectants on some stored product insects. Annals of Agric. Sci., Moshtohor, 38 (1): 565-571.
- El-Lakwah, F.A.M.; Mohamed, R.A. and Darwish, A.A.**
(1995): Evaluation of the toxic effect of chinaberry (*Melia azedarach*) on *Sitophilus oryzae*. Annals of Agric. Sci., Moshtohor, 33 (1): 389-398.
- El-Lakwah, F.A.M.; Mohamed, R.A. and Omnia, M. Khaled**
(1994b): Toxic effect of Chinaberry tree (*melia azedrach*): meliaceae on *Rhizopertha dominica*. Annals of Agric. Sci., Moshtohor, 32 (4): 2149-2204.

- El-Lakwah, F.A.M.; Omnia, M. Khaled and Darwish, A.A. (1993b):** Laboratory studies on the toxic effect of some plant seed extracts on some stored product insects. *Annals of Agric. Sci. Moshtohor*, 31 (1): 593-603.
- El-Lakwah, F.A.M.; Omnia, M. Khaled and Darwish, A.A. (1994a):** Determination of organochlorine and organophosphorus pesticide residues in home-produced wheat during its transport from shounas to mills: *Annals of Agric. Sci. Moshtohor*, 32 (4): 2117-2123.
- El-Lakwah, F.A.M.; Omnia, M. Khaled and Halawa, Z.A. (1996b):** Effect of modified atmosphere of more than 99% nitrogen or enriched with carbon dioxide on mortality of some stored product insects, seed germination, chlorophyll and carotene contents of certain crop seedlings. *Annals of Agric., Sci., Moshtohr*, 34 (4): 1869-1877.
- El-Lakwah, F.A.M.; Omnia, M. Khaled and Mohamed, R.A. (1993c):** Evaluation of the toxic effect of neemazal-F (powder contains 33% azadirachtin) on adults of *Sitophilus oryzae* (L.), *Rhizopertha dominica* (F.) and *Tribolium castaneum* (Herbst). *Egypt. J. Appl. Sci.*, 8 (7): 43-59.
- El-Nahal, A.K.M. and El-Halafawy, M.A. (1974):** The Effect of sublethal treatments with pyrethrins and certain inert dusts on some biological aspect of *S. oryzae* L. and *S.*

granarius L. (coleoptera). Bull. Entomol. Soc. Egypt. Econ. Ser., 7: 253-260.

Figueiredo, F.J.C.; Frazaor, D.A.C.; Oliveira, R.P. and Correa, J.R.V. (1982): Storage of cowpea seed in the physiographic regions of the state of para, circular-Technica. Centro-de-pesquisa-Agropecuaria-do-Tropico-Umido No. 30, 48 pp.

Finney, D.J. (1971): Probit analysis 3rd Edt., Cambrudge Univ. Press: Cambridge, UK 318 pp.

Gadavari-Bio, S.; Krishnomurthy, K. and Mojumders, S.K. (1964): Malathion for stored product insects control. Int. Pest Control, 6: 9-10.

George, J.; Ravishankar, G.A.; Pereira, J. and Divakar, S. (1999): Bioinsecticide from swallowroot (*Decalepis hamiltonii*) wight & an protects food grains against insect infestation. Current Sci., 77 (4): 501-502.

George, T. and Dikshit, A.K. (1995): Persistence of deltamethrin on stored blackgram grains and its decontamination. Pesticide-Research-Journal. 1995, 7 (1): 76-80.

Giga, D.P.; Mutemerewa, S.; Moyo, G. and Neeley, D. (1991): Assessment and control of losses caused by insect pests in small farmers' stores in Zimbabwe. Crop protection, 10 (4): 287-292.

Gouhar, K.A.; Mansour, M.M. and Amir, M.M.I. (1975): Biological changes of *Callosobruchus maculatus* (F.)

achieved during selection with certain insecticides.
Zagazig J. of Agric. Res., 2: 231-239.

Grant, J.A.; Parker, B.L. and Damardjati, D.S. (1990): Controlled-release insecticides for stored grain pests control in Indonesia II Ware house trial. Asean Food J., 5 (2): 71-78.

Gujar, G.T. and Meherotra, K.N. (1983): Inhibition of growth and development of the tobacco caterpillar, *Spodoptera litura*, Fabr. due to azadirachtin and other neem products. Indian J. Ent., 45: 431-435.

Gupta, H.C.; Verma, J.R.; Bareth, S.S. and Mathur, B.N. (1988): Evaluation of some non-edible oils as grain protectant in wheat and their subsequent effect on germination. Ind. J. of Entomol., 50 (2): 147-150.

Hamel, D. (1987): Effectiveness of organophosphorus insecticides against stored product insects in wheat. Inst. Plant Protection, Zagreb, Yugoslavia. 637. Bet. Dagan Israel; Agricultural research Organization.

Harish-Chander; Kulkarni, S.G. and Berry, S.K. (1990): *Acorus calamus* rhizomes as a protectant of milled rice against *Sitophilus oryzae* and *Tribolium castaneum*. J. of Food Sci. and Tech. India, 27 (3): 171-174.

Hashem, M.Y. and Richmuth, Ch. (1994): Interactive action of high carbon dioxide or low-oxygen atmospheres and temperature on hatchability of eggs of three stored product moths. J. Plant Diseases and Protection, 101 (2): 178-182.

- Haubruge, E.; Schiffers, B.; Gabriel, E. and Verstraeten, C. (1988):** Study of the dose-efficacy relationship for six insecticides with regard to *Sitophilus granarius* L., *S. oryzae* L. and *S. zeamais* Mots. (Col., Curculionidae). Mededlingen-van-de-Faculteit-Landbouwwetenschappen-Rijksuniversiteit-Gent. 53 (26): 719-726. In International Symposium on Crop Protection.
- Ho, S.H.; Lee, L.S.; Tong, Y.; Ma, Y. and Sim, K.Y. (1995):** Effects of non-polar extracts of clove flower buds on some life stages of *Tribolium castaneum*. Singapore Int. Pest. Control, 37 (4): 112-113.
- Hung, C.C.; Hsieh, F.K. and Hwang, J.S. (1990):** Stored grain insects monitoring and their chemical control. Chinese. J. of Entomol., 10 (2): 169-179.
- Ivbijaro, M.F. (1984):** Toxic effects of groundnut oil on the rice weevil *S. oryzae* insect. Sci. Appl., 5 (4): 251-252.
- Jacob, S. and Sheila, M.K. (1993):** A note on the protection of stored rice from the lesser grain borer, *Rhizopertha dominica* Fabr. by indigenous plant products. Ind. J. of Entomol., 55 (3): 337-339.
- Jaipal, S.; Singh, Z. and Malik, O.P. (1984):** Insecticidal activity of various neem leaf extracts against *Rhizopertha dominica*, a stored grain pest. Neem Newsl., 1: 35-36.
- Javaid, I. And Poswal, M.A.T. (1995):** Evaluation of certain spices for the control of *Callosobruchus maculatus*

(Fabricius) (Coleoptera: Bruchidae) in cowpea seeds. African Entomol., 3 (1): 87-89.

Jembere, B.; Obeng-Ofori, D.; Hassanali, A. and Nyamasyo, G.N.N. (1995): Products derived from the leaves of *Ocimum kilimandscharicum* (Labiatae) as post-harvest grain protectants against the infestation of three major stored product insect pests. Bulletin of Entomol. Res., 85 (3): 361-767.

Jilani, G. and Amir, P. (1987): Economics of neem in reducing wheat storage losses. Policy Implications. Tech. Bull. No. 2. SEARCA, Phil., P15.

Jilani, G. and Malik, M.M. (1973): Studies on neem plant as repellent against stored grain insects. Pakistan J. Sci. Ind. Res., 16 (6): 251-254.

Kansouh, A.S.H.; Al-Rajhi, D.H.; Hashem, S.M. and Hamada, A.K. (1989): Residues of certain organophosphorus insecticides in the treated bags of wheat flour. J. of king Saud Univ., Agric. Sci., 2: 137-142.

Kelly, M.R. and Amos, K.M. (1993): The control of insects in export grains by admixture chemicals. HGCA-Project-Report. 69, 25 pp.

Kishino, H.; Goto, M.; Imamura, M. and Soma, Y. (1996): Responses of stored grain insects to carbon dioxide. 2 Effects of temperature and exposure period on the toxicity of carbon dioxide to *Sitophilus granarius*, *Lasioderma serricorne*, *Plodia interpunctella*, *Ephestia*

cautella and *Ephestia kuehniella*. Res. Bull. of the plant protection service. Japan, 32: 57-61.

Korunic, Z.; Hamel-Koren, D. and Koren, D. Hamel (1985):

The effect of deltamethrin and pririmiphos methyl on pests of stored products. Zastita Bilja, 36 (4): 417-423.

Koul, O.; Isman, M.B. and Ketkar, C.M. (1990): Properties and uses of neem, *Azadirachta indica*. Can. J. Bot., 68: 1-11.

Lalah, J.O. and Wandiga, S.O. (1996): The persistence and fate of malathion residues in stored beans (*Phaseolus vulgaris*) and maize (*Zea mays*), Pesticide Science. 1996, 46: 3, 215-220.

Lindgren, D.L. and Vincent, L.E. (1970): Effect of atmospheric gases alone or in combination on the mortality of granary and rice weevils. J. Econ. Ent., 63: 1926-1929.

Locatelli, D.P. and Daolio, E. (1991): Effectiveness of CO₂ in avacum autoclave at different temperatures on *Rhizopertha dominica* (F.) and *Sitophilus oryzae* (L.). Istituto di Entomol. Agraria, Univ. degli studi, via celoria 2, 20133 Milan, Italy. Tecnica-Molitaria, 42 (7): 605-610.

Maheswaran, P. and Ganesalingam, V.K. (1988): Effect of volatile substances from *Azadirachta indica* (neem) seeds on the reproductive biology of *Tribolium castaneum* (Coleoptera: Tenebrionidae). Univ. of Jaffna, Srilanka, Vingnanam J. of Sci., 1 (2): 20-26.

- Mahgoub, S.M. and Ahmed, S.M.S. (1996):** *Ricinus communis* seeds extract as protectants of wheat grains against the rice weevil *Sitophilus oryzae* L. Annals of Agric. Sci., Cairo, 41 (1): 483-491.
- Makanjuola, W.A. (1989):** Evaluation of extracts of neem (*Azadirachta indica*). J. of stored products Res., 25 (4): 231-237.
- Meguenni, H.; Bennaceur, M. and Zayed, S.M.A.D. (1996):** Persistence of malathion residues in stored wheat; Bulletin of the National research center Cairo. 21 (3): 239-244.
- Mishra, B.K.; Mishra, P.R. and mohapatra, H.K. (1992):** Studies on some plant product mixtures against *Sitophilus oryzae* (L.) infesting wheat seeds. Ind. J. of Plant Protection, 20 (2): 178-182.
- Mohamed, R.A. (1999):** Effectiveness of some plant extracts under modified atmospheres against the cowpea beetle, *Callosobruchus maculatus* (F.). Annals of Agric. Sci., moshtohor, 37 (1): 709-720.
- Mohapatra, S.; Sawarkar, S.k. and Patnaik, H.P. (1996):** Comparative bioassay of different solvent extracts of neem seed kernels against *Sitophilus oryzae*, L. in rice. Current Agric. Res., 1 (2): 89-93.
- Morner, J. Hyvonen, A. and Simonsson, M. (1987):** Experiments with carbon dioxide against stored products pests. Vaxtskyddsnotiser, 51 (3): 87-92.

- Mostafa, T.S. (1993):** Effects of certain plant powders on mortality, disrupting growth and metamorphosis of *Trogoderma granarium* Everts fourth instar larvae. Bull. Entomol. Soci. of Egypt, Econom. Seri., 20: 67-75.
- Muda, A.R. (1985):** Pest problems and the use of pesticides in grains storage in Malaysia. ACIAR-proceedings-series, Australian center for International Agric. Res., 14: 11-16 In Pesticides and humid tropical grains storage systems. Proceedings of an international seminar, Manila, Philippines, 27-30 May.
- Navarro, S. and Jay, E.G. (1987):** Application of modified atmospheres for controlling stored grains insects. Monograph British Crop protection Council, 37: 229-236.
- Navarro, S.; Dias, R. and Donahaye, E. (1985):** Induced tolerance of *Sitophilus oryzae* adults to carbon dioxide. J. stored prod. Res., 21: 207-213.
- Niber, B.T.; Helenius, J. and varies, A.L. (1992):** Toxicity of plant extracts to three storage beetles (Coleoptera). J. of Applied Entomol., 133 (2): 202-208.
- Noack, S. and Reichmuth, Ch. (1978):** Ein rechnerisches verfahren zur bestimmung von beliebigen Dosiswerten eines wirkst offesaus empirischen Dosis wirkungs Daten: Mitt, Biol. Bundesanstalt fuer land u. Forts wirtschaft, Berlin Dahlem, Heft, 185: 1-49.

- Ofuya, T.I. and Reichmuth, Ch. (1993):** Mortality of the bean bruchid *Acanthoselides obtecus* (say) in some high carbon dioxide atmospheres. J. Plant Diseases and protection, 100 (2): 165-569.
- Ogunwolu, O. and Idowu, O. (1994):** Potential of powdered *Zanthoxylum zanthoxyloides* (Rutaceae) root bark and *Azadirachta indica* (meliaceae) seed for control of the cowpea seed bruchid, *Callosobruchus maculatus* (Bruchidae) in Nigeria. J. of African Zoology, 108 (6): 521-528.
- Omar, E.E.; Hashem, M.Y. and Ismail, I.I. (1995):** Effect of carbon dioxide in air on different stages of the rice weevil *Sitophilus oryzae* (L.) (Curculionidae: Coleoptera). Egypt J. of Agric. Res., 73 (1): 119-126.
- Pandey, G.P.; Shanker, S. and Srivastava, J.L. (1983):** Relative toxicity of some newer organophosphorus insecticides against stored grain insect pests. Bull. of grain Techno., 21 (1): 19-23.
- Paneru, R.B.; Duwadi, V.R. and Bhattaral, M.R. (1993):** second year of testing locally available plant materials against weevil *Sitophilus oryzae* Linn. in stored wheat. Pakhribas Agricultural Center No. 44, 7, (c.f. Rev. of Agric. Entomol., 82 (6): 652, 5718).
- Papadopoulou-Mourkidou, E. and Tomazou, T. (1991):** Persistence and activity of permethrin in stored wheat and its residues in wheat milling fractions. J. of stored products Res., 1991, 27 (4): 249-254.

- Shawir, M.S. (1995):** Measurement of joint action of certain chemicals and amorphous silica mixtures as wheat protectant against rice weevil *Sitophilus oryzae* (L.). Alexandria J. of Agric. Res., 40 (1): 243-253.
- Shemais, Sawsan, A. and Al-Moajel, Nadra, H. (2000):** Efficiency and persistence of extracted capparid, *Capparis spinosa* seeds against the rice weevil, *Sitophilus oryzae* L. (Curculionidae: Coleoptera). Egypt J. Appl. Sci., 15 (2): 267-274.
- Shivanna, S.; Lingappa, S. and Patil, B.V. (1994):** Effectiveness of selected plant materials as protectants against pulse beetle, *Callosobruchus chinensis* (Linn.) during storage of redgram. Karnataka J. of Agric. Sci., 7 (3): 285-290.
- Singh, D. and Yadav, T.D. (1996):** Persistence of deltamethrin, chlorpyrifos methyl, etrimfas, malathion and flunvalinate on five surfaces. Ind. J. of Entomol., 57 (3): 206-212.
- Singh, H.; Mrig, K.K. and Mahla, J.C. (1996):** Effect of different plant products on the fecundity and emergence of lesser grain borer, *Rhizopertha dominica* (F.) in wheat grains. Annals of Bio. Ludhiana, 12 (1): 96-98.
- Singh, S.R.; Luse, R.A.; Leuschner, K. and Nangtu, D. (1978):** Groundnut oil treatment for control of *Callosobruchus maculatus* F. J. stored product Res., 14: 77-80.

- Singh, Y.P.; Gangwar, S.K. and Thakur, N.S.A. (1992):** Persistence of endosulfan in/on maize at medium high altitude hills of Meghalaya. Indian J. of Plant Protection. 20 (2): 195-197.
- Snedecor, G.W. and Cochran, W.G. (1967):** Statistical Methods, Sixth edition, Iowa State College press, Ames. Iowa, USA.
- Spratt, E.C. (1984):** The effects of atmospheres containing 5-20% oxygen with and without 10% carbon dioxide on the productivity of *Tribolium castaneum* (Herbst.) and *T. confusum* (Duval) (Coleoptera: Tenebrionidae). U.K. J. of stored products Res., 20 (4): 199-203.
- Srivastava, J.L.; Requpathy, A. and Jayaraj, S. (1985):** Use of controlled atmosphere for the control of stored product insects. Indian grain storage Institute, Hapur 205101, Uttar Pradesh, India Behavioural and Physiological approaches in pest management. 1985: 205-207.
- Strong, R.G.; Partoa, G.J. and Archel, I.L. (1969):** Comparative susceptibility of confused and red flour beetles from various areas of California to malathion. J. Econ., 62: 470-474.
- Su, H.C.F. (1984):** Evaluation of black pepper as a protectant of wheat in storage against rice weevils. J. Entomol. Sci., 19: 45-48.

- Su, H.C.F. (1985):** Laboratory evaluation of activity of *Cinnamomum cassia* to four species of stored product insects. J. Sci., 20: 247-253.
- Talukder, F.A. and Howse, P.E. (1994):** Laboratory evaluation of toxic and repellent properties of the pithraj tree, *Aphanamixis polystachya* Wall & Parker, against *Sitophilus oryzae* (L.). Int. J. of Pest Management, 40 (3): 274-279.
- Talukder, F.A. and Howse, P.E. (1995):** Evaluation of *Aphanamixis polystachya* as a source of repellents, antifeedants, toxicants and protectants in storage against *Tribolium castaneum* (Herbst). J. Stored Prod. Res., 31 (1): 55-61.
- Thaung, M. and Collins, P.J. (1986):** Joint effects of temperature and insecticides on mortality and fecundity of *Sitophilus oryzae* (Coleoptera: Curculionidae) in wheat and maize. J. of Econ. Entomol., 79 (4): 909-914.
- Tiwari, S.N. (1994):** Efficacy of some plant products as grain protectants against *Rhizopertha dominica* (F.) (Coleoptera: Bostrychidae). Int. J. of Pest management, 40 (1): 94-97.
- Toitil, T.P.S. and Pandey, G.P. (1979):** Insecticidal properties of rhizomes of sweet flage, *Acorus calamus* against rice weevil *Sitophilus oryzae* L. Indian J. Entomol., 41 (1): 91-94.

- Tune, I. (1983):** Mortality of *Tribolium confusum* (Col., Tenebrionidae) adults in various atmospheric gases compositions. Zeitschrift fur Angewandte Entomologie, 95 (3): 263-264.
- White, N.D.G. (1953):** Weight loss in stored wheat caused by insect feeding. J. Econ. Entomol., 46: 609-610.
- White, N.D.G. (1984):** Residual activity of organophosphorus and pyrethroid insecticides applied to wheat stored under simulated western Canadian conditions. Canadian Entomol., 116 (10): 1403-1410.
- White, N.D.G. (1988):** Residual activity of insecticides on freshly harvested and previously stored wheat, and on various carriers exposed to concrete surfaces. Proceeding of the Entomol., Soci., of Ontario, 119: 35-41.
- White, N.D.G. and Sinha, R.N. (1990):** Effects of sub-lethal exposures to malathion or chlorpyrifos-methyl on offspring production by some stored-product beetles. Agric. Canada Res. Station, 195 Dafoe Road, Winnipegman. R3T2Mg, Canadian Entomologist, 122 (11-12): 1187-1191.
- White, N.D.G.; Jayas, D.S. and Sinha, R.N. (1994):** Impact of pirimiphos-methyl and cold temperatures on arthropod populations in stored wheat. Phytoprotection Agric., and Agri-Food Canada, Res., Centre, 75 (2): 79-90.

- Williams, P.; Amos, T.G. and Gueschin, P.B. (1978):** Laboratory evaluation of malathion, chloropyrifos and chloropyrifos methyl for use against beetles infesting stored wheat. *J. Stored Prod. Res.*, 4: 163-168.
- Xie, Y.S.; Fields, P.G.; Isman, M.B.; Chen, W.K. and Zhang, X. (1998):** Insecticidal activity of *Melia toosendan* extracts and toosendanin against three stored product insects. *J. of stored products Res.*, 31 (3): 259-265.
- Yadav, T.D. (1984):** Efficacy of neem (*Azadirachta indica A. juss*) kernel powder as seed treatment against pulse beetles. *Neem Newsl.*, 1: 13-15.
- Yadav, T.D. and Singh, S. (1994):** Persistence toxicity and efficacy of four insecticides as jute fabric treatment to protect cereal and legume seeds. *Ind. J. of Entomol.*, 56 (2): 146-155.
- Yung, T. and Burkholder, N. (1981):** Protection of stored wheat from the granary weevil by vegetable oils. *J. Econ. Entomol.*, 74: 502-505.
- Zabel, A.; Kostic, M. and Oreskovic, M. (1987):** Results of evaluation of the persistence of action of the preparation of Reldan 2E and Actellic 50 EC in the control of the rice weevil (*Sitophilus oryzae* L.). *Zastita Bilja*, 38: 263-271.
- Zang, G.L.; Xu, B.Z.; Li, Y.H.; xue, L.S.; Ye, Z.X. and Shen, Z.c. (1982):** effect of pesticides on stored seeds quality. *Food Res. Inst. Zhejiang Province, Hangzhou*,

Zhejiang, China. Grain storage Liangshi chucang No. 3, 55: 11-16.

Zdarkova, E. and Horak, E. (1984): "Actellic" an insecticide for control of pests in food and from product stores. Sbornik-UVTIZ, potravnarske-ved, 2 (1): 41-48.