




REFERENCES

REFERENCES

- Abbott, W. S. (1925):** A method of computing the effectiveness of insecticides. *J. Econ. Entomol.*, 18: 265-207.
- Abd-Rabou, S.; Hussein, N.; Sewify, G. H. and Elnagar, S. (2000):** Seasonal abundance of the whitefly *Trialeurodes ricini* (Misra) (Homoptera: Aleyrodidae) on some weeds and on castor plants in Qalyubia, Egypt. *Bulletin of Faculty of Agriculture, University of Cairo*, 51(4): 501-510.
- Ammer, I. M. A.; Darwish, E. T. E.; Farag, A. I. and Eisa, A. A. (1986):** Detrimental effects of five moult-inhibiting insect growth regulators on the development and reproduction of the cabbage aphid, *Brevicoryne brassica* (l.). *J. of Appl. Entomology*, 102(4): 417-422.
- Assem, M. A. (1966):** Studies on vegetable leafminers. *Ph.D. Thesis, Fac. of Agric., Cairo Univ.*
- M.A. and El-Atrouzy, N.A. (1972):** Studies on the population density of mites associated with truck crops. *Agric. Res. Rev., Cairo*, 50(1):71 – 88.
- Attia, M. B. (1989):** Studies on the ecology and control of the broad bean leafminer, *Liriomyza trifolii* (Burgess): (Diptera: Agromyzidae) infesting bean fields in Egypt. *Third National Conf. of Pests & Dis. of Veg. & Fruits, Ismailia, Vol. 1*: 278-288.
- Azmi, P. R. and Rathi, Y. P. S. (1992):** Efficacy of insecticides against whitefly-a vector of French bean crinkle stunt virus. *Plant Disease Res.*, 7(2): 257-258.
- Barakat, A. A.; Shereef, G. M.; Abdallah, S. A. and Amer, S.A. (1984):** Joint action of some pesticides and plant extracts

- against *Tetranychus urticae* Koch. *Bulletin of the Entomological Society of Egypt, Economic Series*. 1984-1985, 14: 243-249.
- Bari, M. N. and Sardar, M. A. (1998):** Control strategy of bean aphid with predator, *Menochilus sexmaculatus* (F.) and insecticides. *Bangladesh Journal of Entomology*, 8(1,2): 21-29.
- Becquer, A.; Ferrandiz, R.; Hernandez and A.; Puga, R. (1981):** Population dynamics of aphid populations in fields of beans (*Phaseolus vulgaris* L.). *Ciencias de la Agricultura, Cuba.*, No. 8 : 23-28.
- Beitia, F.; Castaner, M. and Garrido, A. (1992) :** Incidencia de diversos plaguicidas aplicados sobre elestado larvario de *Diglyphus isaea* (Walker) (Hymn., Eulophidae). *Boletin de Sanidad Vegetal, Plagas* 18 (2): 303-308.
- Beitia, F.; Tormos, J. and Garrido, A. (1994):** Some aspects on the superparasitism of *Diglyphus isaea* (Walker, 1838). *Boletin de Sanidad Vegetal Plagas*, 20(3): 793-796.
- Bene, G. Del (1990):** Use of *Diglyphus isaea* (Walk.) (Hymn., Eulophidae) for the control of *Liriomyza trifolii* (Burgess), *Chromatomyia horticola* (Goureau) and *Chromatomyia syngenesiae* Hardy (Diptera: Agromyzidae) in greenhouses of chrysanthemums and gerbera. *Redia* 73(1): 63-78.
- Bene, G. Del.; Landi, S. and Del Bene, G. (1991):** Biological pest control in glasshouse ornamental crops in Tuscany. *Bulletin SROP*, 14(5): 13-21.
- Bishara, S. I.; Fam, E. Z.; Attia, A. A. and El-Hariry, M. A. (1984):** Yield losses of faba bean due to aphid attack. *FABIS-Newsletter*,

-Faba-Bean-Information-Service,-ICARDA. 1984, No. 10, 16-18.

- Bock, K. R.; Guthrie, E. j. and Woods, R. D. (1974):** Purification of maize streak virus and its relationship to viruses associated with streak diseases of sugarcane and *Panicum maximum*. *Annals of Applied Biology*, 77:289-296.
- Bright, A. A. and Regupathy, A. (2000):** Bioefficiency of Nimbecidine and TNAU neem on cowpea aphid *Aphis craccivora* Koch. applied to field bean. *Madras Agricultural Journal, Publ.*, 87(1-3): 116-122.
- Cabitza, F.; Cubeddu, M. and Ballore, S. (1993):** Two years of observation the application of biological control techniques against tomato pests on spring crops in greenhouses. *Informatore Agrario* 49(18): 103-106.
- Carballo, M.; Leon, G. R. and Ramirez, A. (1990):** Biological control of *Liriomyza* sp. (Diptera: Agromyzidae) in horticultural crops of Cost Rica. *Manejo Integrado de Plagas*, 16:4-11.
- Chalfant, R. B. and Young, J. R. (1984):** Management of insect pests of broccoli, cowpeas, spinach, tomatoes, and peanuts with chemigation by insecticides in oils, and reduction of watermelon virus 2 by chemigated oil. *J. Econ. Entomol.*, 77(5): 1323-1326.
- Chang, D. C. and Chen, C. C. (1993):** Population fluctuation of major insect pests on kidney bean and the proper time of control. *Bulletin of Taichung District Agricultural Improvement Station*, 38: 11-22.

- Chaudhuri, N. and Senapati, S. K. (2001):** Evaluation of pesticides from different origin-synthetic and biological, against pest complex of tomato under terai region of West Bengal. *Haryana Journal of Horticultural Sciences*. 2001, 30: 3-4, 274-277.
- Chauhan, R.; Sucheta, K.; Ombir-Singh; Khokhar, K. S. and Singh, O. (1988):** Bioefficacy and persistence of some insecticides against bean aphid, *Aphis craccivora* Koch. infesting chickpea. *Haryana J. of Horticultural Sci.*, 17(1-2): 102-105.
- Colombo, M. and Fasce, D. (1994):** Integrated control of spring crops of tomato in protected cultivation. *Informatore Agrario* 50(29): 37-41.
- Cruz-Amr, de; Cardona, C. J. and Cruz, J. L. de. (1989):** Life cycle, habits and natural enemies of the bean leaf miner *Liriomyza sativae* (Blanchard) (Diptera: Agromyzidae). *Acta Agronomica, Universidad Nacional de Colombia*, 39(3,4): 133-141.
- Dhaliwal, G. S.; Gill, R. S.; Dilawari, V. K.; Dhaliwal, G. S. (ed.); Randhawa, N. S. (ed.) and Dhawan, A. K. (1998):** Management of insect pest complex of cabbage with neem based insecticides. *Ecological Agriculture and Sustainable Development*, 2: 306-314.
- Dharmendar, S.; Jaglan, R. S.; Ramesh, C.; Singh, D. and Chauhan, R. (2003):** Field studies on the efficacy of insecticides against brinjal whitefly (*Bemisia tabaci* Genn.). *Annals of Biology*. 2003, 19: 1, 109-112.

- Dimetry, N. Z. (1971):** Biological studies on a leaf mining Diptera, *Liriomyza trifolii* (Burgess) attacking beans. *Bull. Soc. Entomol. Egypt* 55: 55-69.
- Dimetry, N. Z. and El-Hawary, F. M. A. (1995):** Neem Azal-F as an inhibitor of growth and reproduction in the cowpea aphid *Aphis craccivora* Koch.. *J. of Appl. Entomol.* 119(1): 67-71.
- Dimetry, N. Z. and El-Hawary, F. M. A. (1997):** Synergistic effect of some additives on the biological activity and toxicity of neem – based formulations against the cowpea aphid *Aphis craccivora* Koch. *Insect Science and its Application*, 17(3-4): 395-399.
- Dimetry, N. Z.; Barakat, A. A.; Abdalla, E. F.; El-Metwally, H. E. and El-Salam, A. M. E. A. (1995):** Evaluation of two neem seed kernel extracts against *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae). *Anzeiger Fur Schadlingskunde, Pflanzenschutz, Umweltschutz*, 68(2): 39-41.
- Doss, A. S.; Adam, K. M.; Herakly, F. A. and El-Hamaky, M. A. (1992):** Population densities of the broad bean leafminer, *Liriomyza trifolii* (Burgess) and the cotton whitefly, *Bemisia tabaci* (Genn.) on protected cultivation, *Minia. J. Agric. Res. & Dev.*, 14(3): 787-797.
- El-Defrawi, G. M.; Emam, A. K.; Marzouk, L. A. and Rizkalla, L. (2000):** Population dynamics and seasonal distribution of *Aphis craccivora* Koch. and associated natural enemies in relation to virus disease incidence in faba bean fields. *Egyptian J. of Agric. Res.*, 78(2): 627-641.

- El-Ghar, G. E. S. A. and Esmail, S. E. (1991):** Effects of synthetic pyrethroid insecticides on aphid population *Aphis craccivora* Koch. and yield of faba bean *Vicia faba* L. *Archiv Fur Phytopathologic und Pflanzenschutz*, 27(4): 279-288.
- El-Hemaesy, A. H. Hammad; Zeid, S. M. and Tantawy, M. L. (1974):** Control of the leafminer *Liriomyza trifolii* (Burgess) on broad bean. *Bull. Entomol. Soc. Egypt, Econ., Ser.*, 8: 197-205.
- El-Khayat, E. F.; El-Sayed, A. M.; Shalaby, F. F. and Hady, S. A. (1994):** Infestation rates with *Bemisia tabaci* (Genn.) to different summer and winter vegetable crop plants. *Ann. of Agric. Sci., Moshtohor*, 32(1): 577-594.
- El-Lakwah, F. A. and Mohamed, R. A. (1999):** Efficiency of datura fruit extracts and their mixtures with Malathion against the cowpea beetle, *Callosobruchus maculatus* (F). *Annals of Agric. Sci., Moshtohor*, 37(3):1967-1978.
- El-Lakwah, F. A. M. (1997):** Toxicity of the botanical insecticide, Neemazal-W (powder contains 10% Azadirachtin) alone and in mixtures with two insecticides against the lesser grain borer, *Rhizopertha dominica* (F.). *Annals of Agric. Sci., Moshtohor*, 35(4): 2583-2594.
- El-Lakwah, F. A.; Abdel-Latif, A. M. and Ferial M. El-Sayed (1999):** Toxicity and joint action of the botanical insecticide, Neemazal-W with Malathion to the cowpea weevil, *Callosobruchus maculatus* (F). *Egypt. J. Agric. Res.*, 77(3): 1023-1034.
- El-Lakwah, F. A.; Darwish, A. A.; Horeya, A. Abdel-Wahab and Shaalan, H. S. (1998):** Efficacy of two plant extracts,

Neemazal-T and their mixtures with Chlorpyrifos-methyl against some insect pests of cabbage plants. *Annals of Agric. Sci., Moshtohor*, 36(2): 1171-1209.

El-Lakwah, F. A.; Hamed, M. S. and Abdel-Latif, A. M. (1996): Effectiveness of *Lantana camara* and *Nerium oleander* extracts alone and in mixtures with two insecticides against *Rhizopertha dominica* (F.). *Annals of Agric. Sci., Moshtohor*, 34(4): 1879-1905.

El-Lakwah, F. A.; Omnia, M. Khaled; Abdel-Gawaad, A. A. and Wesam, M. K. Ebrahim (1997): Effectiveness of two plant extracts and their mixtures with insecticides against the red flour beetle, *Tribolium castaneum* (Herbst.). *Annals of Agric. Sci., Moshtohor*, 35(3): 1799-1820.

El-Nahal, A. K. M. and Assem, M. A. (1970): The chemical control of broad bean leaf miner, *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae). *Bull. Entomol. Soc. Egypt, Econ., Ser.*, 4: 265-271.

El-Sayed, A. M. and El-Gbar, G. E. S. A. (1993): Effect of selected insecticides on population, adult longevity and reproduction of the whitefly, *Bemisia tabaci* (Genn.) (Homoptera: Aleyrodidae). *Bull. Entomol. Soc. Egypt, Econ., Ser.*, 20: 161-171.

El-Sayed, A. M.; Shalaby, F. F. and Abdel-Gawad, A. A. (1991): Ecological studies on *Bemisia tabaci* (Genn.) (Homoptera- Homoptera: Aleyrodidae). Infesting different host plants. 1- Fluctuation and population density of *Bemisia tabaci* on different host plants. *Egypt J. Agric. Res.*, 69(1): 193-207.

- El-Serwy, S. A. (1993):** Preliminary field observations on the infestation, mine density, larval and pupal parasitism of *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae). *Bull. Entomol. Soc. Egypt*, 71: 75-81.
- Esbjerg, P. (1977):** The tomato mining fly *Liriomyza bryoniae*. *Manedsoversigt over Plantesygdomme*, 503: 122-123. (c.f. *R.A.E. (A)*, 66: 4, Abs. 2175 p. 276, 1978).
- Fagoonee, I. and Toory, V. (1983):** Preliminary investigations of host selections mechanisms by leaf miner *Liriomyza trifolii* (Burgess). *Insect Science and its Application*, 4(4): 337-341.
- Ferrer, X.; Casadevall, M. and Sorribas, R. (1987):** Trials to quantify the influence of pesticide products used to control *Liriomyza trifolii* (Burgess), on its parasites. *Fulls d'Informacio Tecnica*, No. 132: 4 pp.
- Finidori, Logli V.; A.G. Bagneres and J.L.Clement (1996):** Role of plant volatiles in the search for a host by the parasitoid *Diglyphus isaea* (Hymenoptera: Eulophidae). *J. Chemical Ecology* 22 (3): 541-558.
- Fowler, H. D. (1956):** Some physiological effects of attack by whitefly *Bemisia tabaci* (Genn.) and spraying parathion on cotton in the Sudan Gezira Empire Cotton Growing. *Review*, 33: 288-299.
- Freitas, J. C.; Bueno, M. A.; Calafiori, M. H. and Salvo, S. (1988):** Study on the control of the tropical mite, *Polyphagotarsonemus latus* (Banks) and the two spotted spider mite, *Tetranychus urticae* Koch on beans (*Phaseolus vulgaris* L.). *Ecossistema*, No. 13, 131-136.

- Gaffar, S. A.; Rather, A. J. and Mattoo, M. S. (1990):** Evaluation of various insecticides for the management of cowpea aphid, *Aphis craccivora* Koch. infesting moong. *J. of Aphidology*, 4(1-2): 36-38.
- Gameel, O. I. (1973):** Field evaluation of insecticides for jassid, *Empoasca lybica* De Berg and whitefly, *Bemisia tabaci* (Genn.) control on cotton. *Bull. Entomol. Soc. Egypt, Econ. Ser.*, 7: 113-122.
- Ghallab, M.M.A. (2001):** Ecological, Biological and Morphological studies on some predaceous mites (Suborder: Actinedida). *Ph. D. Thesis, Fac. of Science, Cairo Univ.*, 244 PP.
- Giray, H. (1970):** Investigations on the morphological characters brief biology and type of damage of *Liriomyza cicerina* Rond. (Diptera: Agromyzidae). *Ege Universitesi Ziraat Fakultesi Yayinlari No. 171*, 34 pp., Turkey. (c. f. R.A.E. (A), 60: 3, Abs. 743 p. 156, 1972).
- Grayer, R. J.; Kimmins, F. M.; Padgham, D. E.; Harborne, J. B. and Ranga-Rao, D. V. (1992):** Condensed tannin levels and resistance of groundnuts (*Arachis hypogaea*) against *Aphis craccivora* Koch.. *Phytochemistry*, 31(11): 3795-3800.
- Hafez, M. A. H.; El-Ziady, S. and Dimetry, N. Z. (1970):** Leaf mining Diptera of vegetables and crops in Egypt. *Bull. Entomol. Soc. Egypt*, 59: 249-259.
- Hammad, E. M. A. F.; Nemer, N. M.; Hawi, Z. K. and Hanna, L. T. (2000):** Responses of the sweet potato whitefly, *Bemisia tabaci* (Genn.), to the chinaberry tree (*Melia*

azedarach L.) and its extracts. *Ann of Appl. Biology*, 137(2): 79-88.

- Hammad, K. A. A. (2000):** The broad bean leafminer, *Liriomyza trifolii* (Burgess) and its parasitoids on different broad bean cultivars in newly reclaimed sandy land at El-Khattara district, Sharkia Governorate, Egypt. *Zagazig J. Agric. Res.*, 27(4): 1137-1159.
- Hammad, S. M. (1955):** On some dipterous leafminers from Egypt. *Bull. Entomol. Soc. Egypt*, 39: 391-394.
- Hannou, M. A. and Hegazi, E. M. (1996):** Effects of faba bean cultivars and potassium fertilization on population of *Liriomyza* sp. (Diptera: Agromyzidae). *J. Agric. Sci. Mansoura Univ.*, 21(12): 4565-4574.
- Hannou, Magda A. (1992):** Studies on some vegetable pests and their natural enemies. *Ph.D. Thesis, Fac. Agric., Alex. Univ.*
- Hassan, A.S.; El-Nahal, A.K.M. and El-Badry, E.A. (1959):** Infestation of cotton with spider mites. *Bull. Soc. Entomol., Egypte*, 40(1): 357 – 364.
- Heinz, K. M. and Parrella, M. P. (1990a):** Biological control of insect pests on greenhouse marigolds. *Environmental Entomol.*, 19(4): 825-835.
- Heinz, K. M. and Parrella, M. P. (1990b):** Holarctic distribution of the leafminer parasitoid *Diglyphus begini* (Hymenoptera: Eulophidae) and notes on its life history attacking *Liriomyza trifolii* (Diptera: Agromyzidae) in chrysanthemum. *Ann. Entomol., Soc. Amer.*, 83(5): 916-924.

- Heinz, K. M.; Newman, J. P. and Parrella, M. P. (1988):** Biological control of leafminers on greenhouse marigolds. *California Agric.*, 42(2): 10-12.
- Helal, H. A.; El-Khouly, A. S.; Khalafalla, E. M. E.; Metwally, M. M. and El-Mezaien, A. B. (1997):** Preliminary studies on the population of legume aphid, *Aphis craccivora* Koch. and jassid, *Empoasca* on different faba bean varieties and breeding lines under the field conditions of Kafr El-Sheikh Governorate, Egypt. *Egyptian J. of Agric. Res.*, 75(4): 951-959.
- Helaly, M.M.; Ibrahim, A.E. and Saleh, M.R.A. (1982):** Fluctuations of population densities of *Empoasca* sp., *Aphis craccivora* Koch, and *Tetranychus arabicus* Attiah attacking cowpea plants at Zagazig, Egypt. *Bulletin de la Societe Entomologique d'Egypte*. No. 64: 35-43.
- Hesselein, C.; Robb, K.; Newman, J.; Evans, R. and Parrella, M. P. (1993):** Demonstration/integrated pest management program for potted chrysanthemum in California. *Bulletin OILB/SROP*, 16(2): 71-76.
- Heyer, W.; Lok, M. L. C. and Cruz, B. (1989):** The population dynamics of the cotton whitefly *Bemisia tabaci* (Genn.), in bean fields in the Republic of Cuba. *Archiv Fur Phytopathologie und Pflanzenschutz*, 25(5): 473-479.
- Hightower, B.G. and Martin, D.F. (1956):** Ecological studies of spider mites attacking cotton in central Texas. *J. Econ. Entomol.*, 49: 424.
- Hilije, L.; Cubillo, D. and Segura, L. (1993):** Ecological observations on the whitefly *Bemisia tabaci* (Genn.), in Costa Rica. *Manejo Integrado de Plagas*, 30: 24-30.

- Hincapie, C. M. C.; Saavedra, H. M. E. and Trochez, P. A. L. (1993):** Life cycle, behavior and natural enemies of *Liriomyza huidobrensis* (Blanchard) on bulb onion (*Allium cepa* L.). *Revista Colombiana de Entomologia*, 19(2): 51-57.
- Hinderson, C. F. and Tilton, E. W. (1955):** Test with acaricides against the brown white mite. *J. Econ. Entomol.*, 48: 157-161.
- Hollingsworth, R. G.; Tabashink, B. E.; Johnson, M. W.; Messing, R. H. and Ullman, D. E. (1997):** Relationship between susceptibility to insecticides and fecundity across populations of cotton aphid (Homoptera: Aphididae). *J. of Economic Entomology*, 90(1): 55-58.
- Hurej, M. (1991):** Susceptibility of aphids (Homoptera: Aphididae) to extreme temperatures. *Wiadomosci Entomologiczne*, 10(1): 43-50.
- Ibrahim, Y. B. and Yeow, W. K. (1989):** The effects of different concentrations of carbaryl on the life-table parameters of *Aphis craccivora* Koch.. *Pertanika*, 12(1): 43-46.
- Ichikawa, K.; Ohno, T. and Nakagome, T. (1996):** Control of the legume leafminer *Liriomyza trifolii* (Burgess) in tomatoes. *Res. Bulletin Aichi Ken Agric. Res. Center* (28): 177-187.
- Ignatowicz, S. (1979):** Effect of inorganic salts upon biology and development of acarid mites. III- Effect of relative humidity and mineral salts upon fecundity and longevity of copra mite, *Tyrophagus putrescentiae* (Schrank) (Acarina, Acaridae). *Bull. Entomol. Pol.*, 49: 577-581.
- Ismail, A. I.; Farag, N. A. and Ragaiei, M. A. (1999):** Evaluation of certain insecticides and plant extracts on *Bemisia tabaci*

- (Genn.) and their side effects on some natural enemies. *J. Egypt Ger. Soc. Zool.*, 28(E); *Entomology*: 117-124.
- Issa, S. and Marcano, R. (1994):** Population dynamics of *Liriomyza sativae* (Blanchard) and its parasites on tomato. *Turrialba*, 44(1): 24-30.
- Johnson, M. W. (1993):** Biological control of *Liriomyza* leafminers in the Pacific Basin. *Micronesica*, (4): 81-92.
- Kaya, N. and Hincal, P. (1991):** Survey studies on *Liriomyza trifolii* (Burgess) and *Phytomyza horticola* Goureaux (Diptera: Agromyzidae) found on leguminous plants in the Aegean Region. *Turkiye Entomoloji Dergisi*, 15(4): 241-246.
- Kerns, D. L. and Gaylor, M. J. (1992):** Sublethal effects of insecticides on cotton aphid reproduction and color morph development. *Southwestern Entomologist*, 17(3): 245-250.
- Kerns, D. L. and Stewart, S. D. (2000):** Sublethal effects of insecticides on the intrinsic rate of increase of cotton aphid. *Entomologia Experimentalis et Applicata*, 94(1): 41-49.
- Kim, D. I.; Paik, C. H.; Park J. D.; Kim, S. S. and Kim, S. G. (2000):** Relative toxicity of NeemAzal-T/S to the predacious mite, *Amblyseius womersleyi* (Acari: Phytoseiidae) and the two spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). *Korean Journal of Applied Entomology*, 39(1): 53-58.
- Knodel, J. J.; Larew, H. G. and Webb, R. E. (1986):** Margosan-O, a commercial formulation of neem seed extracts, controls *Liriomyza trifolii* (Burgess) on chrysanthemums. *J. of Agric. Entomol.* 3(3): 249-254. (c.f. R.A.E. A., 1993).

- Kumar, V. and Sharma, D.D. (1993):** Bio-ecology and chemical control of spider mites, *Tetranychus ludei* Zacher on okra. *Indian J. of plant Protection*, 21(1): 68 – 71.
- Kumawat, K. C. and Kumawat, G. L. (1995):** Effect of some insecticides on sucking pests and yellow mosaic incidence of mothbean, *Vigna aconitifolia* (Jacq.) Marechel. *Indian J. of Virology*, 11(2): 69-72.
- Kundu, G. G. and Trimopan (1992):** Preliminary observations of neem products against *Melanagromyza sojae* (Zehntner). *Pesticide Res. J.*, 4(1): 65-68.
- Kuroli, G.; Pocsai, K. and Nemeth, L. (1999):** Flight activity and population changes of aphids in faba bean. *Novenyvedelem*, 35(11): 545-554.
- Lal, O. P. (1992):** Evaluation of certain insecticides against the pest complex of French bean seed crop under field conditions. *J. of Entomological Res.*, 16(1): 57-61.
- Lal, S. S. and Pillai, K. S. (1982):** Ecological studies on whitefly, *Bemisia tabaci* (Genn.) infesting cassava in Kerala. *Entomol.*, 7(1): 101-102.
- Lal, S.S. (1984):** Influence of weather factors on the population of spider mites (Acari: Tetranychidae) and Thrips on Cassava in Kerala. *Indian J. Acarol.*, 7 (1): 5 – 10.
- Laska, P. (1986):** Test of Buprofezin, Cyhalothrin and other chemicals used against glasshouse whitefly. *Ann. of Appl. Biology*, 108: Suppl., *Tests of Agrochemicals and cultivars*, No. 7: 2-3.
- Leibee, G. L. (1984):** Influence of temperature on development and fecundity of *Liriomyza trifolii* (Burgess) (Diptera:

- Agromyzidae) on celery. *Environ. Entomol.*, 13(2): 494-501.
- Li, D. X.; Tiang, J. and Zhang, Z. Y. (1994):** A study on the bionomics and control techniques of *Aphis craccivora* Koch. on mung beans. *Plant Protection*, 20(6): 4-5.
- Linden, A. Van-der; Van-der, L. A. and Der-Linden, A. V. (1991):** Biological control of the leafminer *Liriomyza huidobrensis* (Blanchard) in Dutch glasshouse tomatoes. *Mededelingen Van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent.*, 56(2a): 265-271.
- Lloyd, B. J.; Ming, G. F.; Jennifer, E. F. and George, G. K. (1997):** Infection of the Migratory Grasshopper (Orthoptera: Acrididae) by ingestion of the entomopathogenic fungus *Beauveria bassiana*. *J. of Economic Entomol.*, 90(2): 383-390.
- Lourencao, A. L. and Nagai, H. (1994):** Outbreaks of *Bemisia tabaci* (Genn.) in Sao Paulo State. *Bragantia*, 53(1): 53-59.
- Lyon, J. P. (1986):** Specific problems posed by *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) and biological control of this new pest of protected crops. *Colloques de L'INRA* (34): 85-97.
- Mansour, N. A.; El-Defrawi, M. E.; Toppozada, A. and Zeid, M. (1966):** Toxicological studies on the Egyptian cotton leafworm, *Prodenia litura*. VI. Potential and antagonism of organophosphorus and carbamate insecticides. *J. of Econ. Entomol.*, 59(2): 307-311.
- Martell, C. G. (1968):** Some observations on the bionomics of *Liriomyza minutiseta* in Hawaii. *R. A. E. (A)* Vol. 56: 12 Abs. 2595 p.660.

- Mcclanahan, R. J. (1975):** Notes on the vegetable leafminer *Liriomyza sativae* (Diptera: Agromyzidae), in Ontario. *Proc. Entomol. Soc.*, (1974, publ. 1975), 105: 40-44, Ontario, Canada.
- Mehta, P. K. (1999):** New record of parasitoids of pea leafminer in Himachal Pradesh. *Insect Environment*, 4(4): 154-155.
- Meisner, J.; Yathom, S.; Tal, S. and Ascher, K. R. S. (1986):** The effect of various extracts of neem seed kernel on *Liriomyza trifolii* Burgess (Diptera: Agromyzidae). *Zeitschrift fur Pflanzenkrankheiten und Pflanzenschutz*, 93(2): 146-152.
- Milevoj, L. (1998):** The effect of some fungicides and insecticides on aphids (Aphididae). *Res. Rep., Biotechnical Faculty Univ. of Ljubljana, Agric. Issue*, 71: 21-27.
- Ministry of Agriculture (2003):** report of common bean production. *Agriculture Research Center, Dokki, Giza, Egypt*, NO. 777/2003.
- Mohamed, E. F.; Bahaa, A. O. and Mahmoud, A. Z. (1999):** Evaluation of Bemistop against whitefly *Bemisia tabaci* (Genn.) on tomato plants at Fayoum Governorate. *Fayoum J. Agric. Res. & Dev.*, 13(1): 179-184.
- Mohamed, R. A. (1997):** Effectiveness of datura leaves extracts and their mixtures with Malathion against the cowpea beetle, *Callosobruchus maculatus* (F). *Annals of Agric. Sci., Moshtohor*, 35(1): 589-604.
- Murguido, C. (1983):** Comparative effect of some organophosphorus insecticides for the control of the whitefly *Bemisia tabaci* (Genn.) and the leafhopper (*Empoasca* sp.) in bean crops. *Ciencia Y Tecnica en La*

Agricultura, Proteccion de Plantas, recod. 1985, 6(4) 59: 65.

- Nardo, E. A. B.; Costa, A. S.; Lourenvao, A. L. and De Nardo, E. A. B. (1997):** *Melia azedarach* extract as an antifeedant to *Bemisia tabaci* (Homoptera: Aphididae). *Florida Entomologist, 80(1): 92-94.*
- Nasser, M. A. K.; El-Ghareeb, A. M.; El-Sayed, A. M. K. and Mohamed, G. A. (2000):** Toxicological effects of some insecticides against susceptible and field populations of the cowpea aphid, *Aphis craccivora* Koch. *Assiut Journal of Agricultural Sciences. 2000, 31: 1, 217-230.*
- Nicoli, G. (1997):** Biological control of exotic pests in Italy: recent experiences and perspectives. *Bulletin OEPP 27(1): 69-75.*
- Nishijima, Y.; Honma, K. and Okuyama, S. (1963):** Mortality and its factors in the immature stages of the black wheat leafminer, *Agromyza albipennis* Meigen. *Res. Bull. Zotech. Univ. 1(3) No. 4 pp. 487-504.*
- Noldus, L. P. J. J.; XU, R.M. and Van, J.C. (1986):** The parasite-host relationship between *Encarsia formosa* Gahan (Hymenoptera: Aphelinidae) and *Trialeurodes vaporariorum* (West wood) (Homoptera: Aleyrodidae). XIX, Feeding-site selection by the greenhouse whitefly. *J. Apple-Entomol. 101: 492-507.*
- Oatman, E. R. and Michelbacher, A. E. (1958):** The melon leaf miner, *Liriomyza pictella* Thomson (Diptera: Agromyzidae). *Ann. Entomol. Soc. Amr. Vol. 51(6): 557-566.*

- Oatman, E. R. and Michelbacher, A. E. (1959):** The melon leaf miner, *Liriomyza pictella* Thomson (Diptera: Agromyzidae). II. Ecological studies. *Ann. Entom. Soc. Amr. Vol. 52(1): 83-89.*
- Oetting, R. D. (1986):** Methods of cyromazine application for leafminer, *Liriomyza trifolii*, control in glasshouses. 1986 *British Crop Protection Conf. Pests and diseases. Volume 1. Proceeding of a Conference held at Brighton Metropole, England, November (17-20): 263-269.*
- Ofuya, T. I. and Okuku, I. E. (1994):** Insecticidal effect of some plant extracts on the cowpea aphid *Aphis craccivora* Koch. (Homoptera: Aphididae). *Anzeiger fur Schadlingskunde, Pflanzenschutz, Umweltschutz, 67(6): 127-129.*
- Omar, B. A. and Faris, F. S. (2000):** Bio-residual activity of different insecticides on the leaf miners and yield components of snap bean *Phaseolus vulgaris* L. *Egypt J. of Agric. Res., 78(4): 1485-1497.*
- Omara, S. M.; Kelany, I. M.; Kleeberg, H. and Zebitz, C. P. W. (1997):** Effect of an aqueous neem seed kernel powder extract (ANSKPE) and Neem Azal-F on the *Liriomyza congesta* (Becker) and *Aphis craccivora* Koch. infesting broad bean at Zagazig Region, Sharkia Governorate, Egypt. *Practice oriented results on use and production of neem-ingredients and pheromones. Proceedings 5th Workshop Wetzlar, Germany, 22-25 Jan. 1996, 223-235.*
- Ozawa, A.; Kobayashi, H.; Amano, T.; Ikari, T. and Saito, T. (1993):** Evaluation of imported parasitic wasps as biological control agents of the legume leafminer, *Liriomyza trifolii* Burgess, in Japan. II. A field test on

cherry tomatoes in a plastic greenhouse, in Shizuoka Prefecture. *Proceedings of the Kanto Tosan Plant Protection Society*, 40: 239-241.

- Ozawa, A.; Saito, T. and Ikeda, F. (1998):** Effects of pesticides on *Diglyphus isaea* (Walker) and *Dacnusa sibirica* Telenga, parasitoids of *Liriomyza trifolii* Burgess. *Japanese J. Appl. Entomol. & Zoology*, 42(3): 149-161.
- Parkman, P. and Peinkowski, R. L. (1990):** Sublethal effects of neem seed extracts on adults of *Liriomyza trifolii* Burgess (Diptera: Agromyzidae). *J. Econ. Entomol.*, 83(4): 1246-1249.
- Parkman, P.; Dusky, J. A. and Waddill, V. H. (1989):** Leaf miner and leafminer parasitoid incidence on selected weeds in south Florida. *Florida Entomologist*, 72(3): 559-561.
- Parrella, M. P. (1984):** Effect of temperature on oviposition, feeding, and longevity of *Liriomyza trifolii* (Diptera: Agromyzidae). *Canada. Entomol.*, 116(1): 85-92.
- Parrella, M. P. and Bethke, J. A. (1988):** Larval development and leaf mining activity of *Liriomyza trifolii* Burgess (Diptera: Agromyzidae). *Pan-Pacific Entomol.*, 64(1): 17-22.
- Parrella, M. P.; Yost, J. T.; Heinz, K. M. and Ferrentino, G. W. (1989):** Mass rearing of *Diglyphus begini* (Hymenoptera: Eulophidae) for biological control of *Liriomyza trifolii* (Diptera: Agromyzidae). *J. Econ. Entomol.*, 82(2): 420-425.
- Patil, S. B.; Hugar, P. S.; Udikeri, S. S. and Kambar, N.S. (2001):** Chemical suppression of serpentine leafminer.

Karnataka-Journal-of-Agricultural-Sciences. 2001, 14: 3, 639-641.

- Pena, M. A. (1988):** First biological control experiments against *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) with *Diglyphus isaea* Walker (Hymenoptera: Eulophidae) in the Canary Islands. *Boletin de Sanidad Vegetal, Plagas*, 14(3): 439-445.
- Perkins, H. H. (1983):** Identification and processing of honeydew contaminated cotton. *Textile Research Journal* pp. 508-512.
- Perkins, H. H. (1987):** Sticky cotton. *Proceedings of the Western cotton production conference* pp. 53-55.
- Prieto, M. A. J. and Chaco De Ulloa, P. (1982):** Biology and ecology of the chrysanthemum miner *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) in the Department of Valle del Cauca. *Revista Colombiana de Entomologia* (1980, publ. 1992) 6(3/4): 77-84. (R.A.E. (A) 71:4 Abs. 2871 p. 337 (1983)).
- Puttaraju, H. R.; Prakash, H. S. and Shetty, H. S. (2002):** Contribution of seedborne Blackeye cowpea mosaic potyvirus to disease dynamics and loss of yield. *Tropical-Science*. 2002, 42: 4, 147-152.
- Richter, S. and Caceda, F. (1984):** Studies on improving the effectiveness of organophosphorus insecticides by combination with regulators of biological processes. *Archiv fur Phytopathologie und Pflanzenschutz*, 20(5): 409-418.
- Richter, S. and Tsegaye, Y. (1988):** Rearing *Liriomyza trifolii* Burgess on *Phaseolus vulgaris* L. in the laboratory and the

development of a test method. *Beitrage zur Tropischen Landwirtschaft und Veterinarmedizin*, 26(4): 387-394.

Robb, K. L. and Parella, M. P. (1985): Antifeeding and oviposition-detering effects of insecticides on adult *Liriomyza trifolii* Burgess (Diptera: Agromyzidae). *J. Econ. Entomol.*, 78: 709-713.

Roditakis, N. E.; Golfopoulou, N. G.; Albajes, R. (ed.) and Carnero, A. (1997): Bioecological studies on South American leafminer *Liriomyza huidobrensis* (Blanchard) in Crete. *Bulletin OILB/SROP*, 20(4): 225-231.

Saito, T.; Ikeda, F. and Ozawa, A. (1996): Effect of pesticides on parasitoid complex of serpentine leafminer *Liriomyza trifolii* Burgess in Shizuoka Prefecture. *Japanese J. Appl. Entomol. & Zoology*, 40(2): 127-133.

Saito, T.; Oishi, T.; Ikeda, F. and Sawak, T. (1992): Effect of insecticides on the serpentine leafminer *Liriomyza trifolii* Burgess (Diptera: Agromyzidae). *Japanese J. Appl. Entomol. & Zoology*, 36(3): 183-191.

Sammour, E. A. and El-Kabbany, S. M. (1993): Toxicological and biochemecal studies on acylurea, insecticides and their mixtures on the cowpea aphid, *Aphis craccivora* Koch.. *Bulletin of the Entomological Society of Egypt, Economic Series*, 20: 111-121.

Sarao, P. S.; Gursharan, S. and Singh, G. (1998): Sublethal influence of insecticides on reproduction of mustard aphid, *Lipaphis erysimi* (Kltenbach). *J. of Insect Sci.*, 11(1): 5-8.

Schuster, D. J. (1994): Life-stage specific toxicity of insecticides to parasitoids of *Liriomyza trifolii* Burgess (Diptera:

- Agromyzidae). *International J. Pest Management* 40(2): 191-194.
- Schuster, D. J. and Taylor, J. L. (1987):** Residual activity of Abamectin against *Liriomyza trifolii* Burgess (Diptera: Agromyzidae). *Fla. Entomol.* 70(3): 351-354. (C.f. *Entomology Abstract*, 19(2): 1663, 1988).
- Selim, A. A.; El-Refai, S. A.; El-Gantiry, A.; Refai, S. A. and Gantiry, A. E. (1987):** Seasonal fluctuations in the population of *Aphis craccivora* Koch., *Myzus persicae* Sulze., *Aphis gossypii* Glove. and their parasites. *Ann. of Agric. Sci., Ain Shams Univ.*, 32(3): 1837-1848.
- Serantes De Gonzalez, H. (1975):** *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae). *Revista de la Sociedad Entomologica Argentina* (1973, publ. 1975) Vol. 34 No. — pp. 207-216. (R.A.E. (A) 64:6 Abs. 3660 p. 1030 (1976)).
- Shahein, A. and El-Maghraby, M. M. A. (1988):** Studies on the hymenopterous parasitoids of *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) on broad beans in Egypt. *J. of Appl. Entomol.*, 106(4): 377-380.
- Shahein, A. and El-Maghraby, M. M. A. (1993):** Impact of the parasitoids of *Liriomyza trifolii* Burgess on broad bean. *Zeitschrift fur Angewandte Zoologie*, 79(1): 37-43.
- Shalaby, H.H. (1998):** Studies on the principal insect pests attacking common bean in the field. *M. Sc. Thesis, Fac. Agric., Zagazig Uni. (Benha Branch)*.
- Sharaf El-Din, N.; Kannan, H.; Babiker, A.; Weigand, S. and Saxena, M. C. (1997):** The leaf miner, *Liriomyza trifolii* Burgess (Diptera: Agromyzidae) a serious insect pest of faba bean (*Vicia faba* L.) in the Sudan. 1. Identification of

the pest, natural enemies and abundance. *Arab J. Plant Prot.* 15(1): 39-42.

- Sharama, R. P.; Yadav, R. P. and Singh, R. (1991):** Relative efficacy of some insecticides against the field population of bean aphid (*Aphis craccivora* Koch. and safety to the associated aphidophagous coccinellid complex occurring on Lathyrus, Lentil and chickpea crops. *J. of Entomol. Res.*, 15(4): 251-259.
- Sites, R.W. and Cone, W.W. (1985):** Vertical dispersion of two spotted spider mites on hops throughout the growing season. *J. Entomol. Soc. British Columbia*, 82: 22 – 25, U. S. A.
- Snedecor, G. W. and Cochran, W. G. (1967):** *Statistical Methods, Sixth edition, Iowa State College Press, Ames. Iowa, USA.*
- Spencer, K. A. (1973):** Agromyzidae (Diptera) of economic importance. *Series Entomologica, Vol. 9, Dr. W. Junk, B. V. The Hague, Netherlands*, 418 pp.
- Spollen, K.M.; Johnson, M.W. and Tabashnik, B.E. (1995):** Stability of Fenvalerate resistance in the leafminer parasitoid *Diglyphus begini* (Hymenoptera: Eulophidae). *J. Econ. Entomol.* 88(2): 192-197.
- Srinivasan, G. and Babu, P. C. S. (2001):** Field evaluation of neem products against whitefly, *Bemisia tabaci* Gennadius on brinjal. *Annals of Plant Protection Sciences.* 2001, 9: 1, 19-21.
- Srivastava, U. S. and Jaiswal, A. K. (1989):** Precedence II-induced effects in the aphid *Aphis craccivora* Koch.. *Insect Science and its Application*, 10(4): 471-475.

- Steinberg, S.; Gouldman, D.; Kaminski, L. and Bigler, F. (1991):** The effect of storage conditions on adult longevity, fertility and sex ratio of the leafminer parasitoid *Diglyphus isaea* (Walker). *Fifth Workshop of The IOBC Global Working Group on Quality Control of Mass Reared Arthropods*, 95-101.
- Stenseth, C. and Singh, H. M. (1990):** Buprofezin against the glasshouse whitefly and the cotton whitefly. *Gartneryrket*, 80(1): 18-19.
- Suss, L. (1975):** *Liriomyza nitzkei* Spencer (Diptera: Agromyzidae) injurious to onion crops (*Allium cepa*) in Italy. *Bollettino di Zoologia Agraria e-di Bachicoltura* (1974, Publ. 1975) Vol. 12 pp. 205-209, Italy. (R.A.E. (A) 64:10 Abs. 6174 p. 1718 (1976)).
- Thayaalini, S. and Raveendranath, S. (1997):** The efficacy of lakada (*Gardenia cramerii* Ait.) against cowpea aphid, *Aphis craccivora* Koch. (Homoptera: Aphididae). *Pest Management in Hort. Ecosystems*, 3(2): 66-69.
- Trefas, H. and Bujaki, G. (1997):** Experiments on the biology of *Diglyphus isaea*. *Novenyvedlem*, 33(9): 453-457.
- Tryon, E. H. and Poe, S. L. (1981):** Developmental rates and emergence of vegetable leafminer pupae and their parasites reared from celery foliage. *Flo. Entomol.*, 64(4): 477-483.
- Tuncer, C. and Aliniasee, M. T. (1998):** Acute and chronic effects of neem on *Myzocallis coryli* (Homoptera: Aphididae). *International J. of Pest Management*, 44(2): 53-58.

- Ulubilir, A.; Sekeroglu, E.; Albajes, R. and Carnero, A. (1997):** Dept of Plant Protection, CU Agricultural Faculty, 01330 Adana, Turkey. *Bulletin OILB/SROP*, 20(4): 232-235.
- Ushchekov, A. T. (1994):** *Diglyphus* as an efficient parasitoid of mining flies. *Zashchita Rastenii Moskva*, 3: 56-57.
- Ushchekov, A. T. (1996):** Rearing *Diglyphus*. *Zashchita Rastenii Moskva*, 2: 42.
- Vadodaria, M. P. and Vyas, H. N. (1987):** Control of whitefly, *Bemisia tabaci* (Gennadius) and its impact on yellow mosaic virus (YMV) in green gram *Vigna radiata* (L.) Wilczek and the grain yield. *Indian Journal of Agricultural Research*. 1987, 21: 1, 21-26.
- Veire, M. Van De and Van De Veire, M. (1991):** Progress in IPM in glasshouse vegetables in Belgium. *Bulletin SROP*, 14(5): 22-32.
- Vicente, M.; Kanthack, R. D.; Noronha, A. B. and Stradioto, M. F. S. (1988):** Incidence of golden mosaic on bean grown in two plant seasons in the Presidente Prudente Region. *Fitopatologia Brasileira*, 13(4): 373-376.
- Vishwanath, S. and Singh, J. P. (2002):** Bioefficacy of certain insecticidal formulations against bean aphid, *Aphis craccivora* Koch.. *Ann. of Plant Protection Sci.*, 10(1): 136-137.
- Webb, R. E. and Smith, F. F. (1969):** Effects of temperature on resistance in lima bean, tomato and chrysanthemum to *Liriomyza munda*. *J. Econ. Entomol.* 62(2): 458-462.
- Webb, R. E. and Smith, F. F. (1970):** Rearing a leafminer *Liriomyza munda*. *J. Econ. Entomol.* 63(6): 2009-2010.

- Webb, R. E.; Hinebaugh, M. A.; Lindquist, R. K. and Jacobson, M. (1983):** Evaluation of aqueous solution of neem seed extract against *Liriomyza sativae* and *L. trifolii* (Diptera: Agromyzidae). *J. Econ. Entomol.* 76(2): 357-362.
- Woets, J. and Van Der Linden, A. (1985):** First experiments on *Chrysocharis parksi* Crawford (Hymn.: Eulophidae) as a parasite for leafminer control *Liriomyza* spp. (Diptera: Agromyzidae) in European greenhouse tomatoes. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent*, 50(2b): 763-768.
- Wood, J. P.; Godfrey, L. D.; Dugger, P. and Richter, D. (1998):** Effects of whitefly insect growth regulators knack R and Applaud R on cotton aphid reproduction and survival. 1998 *Proceedings Beltwide cotton Conference, San Diego, California, USA, 5-9 January* 2: 1278-1281.
- Xue, M.; Li, Q. A.; Xue, M. and Li, Q. A. (2002):** Studies on selective toxicity of six insecticides between green peach aphid and ladybirds. *Entomologia Sinica*, 9(2): 17-22.
- Yasarakinci, N. and Hincal, P. (1999):** Studies on the protected bean pests, their natural enemies and their population developments in Izmir Province. *Bitki Koruma Bulteni*, 39(3-4): 137-150.
- Yathom, S.; Ascher, K. R. S.; Tal, S. and Nemny, N. E. (1986):** The effect of Trigard (Cyromazine) on different stages of the leafminer *Liriomyza trifolii* **Burgess**. *Hassaded*, 67(2): 372-376.
- Zchori, F. E.; Roush, R. T. and Sanderson, J. P. (1994):** Potential for integration of biological and chemical control of greenhouse whitefly (Homoptera: Aleyrodidae) using

Encarsia formosa (Hymenoptera: Aphelinidae) and Abamectin. *Environmental Entomology*, 23(5): 1277-1282.

Zoebisch, T. G. and Schuster, D. J. (1987): Longevity and fecundity of *Liriomyza trifolii* (Diptera: Agromyzidae) exposed to tomato foliage and honeydew in the laboratory. *Environmental Entomology*, 16(4): 1001-1003.