

REFERENCES

- Abbott, W. W. (1925). A method of computing the effectiveness of an insecticide.
J. Econ. Entomol., 18: 265-267.
- Adler, C. (1990). Resistance of different laboratory and field strain of the granary weevil *Sitophilus granarius* (L.) (Coleoptera: Curculionidae) to control with modified atmospheres.
Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie, 7: 434-438.
- Adler, C. and Reichmuth, C. (1988). The granary weevil *Sitophilus granarius* (L.). Coleoptera, Curculionidae, its biology and its control in grain, especially with modified atmosphere.
Mitteilungen aus der Biologischen Bundesanstalt für Land und Forstwirtschaft Berlin Dahlem, 239: 100pp.
- Ahmed, M. (1976). Effect of phosphine fumigation on the germination of edible legume seeds.
J. Stored Prod. Res., 12: 211-212.
- Ahmed, M.; Shaukat, S. S.; Alam, M. S.; Iqbal, S.; Ahmed, A. and Khan, J. (1993). Comparative cost effectiveness of four integrated pest management methods for storage of bagged wheat in Lahore and Multan.

- Sarhad J. Agriculture, 9(5): 403-414.
- Aliniaze, M. T. (1971).** The effect of carbon dioxide gas alone or in combinations on the mortality of *T. castaneum* (Herbst.) and *T. confusum* (Du Val) (Coleoptera: Tenebrionidae).
J. Stored Prod. Res., 7: 243-252.
- Annis, P. C.; Banks, H. J. and Sukardi, (1984).** Insect control in stacks of bagged rice using carbon dioxide treatment and an experimental PVC-membrane enclosure.
Division of Entomology Technical Paper No. 22; 44 ref.
- Anon, (1984).** Fumigation Standard No. 12 Bulletin OEPP/EPPO Bulletin, 23, 212.
- Aviles-Pacheco, R.; Stammler, C.; Herrera, A. and Pacheco-R-Aviles (1983).** Effectiveness of phosphine on three pests of unhusked rice, stored in sacks.
Ciencias de la Agricultura. No. 17, 51-56.
- Bailey, S. W. (1955).** Air-tight storage of grain: its effects on insect pests. 1. *Calandra granaria*.
Aust. J. Agric. Res., 6: 33-51.
- Bailey, S. W. and Banks, H. J. (1980).** A review of recent studies of the effects of controlled atmospheres on stored product pests.

Developments in Agric. Engineering I. Controlled atmosphere storage of grains, edited by Shejbal, J. pp. 101-118.

Banks, H. J. (1987). Disinfestation of durable foodstuffs in ISO containers using carbon dioxide.

ACIAR Proceedings Series, Australian Centre for International Agricultural Research, 23, 45-54.

Banks, H. J. and Sharp, A. K. (1979). Insect control with CO₂ in a small stock of bagged grain in a plastic film enclosure. Aust. J. Exp. Agric. Anim. Husb., 19: 102-107.

Barbara, D. H.; Bell, C. H. Mills, K. A. and Goodship, G. (1976). The toxicity of phosphine to all developmental stages of 13 species of stored product beetles. J. Stored Prod. Res., 12: 235-244.

Bell, C. H. (1976). The tolerance of developmental stages of four stored product Moths to phosphine. J. Stored Prod. Res., 12: 77-86.

Bell, C. H. (1979). The efficiency of phosphine against diapausing larvae of *Ephistia elutella*, over a wide range of concentration and exposure times. J. Stored Prod. Res., 15: 53-58.

Bell, C. H. (1984). Effect of oxygen on the toxicity of carbon dioxide to stored insects.

- Developments in Agric. Engineering 1: Controlled atmosphere storage of grains, edited by Ripp *et al.*, B. E., 67-74.
- Bell, C. H. and Glanville, V. (1970).** Toxicity of phosphine to insects. Toxicity to moths.
Pest insect. Res., 55-57.
- Bell, C. H. and Glanville, V. (1973).** The effect of concentration and exposure in tests with methyl bromide and with phosphine on diapausing larvae of *Ephestia elutella* (Hubner) (Lepidoptera: Pyralidae).
J. Stored Prod. Res., 9: 165-170.
- Bell, C. H. and Mills, K. A. (1983).** Special factors affecting the toxicity of fumigants to insects.
Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie, 4: 1-3, 120-122.
- Bell, C. H.; Hole, B. D. and Wilson, S. M. (1985).** Fumigant doses for the control of *Trogoderma granarium*.
Bulletin, Organisation Europeenne et Mediterraneenne pour la Protection des Plantes, 15: 1, 9-14.
- Bell, C. H.; Wilson, S. M. and Banks, H. J. (1984).** Studies on the toxicity of phosphine to tolerant stages of *Trogoderma granarium* Everts (Coleoptera: Dermestidae).
J. Stored Prod. Res., 20: 2, 111-117.

- Bell, C. H.; Wilson, S. M.; Banks, H. J. and Smith, R. H.** (1984). An investigation of the tolerance of stages of Khapra beetle (*Trogoderma granarium*, Everts) to phosphine.
Proceedings of the Third International Working Conference on Stored-Product Entomology: October 23-28, 1983, Kansas State University, Manhattan, Kansas USA, 329-340.
- Bond, E. J.** (1980). Sorption of tritiated phosphine by various stages of *Tribolium castanum*.
J. Stored Prod. Res., 16:27-31.
- Bond, E. J. and Monro, H. A. U.** (1967). The role of oxygen in the toxicity of fumigants to insects.
J. Stored Prod. Res., 3: 295-310.
- Bond, E. J.; Robinson, J. R. and Buckland, C. T.** (1969). The toxic action of phosphine: absorption and symptoms of poisoning in insects.
J. Stored Prod. Res., 5: 289-298.
- Bond, E. J. and Buckland, C. T.** (1978). Control of insects with fumigants at low temperatures: toxicity of fumigants in atmosphere of CO₂.
J. Econ. Entomol., 71:307-309.
- Bond, E. J. and Buckland, C. T.** (1979). Development of resistance of carbon dioxide in the Germany weevil.

- J. Econ. Entomol., 72:770-771.
- Bond, E. J. and Monro, H. A. U. (1961). The toxicity of various fumigants to the cadelle *Tenebroides mauritanicus*.
J. Econ. Entomol., 54: 451-454.
- Bond, E. J.; Sellen, R. A and Dumas, T. (1977). Control of insects with phosphine in open-ended bin spouts.
J. Econ. Entomol., Vol. 70, No.1, 22-25.
- Buscarlet, L. A. (1993). Study on the influence of temperature on the mortality of *Tribolium confusum* du val. Exposed to carbon dioxide or nitrogen.
Zeitschrift fur Naturforschung. Section-C, Biosciences, 48:7-8.
- Calderon, M. and Leesch, J. G. (1983). Effect of reduced pressure and CO₂ on the toxicity of methyl bromide to two species of stored-product insects.
J. Econ. Entomol., 76: 5, 1125-1128.
- Calderon, M. and Navarro, S. (1979). Increased toxicity of low oxygen atmospheres supplemented with CO₂ on *Tribolium castaneum* adults.
Ent. Exp. Appl., 25: 39-44.
- Calderon, M. and Navarro, S. (1980). Synergistic effect of CO₂ and O₂ mixtures on two stored grain insect pests.

- Developments in Agric. Engineering I: Controlled atmosphere storage of grains, edited by Shejbal. J. pp.79-84-Italy.
- Campos, T. B.; Lamonica-Imenes, S.; Bergmann, C. E.; Xavier, N. J. D.; Santiago, I. M.; De-Campos, T. B. and De-Lamonica-Imenes, S. (1995).** Evaluation of the use of phosphine in the protection of cotton seeds against infestations of *Tribolium castaneum* Herbst. (Coleoptera-Tenebrionidae).
Revista de Agricultura Piracicaba., 70: 1, 7-15.
- Carlson, S. D. (1966).** Fumigation of *T. confusum* adults with 80: 20 (CCL₄: CS₂) during CO₂ or N₂ induced respiration elevation or dpresion.
J. Econ. Entomol., 59: 870-872.
- Childs, D. P. and Overby, J. E. (1983).** Mortality of the cigarette beetle in high-carbon dioxide atmosphere.
J. Econ. Entomol., 76: 544-546.
- Contessi, A. (1999).** Cereal preservation by using controlled atmosphere produced where the corn is stored.
Tecnica Molitoria, 50: 9, 990-999.
- Corinth, H. G. and Rau, G. (1990).** Protection of stored grain by means of carbon dioxide.
Anzeiger für Schadlingskunde, Pflanzenschutz, Umweltschutz. 63: 7, 121-123.

- Corues, M. A. and Adeyemi A. O. (1969). A comparison of phosphine and ethylene dibrom for fumigation of cowpea in polyethylene lined hessian sacks.
Bulletin of the Entomological Society of Nigeria, 2(1): 45-50.
- Darwish, A. A. (1998). Rol of carbon dioxide on the toxicity of phosphine to larvae of two-stored product insects.
Annals of Agric. Sc. Moshtohor, Vol. 36(2): 1269-1285.
- Desmarchelier, J. M. (1984). Effect of carbon dioxide on the efficacy of phosphine against different stored product insects.
Mitteilung aus der Biologischen Bundesanstalt für land und Forstwirtschaft Berlin-Dahlem Heft 220 Juli, 1984.
- Desmarchelier, J. M.; Wohlgemuth, R. and Ripp, B. E. (1984). Response of several species of insects to mixtures of phosphine and carbon dioxide.
Controlled atmosphere and fumigation in grain storages. Proceedings of an international symposium 'Practical aspects of controlled atmosphere and fumigation in grain storages' held from 11 to 22 April 1983 in Perth, Western Australia. 1984, 75-81.
- Dias, R. and Navarro, S. (1983). Induced tolerance of *Sitophilus oryzae* adults to carbon dioxide.

- Progress report for the year 1981-82 of the Stored Products Division. 1983, 27-33.
- Donahaye, E. and Zalach, D. (1987).** Comparison of sensitivity of developmental stages of three strains of the flour beetle *Tribolium castaneum* Herbst. to modified atmospheres. Progress report for the year 1984-85 of the Stored Products Division, Institute for Technology and Storage of Agricultural Products, Israel. 1987, 3-23.
- Donahaye, E. J.; Navarro, S.; Rindner, M. and Azrieli, A. (1996).** The combined influence of temperature and modified atmospheres on *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae). J. Stored Prod. Res., 32: 3, 225-232.
- El-Lakwah, F. A. (1990).** Fumigation experiments with phosphine in traditional mud silos in Egypt to control stored product insects. Egypt. J. Appl. Sci., 5(3): 51-65.
- El-Lakwah, F. A. and Kattab, M. M. (1989).** Effect of phosphine fumigation on germination of seeds. J. Agric. Sci. Mansoura Univ. 14 (2): 794-796.
- El-Lakwah, F. A. and Halawa, Z. A. (1998).** Efficacy of a modified atmosphere of nearly 99% Nitrogen plus 1% Oxygen on two species of stored products pests. Annals of Agric. Sc. Moshtohor, vol. 36(4): 2565-2572.

- El-Lakwah, F. A.; Darwish, A. A. and Mohamed, R. A. (1994). Effect of carbon dioxide on the different stages of *Callosobruchus chinensis* (L.) and *Sitophilus granarius* (L.).
Annals of Agric. Sc. Moshtohor, Vol. 32(2): 1009-1018.
- El-Lakwah, F. A.; Darwish, A. A. and Mohamed, R. A. (1997). Efficacy of a modified atmosphere of around 1% oxygen plus 99% nitrogen on some stored product insects.
Annals of Agric. Sc. Moshtohor, Vol. 35(1): 579-587.
- El-Lakwah, F. A.; Hamed, M. S. and Abd-El-Aziz, A. E. (1998). Toxicity and joint action of neemazal with certain modified atmospheres against two species of stored product insects.
Annals of Agric. Sc. Moshtohor, Vol. 36(3): 1941-1958.
- El-Lakwah, F. A.; Khaled, O. M. and Halawa, Z. A. (1996). Effect of modified atmosphere of more than 99% nitrogen or enriched with carbon dioxide on mortality of some stored product insects, seed germination, chlorophyll and carotene contents of certain crop seedlings.
Annals of Agric. Sc. Moshtohor, Vol. 34 (4): 1869-1877.
- El-Lakwah, F. A.; Wohlgemuth, R. and Khattab, M. M. (1989). Efficiency of phosphine and combinations of phosphine with carbon dioxide against Khapra beetle larvae *Trogoderma granarium* Everts (Coleoptera, Dermestidae).

Anzeiger für Schadlingskunde, Pflanzenschutz,
Umweltschutz, 62: 5, 85-88.

El-Lakwah, F. A.; Darwish, A. A.; Omnia, M. Khaled and
El-Sayed, H. M. (1998). Efficiency of controlled
atmospheres containing various carbon dioxide
concentrations and nitrogen against the Khapra beetle
larvae *Trogoderma granarium* Everts. (Coleoptera:
Dermestidae).

Annals of Agric. Sc. Moshtohor, Vol. 36(1): 619-641.

El-Lakwah, F. A.; Saleh, M. K.; Omer, E. E. and Mohamed,
R. A. (1992a). Efficiency of phosphine alone and in
combinations with carbon dioxide against the various
stages of cowpea weevil *Callosobruchus maculatus* (F.)
Coleoptera, Bruchidae.

Egypt. J. Appl. Sci. 7(11), 80-98.

El-Lakwah, F. A.; Saleh, M. K.; Omer, E. E. and Mohamed,
R. A. (1992b). Efficiency of phosphine, carbon dioxide and
their mixtures against the various stages of lesser grain
borer (*Rhizopertha dominica*) (F.) Coleoptera,
Bostrychidae.

Egypt. J. Appl. Sci. 7(11), 99-117.

- El-Lakwah, F. A.; Saleh, M. K.; Omer, E. E. and Mohamed, R. A. (1992c). Efficiency of phosphine, carbon dioxide and their mixtures against the various stages of rice weevil *Sitophilus oryzae* (L.) Coleoptera, Curculionidae. Egypt. J. Appl. Sci. 7(11), 203-221.
- El-Lakwah, F. A.; Abdel Gawad, A.; Meuser, F.; Wohlgemuth, R. and Darwish, A. (1991a). Efficiency of phosphine alone and in mixtures with carbon dioxide against the adult of *Tribolium castaneum* and *Sitophilus oryzae*. J. Plant Diseases and Protection, 98(1), 92-102.
- El-Lakwah, F. A.; Meuser, F.; Abdel Gawad, A.; Wohlgemuth, R. and Darwish, A. (1991b). Efficiency of phosphine alone and in mixtures with carbon dioxide against Angoumois Grain Moth *Sitotroga cerealella* (Olivier); (Gelechiidae; Lepidoptera). J. Plant Diseases and Protection, 98(1), 92-102.
- El-Nahal, A. K. M.; Barakat, A. A.; El-Halafawy, M. A. and Hassan, H. (1986). Susceptibility of natural strains of *Sitophilus oryzae* (L.), *S. granarius* (L.), *Rhizopertha dominica* (F.) and *Tribolium castaneum* (Herbst.) collecting from the different governorates of Egypt to Methyl bromide and phosphine. Bull. Ent. Soc. Egypt. Econ. Ser., 5(14): 141-154.

Finney, D. J. (1971). Probit Analysis, 3rd ed.

Cambridge University Press.

Fleurat-Lessard, F.; Torc'-h, J. M. and Le-Torc'-h, J. M.

(1991). The influence of oxygen content in atmospheres enriched with carbon dioxide on the mortality of the juvenile stages of *Sitophilus oryzae* and *S. granarius*.

Entomologia Experimentalis et Applicata, 58: 1, 37-47.

Gibe, A. J. G.; Sayaboc, P. D.; Açda, M. A.; Caliboso, F. M.;

Sidik, M.; Rejesus, B. M.; Garcia, R. P.; Champ, B. R.;

Bengston, M.; Dharmaputa, O. S. and Halid, H. (1997).

Resistance of four major coleopterous stored grain insects to phosphine in selected areas in the Philippines.

Proceedings of the symposium on pest management for stored food and feed, held at Bogor, Indonesia, 5-7 September 1995. BIOTROP-Special-Publication. 1997, No. 59, 231-236.

Goto, M.; Kishino, H.; Imamura, M.; Hirose, Y. and Soma,

Y. (1996). Responses of the pupae of *Sitophilus granarius* L., *Sitophilus zeamais* Motschulsky and *Sitophilus oryzae* L. to phosphine and mixtures of phosphine and carbon dioxide.

Research Bulletin of the Plant Protection Service, Japan. No. 32, 63-67.

- Halawa, Z. A. (1998). Efficacy of a modified atmosphere on certain stored product pests.
Egypt. J. Agric. Rec., 76: (1), 95-103.
- Harein, P. K. and Press, A. F. (1968). Mortality of stored-peanut insects exposed to mixtures of atmospheric gases at various temperatures.
J. Stored Prod. Res., 4:77-82.
- Hashem, M. Y. (1993). Studies on the responses of the eggs of three curculionids to controlled atmospheres.
Bull. Ent. Soc. Egypt, Econ. Ser., 20: 39-46.
- Hashem, M. Y. (1998). Suggested procedures for applying carbon dioxide CO₂ to disinfest stored medicinal plant products from insect pests.
Bull. Ent. Egypt, Econ. Ser., 25: 107-115.
- Hashem, M. Y. and El-Sayed, M. R. (1998). Toxicity of high carbon dioxide atmospheres to various stages of *Callosobruchus chinensis* (L.).
Bull. Ent. Soc. Egypt, Econ. Ser., 25: 99-106.
- Hashem, M. Y. and Reichmuth, Ch. (1989). The efficiency of phosphine against eggs of lesser grain borer *Rhizopertha dominica* (F.) and larger grain borer *Prostephanus truncatus* (Horn.) (Coleoptera: Bostrichidea).
Nachrichtenble. Deut. Pflanzenschutzd. (10): 159-163.

Hashem, M. Y. and Reichmuth, Ch. (1992). Investigation on the response of the adults of the larger grain borer *Prostephanus truncatus* (Horn.) and lesser grain borer *Rhizopertha dominica* (F.) (Coleoptera: Bostrichidea) to controlled atmospheres.

Zeitschrift für Angewandte Zoologie, 4: 469-476.

Hashem, M. Y. and Reichmuth, Ch. (1994). Interaction effects of high carbon dioxide or low oxygen atmospheres and temperature on hatchability of eggs of three stored product moths.

J. Plant Diseases and Protection, 101 (2): 178-182.

Hashem, M. Y. and Reichmuth, Ch. (1996). Responses of different species of stored product insects to mixtures of carbon dioxide and/or nitrogen in air.

Bull. Ent. Soc. Egypt, Econ. Ser., 23: 86-91.

Hashem, M. Y.; Ismail, I. I. and Omar, E. E. (1995). Treatment of the cowpea weevil *Callosobruchus maculatus* F. (Bruchidae: Coleoptera) with low oxygen atmospheres containing carbon dioxide.

Egypt. J. Agric. Rec., 73(1), 127-133.

Hashem, M. Y.; Risha, E. M. and Sharaf El-Din, A. A. (1993). A method for controlling stored product insects by changing the surrounding atmospheres.

J. Egypt. Ger. Soc. Zool., 12 (D): 323-333.

- Howe, R. W. (1973). The susceptibility of the immature and adult stages of *Sitophilus granarius* to phosphine. J. Stored Prod. Res., 8, 241-262.
- Highland, H. A.; Leesch, J. G.; Cline, L. D. and Zehner, J. M. (1984). Phosphine fumigation of thick-film polyethylene food bags and laminated film food packets. J. Econ. Entomol., 77: 4, 1041-1045.
- Iqbal, J.; Irshad and Khalil, S. K. (1993). Sack fumigation of wheat under polyethylene sheets. Sarhad J. Agriculture, 9: 5, 399-402.
- Ismail, I. I.; Omar, E. E. and Hashem, M. Y. (1995). The effects of sublethal concentrations of carbon dioxide (CO₂) in atmosphere on some biological aspects of the cowpea weevil *Callosobruchus maculatus* F. (Bruchidae: Coleoptera). Egypt. J. Agric. Rec., 73(1), 111-118.
- Jay, E. G.; Redlinger, L. M. and Laudani, H. (1970). The application and distribution of CO₂ in a peanut silo for insect control. J. Stored Prod. Res. 6:247-254.
- Jones, R. N. (1938). Toxicity of fumigant carbon dioxide mixtures to the Red flour beetle. J. Econ. Entomol., 31: 298-309.

Kashi, K. P. (1982). Dose mortality responses of 5 species of stored product insects to phosphine.

Int. Pest Control, 24(2): 46-49.

Kashi, K. P. and Bond, E. J. (1975). The toxic action of phosphine: Role of CO₂ on the toxicity of phosphine to *Sitophilus granarius* and *Tribolium confusum*.

J. Stored Prod. Res., 11:9-15.

Krishnamurthy, T. S.; Muralidharan, N. and Krishnakumari, M. K. (1993). Disinfesting food commodities in small storages using carbon dioxide rich atmospheres.

International Pest Control, 35: 6, 153-156.

Krishnamurthy, T. S.; Spratt, E. C. and Bell, C. H. (1986). The toxicity of carbon dioxide to adult beetles in low oxygen atmospheres.

J. Stored Prod. Res., 22: 3, 145-151.

Krishnasamy, V. and Seshu, D. V. (1990). Phosphine fumigation influence on rice seed germination and vigour.

Crop Sci., 30, 82-85.

Lima, C. P. F.; De-Lima, C. P. F. and Ripp, B. E. (1984). Minimal fumigant requirements for long-term airtight storage of grain.

Controlled atmosphere and fumigation in grain storages. Proceedings of an international symposium 'Practical

- aspects of controlled atmosphere and fumigation in grain storages' held from 11 to 22 April 1983 in Perth, Western Australia. 1984, 665-671.
- Lindgren, D. L. and Vincent, L. E. (1966).** Relative toxicity of hydrogen phosphide to various stored-product insects. J. Stored Prod. Res., 2:141-146.
- Lindgren, D. L. and Vincent, L. E. (1970).** Effect of atmospheric gases alone or in combination on the mortality of granary and rice weevils. J. Econ. Entomol., 63:1926-1929.
- Lindgren, D. L. and Vincent, L. E. and Strong, R. G. (1958).** Studies on hydrogen phosphide as a fumigant. J. Econ. Entomol., 51:900-903.
- Locatelli, D. P. and Daolio, E. (1991).** Effectiveness of CO₂ in a vacuum autoclave at different temperatures on *Rhizopertha dominica* (F.) and *Sitophilus oryzae* (L.). Tecnica Molitoria, 42(7): 605-610.
- Marzke, F. O. and Pearman, G. C. (1970).** Mortality of red flour beetle and Indian-meal moth larvae in simulated peanut storages purged for short periods with CO₂ or N₂. J. Econ. Entomol., 63: 817-819.

Navarro, S. and Calderon, M. (1974). Exposure of *Ephestlia cautella* pupae to CO₂ concentrations at different relative humidities the effect on adult emergence and loose in weight.

J. Stored Prod. Res., 10: 237-241.

Navarro, S. and Calderon, M. (1983). The sensitivity of *Tribolium castaneum* (Herbst.) eggs to controlled atmospheres.

Progress report for the year 1981-82 of the Stored Products Division. 1983, 41-50, VII; 2 fig.; 18 ref.

Navarro, S. and Dias, R. (1987). Effect of carbon dioxide on active and quiescent larvae of *Trogoderma granarium* at three temperatures.

Progress report for the year 1984-85 of the Stored Products Division, Institute for Technology and Storage of Agricultural Products, Israel. 1987, 48-56.

Navarro, S. and Jay, E. G. (1987). Application of modified atmospheres for controlling stored grain insects.

Monograph, British Crop Protection Council. 1987, No. 37, 229-236; In Stored Products Pest Control, Proceedings of a Symposium held at the University of Reading, 25-27 March, 1987; 16 ref.

Navarro, S.; Dias, R. and Donahaye, E. (1985). Induced tolerance of *Sitophilus oryzae* adults to carbon dioxide.

- Mansour, N. A.; El-Defrawi, M. E; Toppozada, A. and Zeid, M. (1966).** Toxicological studies on the Egyptian cotton leaf worm, *Prodenia litura*. Potentiation and antagonism of organophosphorus and carbamate insecticides.
J. Econ. Entomol., 59: 307-311.
- Mills, K. A.; Clifton, A. L.; Chakrabarti, B. and Savvidou, N. (1990).** The impact of phosphine resistance on the control of insects in stored grain by phosphine fumigation.
Proceedings of the Brighton Crop Protection Conference, Pests and Diseases. No. 3, 1181-1187.
- Monro, H. A. U. (1969).** Manual of fumigation for insect control.
FAO Agricultural studies no. 79: 381pp.
- Morner, J.; Hyvonen, A. and Simonsson, M. (1987).** Experiments with carbon dioxide against stored products pests.
Vextskyddsnotiser, 51(3):87-92.
- Nakakita, H. and Kuroda, J. (1986).** Differences in phosphine uptake between susceptible and resistant strains of insects.
J. Pesticide Sci., 11: 1, 21-26.
- Nakakita, H.; Saito, T. and Iyatomi, K. (1974).** Effect of phosphine on the respiration of adult *Sitophilus zeamais*.
J. Stored Prod. Res., 10: 87-92.

- J. Stored Prod. Res., 21: 4, 207-213.
- Newton, J.; Wildey, K. B. and Robinson, W. H. (1993).** Carbon dioxide as a fumigant to replace methyl bromide in the control of insects and mites damaging stored products and artifacts.
Proceedings of the 1st International Conference on Insect Pests in the Urban Environment, 329-338.
- Noack, S. and Reichmuth, Ch. (1978).** Einrechnerisches Verfahren Zur Bestimmung von beliebigen Dosis-Werten eines Wirkstoffes aus empirisch Dosis-Wirkungs-Daten-Mitt.
Biol. Bundesanstalt für Land. Forstwirtsch., Berlin Dahlem, Haft 185: 1-49.
- Ofuya, T. I. and Reichmuth, Ch. (1993).** Mortality of the bean bruchid, *Acanthoscelides obtectus* (Say), in some high carbon dioxide atmospheres.
J. Plant Diseases and Protection, 100 (2): 165-169.
- Omar, E. E.; Hashem, M. Y. and Ismail, I. I. (1995).** Effect of carbon dioxide in air on different stages of the rice weevil *Sitophilus oryzae* (L.) (Curculionidae: Coleoptera).
Egypt. J. Agric. Rec., 73(1), 119-126.
- Othman, K. S. A. (1990).** Toxicological studies of phosphine against the lesser grain borer, *Rhizopertha dominica* (F.) (Coleoptera: Bostrychidae).

- Bollettino di Zoologia Agraria e di Bachicoltura, 22(1): 7-15.
- Ozar, A. I. (1974). Investigation on the effect of phostoxin on *Bruchus lemtis* Frohl and *Callosobruchus maculatus* F. Bitki Karuma Bulteni, 14(1): 43-54.
- Pike, V. (1994). Laboratory assessment of the efficacy of phosphine and methyl bromide fumigation against all life stages of *Liposcelis entomophilus* (Enderlein). Crop Protection, 13: 2, 141-145.
- Price, L. A. and Mills, K. A. (1988). The toxicity of phosphine to the immature stages of resistant and susceptible strains of some common stored product beetles, and implications for their control. J. Stored Prod. Res., 24: 1, 51-59.
- Prozell, S. and Reichmuth, C. (1992). Efficacy of controlled atmospheres on *Sitophilus granarius* Linne, *Sitophilus oryzae* Linne, *Oryzaephilus mercator* Fauvel, *Glycyphagus domesticus* de Geer and *Ephesia kuehniella* Zeller. Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie, 8: 1-3, 265-272.
- Quinlan, J. K. and Pierce, R. (1983). Fumigating empty metal grain bins. Agricultural Research, USA. 1983, 32: 2, 14.

- Qureshi, A. H.; Bond, E. J. and Monro, H. A. U. (1965).
Toxicity of hydrogen phosphide to the granary weevil
Sitophilus granarius and other insects.
J. Econ. Entomol., 58: 324-331.
- Rai, L.; Sarid, J. N. and Pinagale, S. V. (1962). Fumigation of
food grains in India with hydrogen phosphide. I. Tests in
concrete bins.
Bull. Grain Tech., 1: 3-17.
- Rajendran, S. (1990). The toxicity of phosphine, methyl
bromide, 1,1,1-trichloroethane and carbon dioxide alone
and as mixtures to the pupae of red flour beetle, *Tribolium*
castaneum Herbst.
Pesticide Sci., 29: 1, 75-83.
- Reichmuth, Ch. (1984). The effect of the fumigant hydrogen
phosphide (phosphine) on stored product pests.
German Biological Research Center/Institute for Stored
Products Protection.
- Reichmuth, Ch. (1986). The significance of changing
concentrations in toxicity of phosphine. "GASGA seminar
on fumigation technology in developing countries".
Tropical development and research institute, storage
department, Slough, 18-21 March 1986, 88-98.

- Reichmuth, Ch. And Ofuya, T. (1992). Low oxygen atmospheres for the control of *Callosobruchus maculatus* (F.) and *Acanthoscelides obtectus* (Say). Proc. Int. Conf. Controlled Atmosphere and Fumigation in Grain Storage, Winnipeg, Canada, June 1992, Caspit Press Ltd, Jerusalem, pp. 115-120.
- Reynolds, E. M.; Robinson, J. M. and Howells, C. (1967). The effect on *Sitophilus granarius* (L.) (Coleoptera: Curculionidae) of exposure to low concentrations of phosphine. J. Stored Prod. Res., 2: 177-186.
- Scudamore, K. A. and Goodship, G. (1986). Determination of phosphine residues in fumigated cereals and other food-stuffs. Pestic. Sci., 17, 385-395.
- Singh, K. N. and Srivastava, B. P. (1983). Relative toxicity of various fumigants to the adults of two species of *Callosobruchus* and the susceptibility of different sexes to these fumigants. Indian. J. Ent., 45:130-138.
- Soderstrom, E. L. and Brandl, D. G. (1984). Modified atmospheres for post-harvest insect control in tree nuts and dried fruits.

- Proceedings of the Third International Working Conference on Stored Product Entomology. October 23-28, Kansas State University, Manhattan, Kansas USA., 487-497.
- Sowunmi, E. O. (1982). Effect of storage length and insecticide treatment on cowpea (*Vigna unguiculata* L. Walp). Tropical Grain Lagume Bulletin, 25:21-27.
- Spratt, E. C. (1984). The effects of atmospheres containing 5 - 20% oxygen with and without 10% carbon dioxide on the productivity of *Tribolium castaneum* (Herbst.) and *T. confusum* Du Val (Coleoptera: Tenebrionidae). J. Stored Prod. Res., 20: 4, 199-203.
- Spratt, E.; Dignan, G. and Banks, H. J. (1985). The effects of high concentrations of carbon dioxide in air on *Trogoderma granarium* Everts (Coleoptera: Dermestidae). J. Stored Prod. Res., 21: 1, 41-46.
- Srivastava, J. L. (1985). Tamil NADU Agric. Univ., Coimbatore (India), pp. 205-207.
- Srivastava, J. L.; Regupathy, A. and Jayaraj, S. (1985). Use of controlled atmosphere for the control of stored product insects. Behavioural and physiological approaches in pest management, 205-207.

- Stahl, E. and Rau, G. (1985). A new method of disinfection.
Anzeiger für Schadlingskunde, Pflanzenschutz,
Umweltschutz., 58: 7, 133-136.
- Storey, C. L. (1977). Effect of low oxygen atmosphere on
mortality of red and confused flour beetle.
J. Econ. Entomol., 70: 253-255.
- Stoyanova, S. and Shikrenov, D. (1976). Storage of cereals in
an atmosphere with a high CO₂ concentration. Effect of 20
and 40% CO₂ on insect pests.
Development in Agric. Engineering. I- Controlled
atmosphere storage of grains, edited by Shejbal, J. pp. 101-
118-Italy.
- Strong, R. G. and Lindgren, D. L. (1960). Germination of
small legume seeds after fumigation with hydrogen
phosphide.
J. Econ. Entomol., 53: 1-4.
- Suss, L.; Locatelli, D. P. and Frati, M. (1991). The use of
carbon dioxide in cereal pest control.
Tecnica Molitoria, 42: 4, 333-338.
- Torc'-h, J. M.; Fleurat, Lessard, F. and Le-Torc'-h, J. M.
(1991). A study of the adsorption-desorption interactions
between grain and atmospheres enriched with carbon
dioxide, for insect control in rice paddy.
Agronomie, 11: 4, 305-314.

- Tunc, I. (1983). Mortality of *T. confusum* adults in various atmospheric gas compositions.
Zeitschrift für angewandte Entomologie. Sonderdruck aus Bd. 95, H. 3, S. 263-267.
- Tunc, I. and Navarro, S. (1983). Sensitivity of *Tribolium castaneum* eggs to modified atmospheres.
Entomologia Experimentalis et Applicata, 34: 3, 221-226.
- Udeaan, A. S. (1990). Susceptibility status of *Trogoderma granarium* Everts populations to phosphine in Punjab.
Indian J. Ecology, 17(2): 195-196.
- Valentini, S. R. T.; Gomes, R. A. R.; Athie, I.; Castro-MFPPM-de; Verdade- (1997). Efficiency of PVC and polyethylene sheets for the fumigation of grains with phosphine.
Revista Brasileira de Armazenamento, 22: 1, 3-8.
- Vincent, L. E. and Lindgren, D. L. (1972). Toxicity of phosphine to the life stages of four species of dermestidae.
J. Econ. Entomol., 65:1429-1431.
- Vincent, L. E. and Lindgren, D. L. (1975). Toxicity of phosphine and methyl bromide at various temperatures and exposure periods to the four metamorphic stages of *Trogoderma variable*.
J. Econ. Entomol., 68(1):53-56.

- Vincent, L. E. and Lindgren, D. L. (1977). Toxicity of methyl bromide and phosphine at various temperatures and exposure periods to the metamorphic stages of *Lasioderma serricorne*.
J. Econ. Entomol., 70(4):497-500.
- White, N. D. G. and Jayas, D. S. (1991). Control of insects and mites with carbon dioxide in wheat stored at cool temperatures in non-airtight bins.
J. Econ. Entomol., 84:1933-1942.
- Wigglesworth, V. B. (1965). The principles of insect physiology.
Menthuen, London (C. F. Bond and Monro, 1967)
- Winks, R. G. (1973). Some aspects of the response of *T. castaneum* to phosphine.
Ph. D. Thesies, University of London.
- Winks, R. G. (1982). The toxicity of phosphine to adults of *T. castaneum*: Time as a response factor.
J. Stored Prod. Res., 18: 159-169.
- Winks, R. G. (1984). The toxicity of phosphine to adults of *Tribolium castaneum* (Herbst.): Time as a dosage factor.
J. Stored Prod. Res., 20: 1, 45-56.
- Winks, R. G. (1985). The toxicity of phosphine to adults of *Tribolium castaneum* (Herbst.): Phosphine-induced narcosis.

- J. Stored Prod. Res., 21: 1, 25-29.
- Winks, R. G. (1986).** The significance of response time in the detection and measurement of fumigant resistance in insects with special reference to phosphine.
Pesticide Science, 17: 2, 165-174.
- Winks, R. G. and Waterford, C. J. (1986).** The relationship between concentration and time in the toxicity of phosphine to adults of a resistant strain of *Tribolium castaneum* (Herbst.).
J. Stored Prod. Res., 22: 2, 85-92.
- Zakladnoi, G. A. (1976).** Regulation of the gas compositions of the atmosphere for eliminating insects in grain.
Development in Agric. Engineering. I- Controlled Atmosphere Storage of Grains, edited by Shejbal, J., pp. 101-118.