

REFERENCES

- Allard, R.W., 1960. Principles of Plant Breeding. John Wiley & Sons, Inc., New York, pp. 485.
- Barton, D.W., Butler L., Jenkins, J.A., Rick, C.M., and Young, P.A., 1955. Rules for nomenclature in tomato genetics including a list of known genes Jour. Heredity 46:22-26.
- Bergh, R.O. 1957. The inheritance of fruit size differences. Tom. Genet. Coop. Rep., 7: P. 3, 1957.
- Bianchi, A., 1954. Selezione di pomodoro con elevato contenuto in zuccheri nei frutti. "Tomato selection for high sugar content in the fruits". Genet. Agr. 4:58-67.
(Cited after Pl. Breed. Abst., 24(4):644, (3455), 1954).
- , Forlani, R. and Manunta, C., 1954. Sul contenuto in zuccheri dei frutti di varie razze di promodoro e di alcuni loro incroci. "The sugar content of various tomato varieties and of some crosses of them". Genet. Agr., 4:42-57.
(Cited after Pl. Breed. Abst., 41:644, (3454), 1954)
- Bohn, G.W., and Tucker, C.M., 1940. Studies on Fusarium wilt of the tomato. I. Immunity in Lycopersicon pimpinellifolium Mill. and its inheritance in hybrids. Mo. Agr. Expt. Sta. Res. Bull. No. 311:82 pp.
(Cited after Pl. Sci. Seminar; Camden New Jersey 57-78, 1960).
- Borzini, G., 1942. Shull'orientamento attuale nella lotta contro le malattie crittogamiche delle piante. "Present tendencies in the control of fungal diseases of plants". Italia Agric. 79(3): 163-166.
(Cited After Biol. Abs., 17(6):1567, (17343), 1943.)

- Bourdouil-Sosa, C., 1935. Remarques sur le poids des fruits de tomates hybrides (Solanum lycopersicum L.)
Bull. Mus. Nation. Hist. Nat. (Paris) 7(1):71-72
(Cited after Biol. Abs., 11(4):996, (9357), 1937).
- Brown, G.B., and Bohn, G.W., 1946. Ascorbic acid in fruits of tomato varieties and F_1 hybrids forced in the greenhouse.
Proc. Amer. Soc. Hort. Sci. 47:255-261.
(Cited after Biol. Abs., 21(4):973, (10012), 1947).
- Burton, G.W., 1951. Quantitative inheritance in Pearl Millet.
(Pennisetum glaucum)
Agron. Jour. 43:409-417.
- Butler, L., 1941. The inheritance of fruit size in the tomato.
Canad. Jour. Res. Sect. C. Bot. Sci., 19(6) : 216-224.
(Cited after Biol. Abs., 15(8) :1487, (16234), 1941.)
- , 1952. The linkage map of the tomato.
Jour. Heredity 43:25-35.
- Capinpin, J. M. and Cauton, S.C., 1948. Study of Mendelian biotypes in the segregating generation of tomato hybrids.
Philipp. Agric., 32:31-44.
(Cited after Pl. Breed. Abs., 21(2):439, (1432), 1951).
- Castle, W.R. and Wright S., 1921. An improved method of estimating the number of genetic factors concerned in cases of blending inheritance "
Science N.S., 54: p.223.
- Charles, D.R., and Smith, H.H., 1939. Distinguishing between two types of gene action in quantitative inheritance.
Genetics, 24(1):34-48.
(Cited after Biol. Abs., 13(1):164, (1734), 1939).

- Chmielewski, I.M. 1963. A factor from L. minuntum affecting the beta-carotene content.
Tom. Genet. Coop. Rep., 13: p 10., 1963.
- Crane, M.B., 1915. Heredity of types of inflorescence and fruits in the tomato.
Jour. Genetics, 5:1-11.
- Currence, T.M., Fogle, H., and Moore, J.F., 1951. Breeding tomatoes for ascorbic acid content.
Proc. Amer. Soc. Hort. Sci., 58:245-53.
(Cited after Pl. Breed. Abs., 22(4):616, (3055), 1952).
- Denisen, E.L., 1951. Carotenoid content of tomato fruits as influenced by environment and variety. II. Effects of plant nutrients, gas storage, and variety.
Iowa State Coll. Jour. Sci. 25(4):565-574.
(Cited after Biol. Abs., 25:(11)3086, (34807), 1951.)
- Dennett, R.K., 1950. The association of resistance to Fusarium wilt and Stemphylium leaf spot in tomato, Lycopersicon esculentum.
Proc. Amer. Soc. Hort. Sci., 56:353-357.
(Cited after Biol. Abs., 26(1):216, (2177), 1952).
-
- _____, 1952. The inheritance of shape index locule number, and stem colour in derivatives of a hybrid between Lycopersicum esculentum and Lycopersicum peruvianum.
Abs. Doct. Diss. Pa. St. Coll. 15:455-457.
(Cited after Pl. Breed. Abs., 24(1):722, (723), 1954.)
- Dokić, A., (1954). Studying dominant characters in intervarietal tomato hybrids.
Rev. Res. Wk. Fac. Agric., Belgrad, 2:28-104 (Serbian)
(Cited after Pl. Breed. Abs., 25(2):253 (1489), 1955.)
- Fleming, H.K., and Myers, C.E., 1937. Tomato inheritance, with special reference to skin and flesh colour in the Orange variety.
Proc. Amer. Soc. Hort. Sci., 35:609-624.
- Fogle, H.W., and Currence, T.M., 1950. Inheritance of fruit weight and earliness in a tomato cross.
Genetics, 35:363-380.

- Forlani, R., 1951. The genetics of the tomato.
Ital. Agric. 88:39-45.
(Cited after Pl.Breed.Abs., 21(3):697,(2230),1951)
- Foster, R.E., and Walker, J.C., 1947. Predisposition of tomato to Fusarium wilt.
Jour.Agric.Res., 74(516):165-158.
(Cited after Biol.Abs., 21(2):1526,(15698),1947).
- Frimmel, F. 1922. ÜBER Die Verebung Der Früchtrgrösse Der Tomaten Zischr.
Pflanzenzüchtung , 8:457-462.
(Cited after Jour.of Agric.Res., 55(2):141-152, 1937.)
- Fryxell, P.A., 1954. Genetics of locule number.
Tom.Genet.Coop. Rep., 4:p.10, 1954.
- Fuizat, Y. and Atkins, R.E., 1953. Genetic and environmental variability in segregating barley populations.
Agron. Jour. 45 : 414 - 420.
- Gaafar, M.I., 1962. Genetical studies in a tomato hybrid between Pearl Harbor and Stoner's exhibition.
M.Sc. Thesis.Fac.Agric. Ain Shams Univ.
- Giles, J.E.and Hutton, E.M., 1958. Combining resistance to the root-knot nematode Meloidogyne javanica (Treul) Chitwood, and Fusarium wilt in hybrid tomatoes.
Australian Jour. Agr. Res., 9(2):182-192.
(Cited after Biol. Abs., 32(9):2646(31703),1958.)
- Griffing, B., 1952. Contribution of genotype and environment upon the relationship existing among tomato yield components.
Genetics, 37:588-589.
-
- 1953. An analysis of tomato yield components in terms of genotypic and environmental effects,
Iowa Agric.Expt.Sta. Res.Bull. No.397:324-480
(Cited by Rick, C.M. and Butler L., 1956).

- Groth, B.H., 1912. The F_1 heredity of size, shape and number of locules in Tomato fruits.
N.J. Agr. Exp. Sta. Bull. No. 242
- , 1914. The "Golden Mean" in the inheritance of size.
Science, 39:581-584.
(Cited after Fleming, H.K. and Mayer, C.E., 1937).
- , 1915 Some results in size inheritance.
N.J. Agr. Exp. Sta. Bull. No. 278: 1-91.
(Cited by Fleming H. K., and Mayer C. 1937).
- Halsted, B.D., Owen, E.J., Shaw, J.K., 1905. Experiments with tomatoes.
N.J. Agr. Exp. Sta. Ann. Rept. 26:447-477.
- Harrison, A.L., 1960. Breeding of disease resistant tomatoes with special emphasis on resistant to Nematodes..
Proceed. Pl. Sci. Seminar, Camden, New Jersey, 57-78.
- Hawthorn, L.R. 1942. Breeding Summer tomatoes for increased size.
Proc. Amer. Soc. Hort. Sci. 40:390-394.
(Cited after Biol. Abs. 17(10):2416, (24987), 1943).
- Hedrick, U.P., and Booth, N.V., 1907. Mendelian characters in tomatoes.
Proc. Amer. Soc. Hort. Sci., 5:19-24.
(Cited after Yeager, A.F. 1937).
- Henderson, W.R. and Winstead, N.N., 1961. Reaction of tomato varieties and breeding lines to Fusarium oxysporum f. lycopersici race 1.
Plant Dis. Repor. 45:272-275.
(Cited after Pl. Breed. Abs., 31(1):1005, (5529), 1961).
- Houghtaling, H.B., 1935. A developmental analysis of size and shape in tomato fruits.
Bull. Terrey Bot. Club, 62(5):243-252.
(Cited after Biol. Abs., 10(3):735, (6815), 1936).
- Hutton, E.M., Margaret, M. and Giles, J.E., 1947. Fusarium wilt of tomato in Australia. 2-Inheritance of field immunity to Fusarium wilt in the tomato (Lycopersicon esculentum)
Austr. Coun. Sci. and Indust. Res. Jour. 20(4):468-474
(Cited after Biol. Abs., 22(8): 1947, (19317), 1948).

- Jenkins, J.A., 1949. The origin of tomato.
Econ. Bot. 2:379-392.
(Cited after Jenkins and Mackinney G., 1953).
- _____, 1951. Inheritance of size and shape of fruit.
Tom. Genet. Coop. Rep. 1: p.8, 1951.
- _____, and Mackinney, G., 1951. Colour in tomatoes.
California Agric., 5(2): 13-14.
(Cited after Biol. Abs., 25(8): 2033, (22498), 1951).
- _____, _____, 1953. Inheritance of carot-
enoid differences in the tomato hybrid yellow x
tangerine.
Genetics 38:107-16.
- _____, _____, 1955. Carotenoids of the
apricot tomato and its hybrids with yellow and
tangerine.
Genetics, 40(5):715-720.
- _____, _____, 1955. Carotenoids of yellow-
apricot and the tangerine apricot tomatoes.
Tom. Genet. Coop. Rep. 5: p.18, 1955.
- Kohler, G. W., Lincoln, R.E., Porter, J.W., Zscheile, F.P.,
Ralph, M.C., Harper, R.H., and Silver, W., 1947.
Selection and breeding for high β -carotene content
(Provitamin A) in tomato.
Bot. Gaz., 109(2):219-225.
(Cited after Biol. Abs., 22(5):1028, (10598), 1948).
- Larson, R.E., 1941. The F_1 combining ability of certain
tomato varieties.
Proc. Amer. Soc. Hort. Sci., 39:313-314.
- _____, and Currence, T.M., 1944. The extent of hybrid
vigour in F_1 and F_2 generations of tomato crosses.
Tech. Bull. Minnesota Agric. Expt. Sta. No. 164:1-32.
(Cited after Biol. Abs., 18(9), 1970, (18599), 1944).
- Le Rosen, A.L., Went F.W. and Zechmeister, L., 1941. Relation
between genes and carotenoids of the tomato.
Proc. Nation. Acad. Sci. U.S.A., 27(5):236-242, 1941.
(Cited after Biol. Abs. 15(8):1489, (16250), 1941).

- Lesley, M., and Lesley, J.W., 1946. A genetic study of flesh colour of fruit in hybrids of Lycopersicon esculentum Mill and L. hirsutum Humb. and Bonple. Genetics, 31(2): 223-232.
(Cited after Pl.Breed.Abs., 16(1):355, (1440), 1945-1946).
- Lincoln, R.E., and Porter, J.W. 1950. Inheritance of beta-carotene in tomatoes. Genetics, 35(2): 206-211.
(Cited after Biol.Abs., 24:2099, (22494), 1950).
- Lindstrom, E.W., 1926. Heredity correlation of size and colour characters in tomatoes. Iowa Agric.Exp.St.Res.Bull. No.93 : 99-108.
(Cited after Biol. Abs., 1(2): 231, (2213), 1927).
- _____, 1926. Linked inheritance in tomatoes. Jour.Sci., 1 (1) : 3-13.
(Cited after Biol.Abs., 1 (4): 467, (4739), 1927).
- _____, 1927. The inheritance of ovate and related shapes of tomato fruits. Jour.Agric.Res. 34(10) : 661-985.
- _____, 1928. Linkage of size, shape and colour genes in Lycopersicum. Zeitschr.Indukt.Abstam. -U.Vererb.Supplementh., 2: 1031-1057.
(Cited after Biol. Abs., 3(4-6):411, (4563), 1929).
- _____, 1928. Inheritance of shape in tomato fruits. Proc.Iowa Acad.Sci., 35:199-200.
(Cited after Biol. Abs., 4(11):2511, (26558), 1930).
- _____, 1929. Fruit size and shape genes on the first chromosome of the tomato. Proc.Iowa Acad.Sci., 36:189-140.
(Cited after Biol.Abs., 5(11):2632, (26697), 1931).
- _____, 1931. Linear order of four genes on the first chromosome of the tomato. Proc.Iowa Acad. Sci., 38:133-134.
(Cited after Biol. Abs. 7(6):1268, (12640), 1932).

- Lindstrom, E.W. 1932. Chromosome genes in the tomato.
Genetics, 17(3): 351-357.
(Cited after Biol. Abs., 7(7):1519,(15163),1938).
-
- _____, 1935. Segregation of quantitative genes in tetraploid tomato hybrids as evidence for dominance relations of size characters.
Genetics, 20(1): 1-11.
(Cited after Biol.Abs., 9(5) : 978, (8751),1935).
- MacArthur, J.W., 1928. Linkage studies with the tomato.
II Three linkage groups.
Genetics, 13(5) : 410-420.
(Cited after Biol.Abs., 2(3-5) : 404, (4565),1928).
-
- _____, 1934. Linkage groups in the tomato.
Jour.Genetics 29(1) : 123-133.
(Cited after Biol.Abs.9(1):14,(103), 1935).
-
- _____, 1934. Linkage group in tomatoes.
Jour. Genetics, 29:123-133.
-
- _____, 1941. Size inheritance in tomato fruits.
Jour.Heredity, 32(9):291-295.
(Cited after Biol.Abs., 16(2):305,(3578),1942).
-
- _____, and Butler, L., 1938. Size inheritance and geometric growth processes in the tomato fruits.
Genetics, 23(3):253-268.
- Mackinney, G., Rick, C.M. and Jenkins, J.A., 1954. Carotenoid differences in Lycopersicon : Hybrids of an unusual race of L. pimpinellifolium.
Proc. Nat.Acad.Sci. U.S., 40:695-699.
(Cited after Pl.Breed.Abs.25(2):253,(1488),1955).
- Mahmoud, I. and Kramer, H.H. 1951. Segregation for yield, height and maturity following a soybean cross.
Agron. Jour. 43 : 605 - 609.
- Manunta, C., 1948. Ricerche sui pigmenti del pomodoro in varie razze ed incroci. " Investigations on the pigments of the tomato in different varieties and crosses"
Genetica Agraria, 1(4): 351-359.
(Cited after Pl.Breed.Abs. 21(1):208,(717),1951).

- Marx, T., 1950. L-Ascorbinsäure und Tomaten " L-ascorbic acid and tomatoes "
Landw-Forsch; 2:74-80
(Cited after Pl.Breed Abs., 21(4):981,(3102),1951).
- McFarlane, J.S., Hartzler, E., and Frazier, W.A., 1946. Breeding tomatoes for nematode resistance and for high vitamin C content in Hawaii.
Proc.Amer.Soc.Hort.Sci., 47:262-270
(Cited after Biol.Abs., 21(1):529,(5156),1947).
- Miller, J.C., and John, J.M., 1939. Linkage studies of certain characters in tomatoes.
Proc.Amer.Soc.Hort.Sci., 37:884-885.
(Cited after Biol.Abs., 14(10):1507,(15682),1940).
- Morgando, F.A. 1948. Observations on the inflorescences of tomatoes.
Nuovo Gior. Bot.Ital., 55(3):374-375.
(Cited after Biol.Abs., 24(1): 210(2542),1950).
- Murphy, E.F., and Mildred, R.C., 1951. Tomatoes in Maine.
Maine Agric. Expt.Stat. Bull. No.489:1-69.
(Cited after Biol. Abs., 25(12), 3335,(37561),1951).
- Muthlendyok, E., 1943. Beiträge Zur Morphologie der Frucht der tomate (Solanum lycopersicum.)
1.Zeitschr.Pflanzenzücht., 25(2):117-163.
(Cited after Biol.Abs., 23:924,(8751),1949).
- Paddock, E.F., 1950. A tentative assignment of the Fusarium-immunity locus to linkage groups in tomato.
Genetics, 35 : 683-84.
- Pearson, D., 1962. The Chemical Analysis of foods. 5th.Ed.
J. and A. Churchill Ltd.
- Perry, F.E., 1915. The inheritance of size in tomatoes.
Ohio Nat. 15:473-495.
(Cited by Fleming , H.K., and Mayers, C.E. 1937).
- Poole, C.F., 1956. Improvement in tomato ascorbic acid content.
Proc.Amer.Soc.Hort.Sci., 68:443-451.

Porte, W.S., Doolittle, S.P., and Wellman, F.L., 1941. Hybridization of a Mosaic-tolerant, Wilt-resistant Lycopersicon hirsutum with Lycopersicon esculentum Phytopathology, 29:757-759, 1939.

_____, and Walker, H.B., 1941. The Pan American tomato a new red variety highly resistant to Fusarium wilt. U.S.D.A. Circ., 611:6 pp.
(Cited after Proceed.Pl.Sci. Seminar; Camden, New Jersey, 57-78, 1960).

_____, and Wellman, F.L., 1941. Development of interspecific tomato hybrids of horticultural value and highly resistant to Fusarium wilt. U.S.D.A. Circ., 584.
(Cited after Walker 1941).

Powers, Le Roy, 1939. Formulas for determining theoretical effects of certain genetic factors upon inheritance of quantitative characters, with special reference to a study of a Lycopersicon hybrid. Jour.Agric.Res. 59(8):555-577.

_____, 1939. Studies on the nature of the interaction of the genes differentiating quantitative characters in a cross between Lycopersicon esculentum and L. pimpinellifolium. Jour.Genetics, 39(1):139-170.
(Cited after Biol.Abs., 14(8):1237, (12919), 1940).

_____, 1941. Inheritance of quantitative characters in crosses involving two species of Lycopersicon. Jour.Agric.Res., 63(3):149-179.

_____, 1942. The nature of the series of environmental variances and the estimation of the genetic variances and the geometric means in crosses involving species of Lycopersicon. Genetics, 27(1):162. (An abstract).
(Cited after Biol.Abs., 16(4):801, (8766), 1942).

_____, 1942. The nature of the series of environmental variances and the estimation of the genetic variances and the geometric means in crosses involving species of Lycopersicon. Genetics, 27(6):561-575.
(Cited after Biol.Abs. 17(3):668, (8031), 1943).

- Powers, Le Roy, 1945. Relative yields of inbred lines and F_1 hybrids of tomato.
Bot.Gaz. 106:247-268.
- _____, 1950. Gene analysis of weight per locule in tomato hybrids.
Bot.Gaz. 112:163-74.
- _____, 1955. Components of variance method and partitioning method of genetic analysis applied to weight per fruit of tomato hybrid and parental populations.
U.S.Dept.Agric.Tech.Bull., 1131:1-64, 1955.
(Cited after Biol.Abs., 30:1839, (18571), 1956).
- _____, Channing, B.L., 1941. Inheritance studies on duration of developmental stages in crosses within the genus Lycopersicon.
Jour.Agric.Res., 63(3):129-148.
(Cited after Biol.Abs.15(10):2142, (23107), 1941).
- Powers, L., and Locke, L.F. and Garrett, J.C., 1950. Partitioning method of genetic analysis applied to quantitative characters of tomato crosses.
U.S.D.A. Tech.Bull. 889, 1950.
- Price, H.L. and Drinkard, A.W., 1908. Inheritance in tomato hybrids.
Virginia Agr.Exp.Sta.Bull. No.177 : 1-53.
(Cited after Adv. Genet.8:280, 1956).
- Reynard, G.B., and Margaret, S.K. 1942. Ascorbic (Vitamine C) contents of some tomato varieties and species.
Proc. Amer.Soc. Hort.Sci., 41:298-300.
- Rick, C.M., and Butler, L., 1956. Quantitative characters 2-The inheritance of fruit size.
Advances in Genetics, 8:330-341.
- Salib, N.S., 1965. Genetic studies in Lycopersicon esculentum.
M.Sc. Thesis, Fac.Agric. Assiut Univ.
- Schultz, J.H. and Kelly, E., 1952. A study of ascorbic acid content and fruit size in tomatoes segregation for those characters.
Proc.Amer.Soc.Hort.Sci. 59:458-62.
(Cited after Pl.Breed.Abs., 23(1):146, (748), 1953).

- Soost, R.K., 1956. A new pigment system.
Tom.Genet.Coop.Rep. 6:28-29, 1956.
- Suzuki, I., Sugohara, Y. and Todoka, S. 1962. (On breeding tomato for resistance to Fusarium oxysporum and the breeding system involved).
Engei Skinkenjo Hokoku (Bull.Hort.Res.Sta.
Ser.B.No. 1:57-73 (Japanese)
(Cited after Pl.Breed.Abs.33(3)487,(3874),1963).
- Thompson, A.E., 1955. Inheritance of high total carotenoid pigments in tomato fruits.
Science, 121:896-97.
(Cited after Pl.Breed.Abs., 26(1):16(763),1956).
- _____, 1957. Additional information on the inheritance of ry, a modifier gene for red colour in yellow tomato fruits.
Tom.Genet.Coop.Rep. 7:14-15,1957.
- Tomes, M.L. and Quackenbush, F.W., 1953. Two factor interaction governing beta-carotene content in the tomato.
Genetics 38:696-697(Abstract)
- _____, _____, and Kargl, T.E., 1956. Action of the gene B in biosynthesis of carotenoid in the tomato.
Bot.Gaz., 117:248-253.
- _____, _____, _____, 1958. Synthesis of B carotene in the tomato fruit.
Bot.Gaz. 119:250-53.
- _____, _____, and McQuisten M., 1954. Modification and dominance of the gene governing formation of high concentration of beta-carotene in the tomato.
Genetics, 39:810-817.
- _____, Nelson, O.E. and North, B.1953.
'The inheritance of' carotenoid pigment system in the tomato.
Genetics, 38(2):117-128.
- Walker, H.B., 1941. Diseases of vegetable crops.
McGraw-Hill Book Company, Inc. pp. 529.

- Walkof, C., and Hyde, R.B., 1955. Acid inheritance under monogenic control.
Tom.Genet.Coop.Rep. 5:p.30,1955.
- Wann, E.V., and McFerran, T., 1960. Studies of the inheritance of red colour in fruits of Lycopersicon pimpinellifolium Mill.
Proc.Amer.Soc.Hort.Sci. 76:524-529.
- Warren, P.A., 1924. Genetic studies in Lycopersicum. 1. The heredity of fruit shape in the garden tomato.
Mich.Acad.Sci., Arts, and Letters, Papers, 4(1): 357-394.
(Cited after Yeager, A.F., 1937).
- Weber, C.R., 1950. Inheritance and interaction of some agronomic and chemical characters in an interspecific cross in soybean (Glycine max x G. ussuriensis) Iowa Agr.Exp.Sta.Res.Bul. 374:765-816.
- Yeager, A.F., 1937. Studies on the inheritance and development of fruit size and shape in the tomato.
Jour.Agric. Res. 55(2):141-152.
- Young, P.A., 1956. ry , a modifier gene for colour in yellow tomato fruits.
Tom.Genet.Coop.Rep., 6:p.33,1956.
- _____, and MacArthur, J.W., 1947. Horticultural characters of tomatoes.
Texas Agr.Expt.Sta. Bull.No.608,1947.
(Cited after Barton, D.W. et al., 1955).
- Zechmeister, L., Le Rosen, A.L., Went, F.W. and Pauling, L., 1941. Prolycopene a naturally occurring stereoisomer of Lycopene.
Proc.Nat.Acad. Sci. 27:468-474.
(Cited after Jenkins and Mackinney, 1953).
- _____, and Went, F.W., 1948. Some stereochemical aspects in genetics.
Nature(London), 162:(4126), 847-848.
- Zielinski, Q.B., 1945. Fasciation in horticultural plants with special reference to the tomato.
Proc.Amer.Soc.Hort.Sci., 46:263-268.
- _____, 1948. Fasciation in Lycopersicon.
1 Genetics. 33 (4) : 405 - 428.
(Cited after Biol. Abs. 23 : 299, (2627), 1949).