

VIII. REFERENCES

- Achkova-Valkova, Z. and P. Stoeva (1978) : Bilateral hybridization *Lycopersicon peruvianum* with some self compatibility species. *Tomato Genetic Cooperative (TGC)*. 28:2.
- Agadzhanyan, A.M. (1985): Inheritance of the character protruding stigma in hybrids between self-compatibility tomato species. *Biologicheskii Zhurnal Armenii*. 38(3) 195-202. (C.F. Plant Breeding Abstr. 1985 55:755).
- Ammati, M.; Thomason, I.J. and Roberts, P.A. (1985): Screening *Lycopersicon* ssp. for new genes imparting resistance to root-knot nematodes. *Plant Disease*. 69(2) 112-115. (C.F. Plant Breeding Abstr. 1985 55: 5744).
- Antonelli, N.M. and J. Stadler (1989) : Chemical methods for efficient direct gene transfer to maize cells: treatment with (PEG). *Gene*. Breed. 43:113-122.
- Arnison, P.G. and W.G. Boll (1975) : Isoenzyme in cell culture of bush bean *Phaseolus vulgaris* : isoenzymatic differences between stock suspension cultures derived from a single seedling. *Can. J. Bot.* 53:261-271.

- Barbano, P. and L. D. Topoleski (1984): Postfertilization hybrid seed failure in *Lycopersicon esculentum* x *Lycopersicon peruvianum* ovules. J. Amer. Soc. Hort. Sci. 109(1): 95-100.
- Bendish, A. J. and E. T. Bolton (1967): Relatedness among plants asured by the DNA-agar technique. Plant Physiol. 42: 959-967.
- Benhki, R. M. and S. M. Lesley (1976) : In vitro plant regeneration from leaf explants of tomato. Can. J. Bot. (54): 2409-2414.
- Blobel, G. and V. R. Potter (1966) : Nuclei from rat liver : isolation method that combines purity with high yield. Science. 154: 1662-1665.
- Bonito, R. DI. (1985) : Embryo culture in tomato a technique to accelerate the breeding for disease. Genetica. Agraria. 39(3): 321-322 (c.f. Plant Breeding Abstr. 1986 56: 11156).
- Bonito, R. DI; and G. Ancora (1986) : Tomato (TMV) resistant lines obtained by hybridization with *Lycopersicon peruvianum*. Genetica. Agraria. 40(4) 446. (c.f. Plant Breeding Abstr. 1986 56: 9321).
- Bookout, D. and R. Noble (1987) : The effects of IAA on tomato plant regeneration. Plant Physiology (poster) .83 (4, supplement) 76. (c.f. Plant Breeding Abstr. 1986 56: 8427).
- Burton, K. (1956) : A study of the condition and mechanism of the dephenylamine reaction for coloremtric estimation of DNA . Biochem. J. 62: 315-323.

- Catherine,A.; Neal Leonard and D. Topoleski(1983): Effects of the basal medium on growth of immature embryos in vitro . J . Amer. Soc.Hort.Sci.108(3) 434-438.
- Charles, P. (1970): Isolation of deoxyribonucleic acids. pp.16 , 23 (ledoux,L.(ed.),uptake of informativemolecules by living cells. North Holland publishing company. Amesterd-am London).
- Chrambach,A.and D.Rodbard(1971): polyacryla-mide gel electrophoresis. Science.172: 449-451.
- D' Allesio,G. and A.R.Trim (1968) : A method for the isolation of nuclei from leaves. J.Exp.Bot.19:831-839.
- Davey,M.R.;E.L.Rech; and B.J.Mulligan (1989) : Direct DNA transfer to plant cells .Plant Molecular Biology . 13: 273-285.
- Davidson ,J.N.(1960): In Biochemistry of the Nucleic Acid.John Wily & Sons Ins.,New Youk ;Methan& Co.Ltd.,London.pp.66.
- Davis,B.J(1964): Disc electrophoresis.II.methods and application to human serum proteins.Ann.New york.Acad.Sci.121: 404-427.(citted after
- Delanghe,E. and E.Debruijine(1976): Continuous propagation of tomato plants by means of callus culture. Hort.Sci.4:221-227.

- Evesyrkin, W. and B. Katrina (1989) : A simple efficient method for the *Agrobacterium mediated* transformation of carrot callus cells. *Plant .Sci* . 61: 253-262.
- Flavell ,R.B. ;M.D. Bennett;J.B.Smith and D.B. Smith (1974) : Genome size and the protortion of repeated nucleotide sequence DNA in plants. *Biochemical. Genetics*. 12: 257-269.
- Gamborg,O.L.;R.A.Miller;and K.Ojima.(1968) : Plant cell cultures.I.Nutiernt reuirements of suspension cultures of soybean root cells. *Exp.Cell.Res*.50: 151-158.
- Gomez,K.A. and A.A.Gomez ,(1984): Statistical procedures for Agricultural Research.2nd Ed. John Wiley and sons pub.pp.139-153(686 pp.).
- Gradziel,T.M. and R.W.Robinson (1989) : Breakdown of self-incompatibility during pistil development in *Lycopersicon peruvianum* by modified bud pollination .*Sexual Plant Reproduction*.2(1): 38-42.
- Gradziel,T.M. and R.W.Robinson (1991) : Overcom-ing unilateral breeding barriers between *Lycopersicon peruvianum* and cultivated tomato, *Lycopersicon esculentum*. *Euphy-tica*.54(1):1-9.(c.f. Plant Breeding Abstr.1991 61:11711).
- Graham,D.E.(1970) : Isolation of DNA satellites by thermal chromatography in the presence of sodium perchlorate. *Analytical. Biochem-istry* .36:315-322.

- Gyldenholm, A.O. (1968): Macromolecular physiology of plastids V. on the nucleic acid metabolism during chloroplast development. *Hereditas*. 59:142-168.
- Heideveld, M.; C.E. Jansen; and A.J. Kool (1984) : Regeneration of tomato cultivar. In a new era in tomato breeding symposes IXth meeting EUCARPIA Tomato Working Group 22-24 May 1984: (c.f. *Plant Breeding Abstr.* 1985 55:5724).
- Hermesen, J.G.T and L.M. Taylor (1979): Successful hybridization of non-tuberos *Solanum tuberosum* Lindl, and tuber bearing *S. pinnatisectum* Dun. *Euphytica*. 28: 1-7.
- Hogenboom, N.G. (1972) : Breaking breeding barriers in *Lycopersicon* : 5. The inheritance of the unilateral incompatibility between *Lycopersicon peruvianum* and *Lycopersicon esculentum* and the genetics of its breakdown. *Euphytica*. 21:405-414.
- Honda, S.I.; T. Hogladorom and G.G. Laties (1966): A new isolation medium for plant organelles. *J. Exp. Bot.* 7 : 460-472.
- Jain, S.M.; R.J. Newton; and N.A. Tuleen (1988): Tissue culture and gene transfer in barley. *Current Science*. 57:59-70.
- Kartha, K.K.; O.L. Gamborg; J.P. Shyluk and F. Constabel (1976): Morphogenetic investigations on in vitro leaf culture of tomato and high frequency plant regeneration. *Z. Pflanzen-physiol.* 77:292-301.
- Kartha, K.K.; S. Champoux; O.L. Gambory and K. Pahl (1977) : I. In vitro propagation of tomato by shoot apical meristem culture. *J. Am. Soc. Hort. Sci.* 102: 346-349.

- Kesicki, E. (1979): New hybrids between *Lycopersicon esculentum* and *Lycopersicon peruvianum*. TGC. 29: 28.
- Kesicki, E. (1981) : Enigmas of species crossing in *Lycopersicon*. TGC. 31:8.
- Kitat, F.M. , (1959): Colohiploidy in *Citrus auratifolia* swingle, *C. Limetta* , Russo and *C. Sinensis* , osbeck. Ph.D. Thesis Fac. of Agric. Alex. Univ. pp.127.
- Kut, S.A ; and D.A. Evans (1982) : Plant regeneration from cultured leaf explants of eight wild tomato species and two related solanum species .In Vitro. 18: 593-598.
- Lamproye, A.; M. Hofinger; J.Y. Berthon and T. Gaspar (1990): (Benzo(b)selenienyl³) acetic acid: a potent synthetic auxin in somatic embryogenesis .Comptes Rendus de l' Academie Sciences. Series 3 , Sciences de la vie .311(3):127-132.
(cited after Hagan, A., M.Sc. Thesis, Fac. of Agric. El-Azhar University, 1990.)
- Laterrot, H. (1983) : Use of pollen Mixture technique in interspecific cross between *Lycopersicon peruvianum* .TGC. 33:3-4.
- Ledoux, L. and R. Huart (1969) : Fate exogenous bacterial deoxyribonucleic acids in .J.Mol.Biol . 43:243-262.
(cited after Hagan, A., M.Sc. Thesis, Fac. of Agric. El-Azhar University, 1990.)
- Leshanu, M.G.; V.A. Smirnov (1984) : Dynamics of pigments in tomato cell culture. otdel genetrasnit.

Akademiyanauk. Kishinev. Mold-avian.
SSR.135.(c.f. Plant Breeding Abstr.1984 54:10019).

Long,C.(1954):Science,progress42:266(citedafter:Davidson,1960) (cited after: Thorpe, T.A., editor, Plant Tissue Culture, Methods and Applications in Agriculture Academic Press,1981).

Lorz,H.;B.Baker;and J.Schell (1985) : Gene transfer to cereal cells ;treatment with(PEG) .Genet. Breed.43:113-122.

Marmur,J.(1961):A procedure for the isolation of deoxyribonucleic acid from micro-organisms.J.Mol.Biol.3:208-218.

Martin,F.W.(1961) : The inheritance of self-incompatibility in hybrids of *Lycopersicon esculentum* X *Lycopersicon chilense*. Genetics. 46:1443-1454.

Meizel,S.and E.R.M.Kay (1965) : Biochim.Biophys. Acta.103:431.(cited after : Ledoux,L.and R. Hurat 1969).
(cited after Hagran, A., M.Sc. Thesis, Fac. of Agric. El-Azhar University, 1990.)

Miltau,O.;D.Zamir;and J.Rudich (1984) : Breeding for chilling tolerance in tomato : an examination of selection criteria.In A new era in tomato breeding .Symopses,IXth meeting EUCARPIA tomato working group .22-24 May 1984.(c.f. Plant Breeding Abstr. 1984 54:5733).

- Murashige ,T.;and Skoog,F.(1962) : A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15:473-497.
- Nettancourt.D.;M.Devreux;U.Laneri;M.Cresti;E.Pacini;andG. Sarfatti(1974):Genetical and ultrastructural aspects of self and cross-inompatibility *Lycopersicon esculentum* and self-incompatibility *Lycopersicon peruvianum*. *Theor. AppleGenet.* 44: 278-288.
- O'Connell, M. A.;L. P. Hosticka and M. R. Hanson (1986) : Examination of genome stability in cultured *Lycopersicon*. *Plant.Cell.Rep.* 5(4):276-279.(c.f. *Plant Breeding Abstr.* 1986 56:761).
- Olivares,F.M.;J.R.Ġruz;A.A.Dela;and L.B.Salisi (1972) : Progress report on whitefly transmission of tobacco leaf curl virus to vegetable and weeds. *Plant Path.* 53: 1287.
- Padmanabhan,V.;E.F.Paddock;and W.R.Sharp (1974) : Plantlet formation from *Lycopersicon esculentum* leaf callus .*Can.J.Bot.* 52: 1429-1432.
- Paszkowski,J.;R.D.Shillito; M.Saul; V.Mandak; B.Hohn;and I.Potrykus (1984) : Direct gene transfer to plants. *The EMBO.Journal.* 3(12) :2717-2722.
- Pilowsky,W. (1976) : Breeding of disease resistant varieties of tomatoes. (Abstr.)*Phytoparasit-ica.* 4: 209.
- Popova-Konstantinova,M.(1974):Inheritance of fruit set percent in tomato interspecific F1 crosses .*TGC.* 25:10-11.

- Puertas, M.J.; De la Pena; Estades, B.; and Merin, F. (1984):
Chromosoma. 89:121-126. (cited after: De la Pena,
Lorz, H. And Schell, J., 1987).
- Rafique Uddin, M.; Stanley, Z. Berry; and A.D. Bisges (1988) :
An improved shoot regeneration system for
somaclone production in tomatoes. Hort. Sci.
23(6):1062-1064.
- Rick, C.M. (1963) : Search for *S-locus*. TGC. 13:22-23.
- Rick, C.M.; and P.G. Smith (1953) : Novel variation in tomato
species hybrids. Am. Nat. 87: 359-373.
- Ris, H.; A.E. Mirsky (1949) : Quantitative cytochemical
Determination of deoxyribonucleic acid with the
feulgen nuclear reaction. J. Gen. Physiol. (33)125-149.
- Schnider, W.C. (1957) : Determination of nucleic acids in
tissues by pentose analysis. In S.P. Colowick and
N.O. Kaplan (eds) Methods In Enzymology. III: 680-684.
- Schnider, P.H.; E.A. Seftor; J. Schell; and H.J. Bohnert (1985):
EMBO.J. 4:25-32. (cited after: De la Pena; A. Lorz
and J. Schell 1987).
- Segeren, M.I.; M.R. Sondahl; W.J. Siqueira; H.P. Mediana. Filho; H.
.Nagai; and A.L. Lourencao (1993) : Tomato to
breeding : I. Embryo rescue of interspecific hybrid
between *L. esculentum* X *L. peruvianum*. Revisia
Brasileira de Genetica. 16(2) :376380. (C.f
J. Announcement, Biological . Abstr. 1993.96(6):
03714).

- Sharon, M. K. and Daniel Lineberger (1983) : Genotypic differences in morphogenic capacity of cultured leaf explants of tomato .J.Amer.Soc.Hort.Sci.108(5) : 710-714.
- Sirotnak, F.M. and D.G. Hutchinson (1959): Biochim. Biophys Acta.36:246.(cited after: Ledoux, L. And R.Huart 1969).
- Smith, P.G. (1944) : Embryo culture of a tomato species hybrid. Am.Soc.Hort.Sci. 44: 413-416.
- Soyfer, V.N.; and Y.B. Titov (1981) : The absence of a definite gene specific effect in wheat after seed treatment with exogenous DNA .Mol.Gen.Genet. 182:361-363.
- Szilagyi, L.; and A.H. Nagy (1977) : Comparative analysis of green spontaneous albino mutant lines of *Nicotiana glauca* and *Nicotiana glauca* .Acta.Bot.Acad.Sci.Hung. 23:219-227.
- Taylor, I.B.; and Al-Kummer, M.K. (1982) : The formation of complex hybrids between *L. esculentum* and *L. peruvianum* and their potential use in promoting interspecific gene transfer. Theor.Apple.Genet. 61:59-63.
- Tanksley, S.D. (1979-a) : An efficient and economical design for starch gel electrophoresis. TGC. 29:37-38.
- Tanksley, S.D. (1979-b): Linkage chromosomal association and expression of *Adh-1* and *Pgm-2* in tomato. Biochem.Genet. 17 : 1159-1167.

- Tanksley, S.D. (1983) : Segregation of isozyme markers and cold tolerance in an interspecific backcross of tomato. *Theor. Appl. Genet.* 66(3/4):241-247.
- Tanksley, S.D.; and C.M. Rick. (1980-a) : Isozyme gene linkage map of the tomato : Application in genetics and breeding. *Theor. Appl. Genet.* 56:161-170.
- Tanksley, S.D. and C.M. Rick. (1980-b): Genetics of estrases in species *Lycopersicon*. *Theor. Appl. Genet.* 56:209-219.
- Thomas, B.R.; and D. Pratta (1981-a) : Efficient hybridization between *L. esculentum* X *L. Peruvianum* via embryo callus. *Theor. Appl. Genet.* 59:215-219.
- Thomas, B.R.; and D. Pratta (1981-b) : Breeding tomato strains for use in cell culture research. *PMB. News. Letter.* 2:102-105.
- Thorpe, L.; and T. Gaspar (1978) : Changes in isoperoxidases during shoot formation in tobacco callus. *In Vitro.* 14:522-526.
- Tokarskaya, V.L.; S.R. Umanskii and P.A. Nelipovich (1968) : Metabolically active DNA fraction. *Biochemistry.* 33 : 445-542.
- Uralets, L.I. (1984) : In vitro embryo culture in tomato. *Fiziologiya-i-Biokhimiya-Kulturnykh-Rastenii.* 16(5):495-499. (c.f. *Plant Breeding Abstr.* 1986 56:11143).

- Uralets, L.I.; M.I. Grat and V.K. Andryushchenko (1992) :
Production of interspecific *L. esculentum* with
L. peruvianum tomato hybrids in embryo callus
culture . Sel Skokhozyaistvennaya. Biologiya. 1:29-
35. (C.f. Plant Breeding Abstr. 1992 62:7983).
- Valkova-Achkova , Z. (1982) : Use of BC1 hybrids for
overcoming of incrossability of some species of the
genus *Lycopersicon* Mill. TGC. 32:50.
- Valkova-Achkova, Z. and V. Sotirova (1981) : Trigenomic
hybrid between *L. esculentum*, *L. chilenses*, and *L.*
peruvianum reproductive relationships and resistance
to *Corynebacterium michiganense*. TGC. 31:19-20.
- Verma, V.S. (1974) : Leaf curl virus . Plant . Path. 53 : 4492.
- Verma, J.; J. Hayati and Poonam (1980) : Resistance in
Lycopersicon spp. to tomato leaf curl disease in india.
Zeitschrift Fur Pflanzenkrankheiten und
Pflanzenschutz. 87: 137-144.
- Vnuchkova, V.A. (1977) : Development of a method for
obtaining regenerate tomato plants under tissue
culture conditions. Fiziol. Rast. 24:1094-1100.
- Wang, T.Y. (1967) : The isolation and purification of
mammalian cell nuclei. In L. Grossman and K. Maldava
(eds). Methods in Enzymology.
XII:417-421.

- Wijbrandi, J; J. G. M. Vos; M. Koornneef. (1988): Transfer of regeneration capacity from *Lycopersicon peruvianum* to *L.esculentum* by protoplast fusion .Plant Cell ,Tissue and Organ Culture. 12(2): 193-196.
- Wijbrandi,W.; Van Capelle.W.; C.J.Hanhart;E.P.Van Loenen Martinet Schuringa;and M.Kooknneef (1990) : Selection and characterization of somatic hybrids between *L.esculentum* and *L.peruvianum* .Plant.Sci. 70: 197-208.
- Yamauchi,A.;M.Imoto;Hibi and T.Komuro (1976) : Evidence that tobacco leaf curl virus 's not transmitted by greenhouse whitefly, trialeurodes vaporariorum. Ann. Phyto. Soc.Japan. 42:424-430.
- Yassin,A.M.(1973) : Leaf curl virus of tomato. Bilo. Abstr. 1973, 39351.
- Zaher,Nahid,A.M. (1973) : Leaf curl virus disease of tomato .M.Sc.Thesis,Fac.Agric.Cario.Univ.