

## REFERENCES

**Abbott, W.S. (1925):**

A method of computing the effectiveness of insecticides.  
J. Econ. Ent. 18, 265-267.

**Abdallah, S.A. (1983):**

Efficacy of certain synthetic pyethroids *Bacillus thuringiensis* and their combination on the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.).  
Bull. Ent. Soc. Egypt. Econ. Ser., 145-155.

**Abd El-Halim , Sawsan M. (1997a):**

Effect of certain bio control agents against the cotton leafworm, *Spodoptera littoralis* (Boisd.) in comparison with methamyl.  
Egypt J. Biol. Pestcontrol, 7 (1): 35-39.

**Abd El-Halim , Sawsan M. (1997b):**

Efficacy and residual effect of some microbial and chemical insecticides against of the cotton leafworm, *Spodoptera littoralis* (Boisd.).  
Egypt J. Biol. Pest Control, 7 (2): 73-78.

**Abd El-Hameed , Naglaa F.(1995):**

Evaluation of the efficacy of some bacterial preparations in controlling the Pink bollworm.  
M.Sc. Thesis, Fac. Agric., Zagazig Univ., Egypt pp. 120 .

**Abd El-Rahman, Amina .M. (1980):**

Microscopic studies on the fourth larval instar of the Mediterranean flour moth *Ephestia kuchiella*.  
Bull. Fac. Sci. Cairo Univ., 53: 341-347.

**Abd El-Rahman, Amina M. (1986):**

Morphological effect of actellic on haemocytes of *Spodoptera littoralis* (Boisd.) caterpillars.

Bull. Fac. Sci. Cairo Univ., 54: 545-559.

**Abdel-Megeed, M.I.; El-Refaei, S.A. and Zidan, A.A. (1984):**

The joint action of microbial and chemical insecticide combinations on the cotton leafworm, *Spodoptera littoralis* (Boisd.).

Bull. Ent. Soc. Egypt, Econ. Ser., 14: 119-126.

**Abdel - Rahman, Amina M.; El- Moursy, Ali A. ; Kares, Esmat A. and El- Mandarawy, Mona B. R. (1994):**

Total haemocyte counts, viscosity and volume of haemolymph in larvae of *Artoglia rapae* (L.) different biological treatments .

Bull. Ent. Soc. Egypt, 72 : 43 – 54 .

**Abou-Bakr, H. (1997):**

Efficacy of different *Bacillus thuringiensis* formulations against the cotton leafworm, *Spodoptera littoralis* (Boisd.).

Egypt J. Biol. Pest. Control, 7 (1): 7-11.

**Abou-Bakr, H.; Shalaby, F.F.; Assar, M.M. and Fahmy, M.F. (1993):**

Effectiveness of *Bacillus thuringiensis* varieties and strains versus L<sub>1</sub> of *Spodoptera littoralis* (Boisd.), (Lepidoptera: Noctuidae).

Egypt J. Biol., Pest Control, 3 (2): 245-252.

**Arnold, J.W. (1952):**

The haemocytes of the Mediterranean flour moth, *Ephestia kuehniella* Zell (Lepidoptera: Pyralidae).

Can. J. Zool., 30: 352-364.

**Arnold, J.W. and Hinks, C.F. (1976):**

Haemopoiesis in Lepidoptera 1-Multiplication of circulating haemocytes.

Can. J. Zool., 54: 1003-1012.

**Ashhurt, D.E. and Richards, A.G. (1964):**

Some histochemical observations on the blood cell of the wax moth *Galleria mellonella* L.

J. Morph., 114: 247-254.

- Azmy, N.M.; Rezk, G.N. and Hamed, A.A. (1981):**  
Histopathology of the cotton leafworm *Spodoptera littoralis* (Boisd.) treated with *Bacillus thuringiensis* Berliner.  
Res. Bull. Fac. Agric., Ain Shams Univ. , 1673, 5 (3): 7-11.
- Bartelt, R.J.; Mcguire, M.R. and Black, D.A. (1990):**  
Feeding stimulants for the European Corn Borer (Lepidoptera: Pyralidae): Additives to a starch-based formulation for *Bacillus thuringiensis*.  
Environ. Ent., 19 (1): 182-189.
- Beroza, M.; Sneh, B.; Yawetz, A.; Oron, U. and Honigman, A. (1984):**  
Commercial application of *Bacillus thuringiensis* var. *entomocidus* to cotton fields for the control of *Spodoptera littoralis* (Boisd.) (Lepidoptera-Noctuidae).  
J. Econ. Ent., 77 (6): 1530-1533.
- Chain, B.M. and Anderson, R.S. (1983):**  
Information of insect: The release of plasmatocyte depletion factor following interaction between bacteria and haemocytes.  
J. Insect Physiol., 29(1): 1-4.
- Charles, J. (1971):**  
Haemolymph. Arthropoda.  
Chem. Zool., 6: 64-118.
- Dunkle, R.L. and Shasha, B.S. (1988):**  
Starch encapsulated *Bacillus thuringiensis*: A potential new method for increasing environmental stability of entomopathogens.  
Environ. Ent., 17 (1): 120-126.
- Eduarda Aranda; Jorge Sanchez; Marnix Peferoen; Leopoldo Guerega and Alejandra Bravo (1996):**  
Interactions of *Bacillus thuringiensis* crystal proteins with the midgut epithelial cell of *Spodoptera littoralis* (Lepidoptera: Noctuidae).  
J. Invert. Path., 68: 203-212.

**El-Husseini, M.M. (1984):**

New approach to control the cotton leafworm *Spodoptera littoralis* (Boisd.) by *Bacillus thuringiensis* Berl. in clover fields.

Bull. Ent. Soc. Egypt, 12: 1-4.

**El-Husseini, M.M.; Agamy, F.A. and Abou-Bakr, H. (1997):**

Microbial control of *Spodoptera littoralis* (Boisd.) in Egyptian clover fields using three different entomopathogens.

Egypt J. Biol. Pest Control, 7 (1): 25-28.

**El-Mandarawy, M.B.R. (1992):**

Studies on the biological effects of *Bacillus thuringiensis* Berliner and gamma radiations on *Pieris rapae* L.

M.Sc. Thesis, Fac. of Sci., Cairo Univ.

**Endo, Y. and Nishiitsutsuje-Uwo, J. (1980):**

Mode of action of *Bacillus thuringiensis*  $\delta$  -endotoxin, histopathological changes in the silk worm midgut.

J. Invert. Path., 36: 90-103.

**Ephraim Cohen; Harel Rozen; Tamar Joseph; Serget Braun and Leon Margulies (1991):**

Photoprotection of *Bacillus thuringiensis kurstaki* from ultraviolet irradiation.

J. Inver. Path., 57: 343-351.

**Gad, A.M. (1951):**

The head capsule and mouth parts in the ceratopogonidae (Diptera-Nematocera).

Bull. Ent. Soc. Egypte, 35: 17-75.

**Gadallah, A.I.; Emara, S.A.; Houssien, N.M.; El-Kordy, M.W. and Abd El-Halim, S.A. (1990):**

Bioactivity of four commercial *Bacillus thuringiensis* formulations on some phytophagous lepidopterous insect pests of vegetables.

3<sup>rd</sup> Conf. Agric. Dev. Res, Fac. Agric. Ain Shams Univ. Cairo Egypt, Dec., 22-24.

**Gagen, S.Y. and Ratcliffe, N.A. (1976):**

Studies on the in vivo cellular reactions and fate of injected bacteria in *Galleria mellonella* and *Pieris brassicae* larvae.

J. Invert. Pathol., 28: 17-24.

**Gerald, R.S. and Earle, S.R. (1966):**

Histopathology of European-Corn-Borer Larvae treated with *Bacillus thuringiensis*.

J. Invert. Pathol., 9: 90-103.

**Hafez, A.A. (1993):**

Effects of a bioinsecticide of *Bacillus thuringiensis* var. *kurstaki* on *Spodoptera littoralis* (Boisd.).

Ann. Agric. Sc., Moshtohor, 31 (3): 1727-1736.

**Hamal, M.; Brownbridge, M.; Broza, M. and Seneh, B. (1991):**

Screening for highly effective isolates of *Bacillus thuringiensis* against *Spodoptera exempta* and *Spodoptera littoralis*.

Phytoparasitica, 19 (1): 9-21.

**Hamed, A.A. (1979):**

Histopathological studies on the effect of certain pathogens on some cotton pests.

M.Sc. Thesis, Fac. Agric., Ain Shams Univ., Cairo, Egypt.

**Himeno, M. (1987):**

Mechanism of action of delta-endotoxin from *Bacillus thuringiensis*.

J. Toxicology, Toxin 6 (1): 45-71.

**Hofmann, C. and Luthy, P. (1986):**

Binding and activity of *Bacillus thuringiensis* delta-endotoxin to invertebrate cells.

Archives Microbiology, 146 (1): 7-11.

**Horohov, D.W. and Dunn, P.E. (1982):**

Changes in the circulating haemocyte population of *Mondaca sexta* larvae following injection of bacteria.

J. Invert. Pathol., 40 (3): 327-339.

**Hunter, D. and Bomford, R.R. (1959):**

Clinical methods classcl and company.  
Ltd. London, 452, PP.

**Ignoffo, G.M. and Batzer, O.F. (1971):**

Microencapsulation and ultraviolet protectants to increase  
sunlight stability of an insect virus.  
J. Econ. Ent., 64 (4): 850-853.

**Jones, J.C. (1956):**

The haemocytes of *Sarcophaga bullata*.  
J. Morph., 99 : 233-257.

**Kares, E. A. ( 1994 ) :**

Effect of parasitism by *Zelet nigricornis* ( Walk. ) on the larval  
haemolymph of *Spodoptera littoralis* ( Boisd . )  
Ann. of Agric. Sc . Moshtohor , 32 ( 4 ) : 2139 – 2147 .

**Kares, E.A.; Abdel-Rahman, A.M.; El-Moursy, A.A. and El-  
Mondarawy, M.B.R. (1992):**

Qualitative and quantitative studies on the haemolymph of  
*Artogeia rapae* L. larvae after a bioinsecticidal and gamma  
treatment.  
Egypt. J. Biol. Pest Control, 2 (2): 131-141.

**Kasoper, H. (1965):**

Erörterungen Zur prafung von fungiziden in obrthau.  
Pflanzenschutz-Nachrichten. "Bayer", 18: 83-92.

**Kinsinger, R.A. and Mcgauchey, W.H. (1979):**

Susceptibility of populations of Indian meal moth and almond  
moth of *Bacillus thuringiensis*.  
J. Econ. Ent., 72(3): 346-349.

**Lai-Fook, J. (1973):**

Classification of blood cells in lepidoptera insects.  
J. Morph., 12: 139-179.

**Levinson, S.A. and Macfate, R.P. (1951):**

Clinical laboratory diagnosis Philadelphia: Lea and Febiger Co.

- Liu, Y.T.; Sui, M.J.; Ji, D.D.; Wu, H.; Chou, C. and Chen, C. (1993):**  
Protection from ultraviolet irradiation by melanin of  
mosquitocidal activity of *Bacillus thuringiensis* var.  
*israelensis*.  
J. Invert. Pathol., 62: 131-138.
- Lynch, R.E.; Lewis, L.C. and Berry, E.C. (1980):**  
Application efficacy and field persistence of *Bacillus*  
*thuringiensis* when applied to corn for European corn Borer  
control.  
J. Econ. Ent., 73: 4-7.
- Martourel, D.; Dllosta, J. and Roche, A. (1975):**  
Action sur le mesentron de *Pieris brassica* de latoxine de  
l'inclusion parasporale de *Bacillus thuringiensis*.  
Entomophaga, 10: 344-365.
- Mathavan, S. and Sudha, P.M. (1989):**  
Effect of *Bacillus thuringiensis israelensis* on the midgut cells  
of *Bombyx mori* larvae: A histopathological and histochemical  
study.  
J. Invert. Pathol., 53: 217-227.
- Mattes, O. (1927):**  
Parasitativе krankheiten der Mehlmoten larven und versuche  
uber ihre verwendbarkeit als biologische Bekämpfungsmittel  
(Zugleich ein Beitrag Zur Zytologie der Backterien).  
Sitzungsber. Ges. Barfarder Ungges.  
Naturwiss., Marburg, 62: 381-417.
- Mcguire, M.R. and Shasha, B.S. (1990):**  
Sprayable self-encapsulating starch formulations for *Bacillus*  
*thuringiensis*.  
J.Econ.Ent., 83 (5): 1813-1817.
- Mcguire, M.R. and Shasha, B.S. (1992):**  
Adherent starch granules for encapsulation of insect control  
agents.  
J. Econ. Ent., 85 (4): 1425-1433.

- Mcguire, M.R. and Shasha, B.S.; Leslie Lewis and Nelsen, T.C. (1994):**  
Residual activity of granular starch-encapsulated *Bacillus thuringiensis*.  
J. Econ. Ent., 87(3): 631-637.
- Miselyunene, I.S. (1976):**  
Pathologic changes in the morphology and number of haemolymph haemocytes of the cabbage butterfly larvae infested with endobacterim-3 (*Pieris brassicae*, Lep., Pieridae).  
Tsitologiya, 18 (10): 1220-1225.
- Morris , O. N. ( 1983 ) :**  
Protection of *B. thuringiensis* from inactivation by sunlight .  
Can. Ent . 115 : 1215 – 1217 .
- Nacht , S. ( 1991 ) :**  
Melanin , nature's own sunscreen polymer . In Cosmotic and Pharmaceutical Applications of Polymers. Edited by **G. G. Gebelein et al** Plenum Press , New York , . pp – 8394 .
- Nathransan, J.A. (1984):**  
Coffeine and related methylxanthine: Possible naturally occurring pesticides.  
Science (Washington, Dc.), 226: 184-187.
- Navon, A.; Wysoki, M. and Keren, S. (1983):**  
Potency and effect of *Bacillus thuringiensis* preparations against larvae of *Spodoptera littoralis* and Boarmia (Ascotis).  
Division of Entomology, ARO, The Volcani center. Bet Dagan Israel.
- Neuwirth, M. (1973):**  
The structural of the haemocytes of *Galleria mellonella* (Lepid .).  
J. Morph., 139: 105-124.
- Noack, S.; Reichmuth, C.H. and Wohlgemuth, R. (1983):**  
RH3-Ruckstande bei vorratsschutzbegasungen in Abhangigkeit von der konzentration, Einwirkungszeit und Lageradauer nach der Begasung.  
Zeitschrift fur Lebensmittel-Untersuchung und Forschung., 177: 87-93.

- Oddou, P.; Hartmann, H.; Radecke, F. and Geiser, M. (1993):**  
Immunologically unrelated *Heliothis* sp and *Spodoptera* sp. midgut membrane-proteins bind *Bacillus thuringiensis* Cry IA (b) delta-endotoxin.  
European. J. Biochemistry, 212 (1): 145-150.
- Possagy, M.; Fast, P.; Kaplan, H. and Carey, P.R. (1987):**  
The effect of sunlight on the protein crystals from *Bacillus thuringiensis* var. *Kurstaki* HD1 and NRD12: A Rman Spectroscopy study.  
J. Invert. Patho , 50 : 620 – 622 .
- Price, C.D. and Ratcliffe, N.A. (1975):**  
A reappraisal of insect haemocytes classification by the examination of blood from fifteen insect orders.  
Zeit. Zell. Mikrosko.Anat., 147 (4): 537-549.
- Ragaei, M. (1998):**  
Laboratory evaluation of shellac as ultraviolet screen for the *Bacillus thuringiensis* var. *entomocidus* against *Spodoptera littoralis*.  
Anz. Schdlingskde, Pflanzenschutz, Umweltschmtz, 71: 132-134.
- Raina, A.K. and Bell, R.A. (1974):**  
Haemocytes of the pink bollworm *Pectinophora gossypiella* during larval development and diapause.  
J. Insect. Physiol., 20: 21471-21480.
- Ramakrishnam, N. and Tiwari, L.D. (1972):**  
Observation on the blood cells of *Prodenia litura* F. in relation to nuclear polyhdrosis.  
Indian J. Ent., 34(4): 290-293.
- Rezk, G.A.; Azmy, N.M. and Hamed, A.A. (1981):**  
Laboratory studies in the use of *Bacillus thuringiensis* against lepidoptera cotton pests.  
Res. Bull. Fac. Agric., Ain Shams Univ. N., 1672, 8(4): 71-169.

- Ridgway, R.L.; Illum, V.L.; Farrar, R.R.; Calvin, D. ;JR; Flescher, S.L. and Inscoe, M.N. (1996):**  
Granular matrix formulation of *Bacillus thuringiensis* for control of the European corn Boror (Lepidoptera: Pyralidae).  
J. Econ. Ent., 89 (5): 1088-1094.
- Rie, J.V.; Jansens, S.; Hofte, H.; Degheele, D. and Mellaert, H.V. (1990):**  
Receptors on the brush border membrane of the insect midgut as determinants of the specificity of *Bacillus thuringiensis* delta-endotoeshior hky.  
Appl. Environ. Microbiol., 56 (5): 1378-1385.
- Rizk, S. A. (1991):**  
Effect of gamma radiation and some insecticides on cotton leafworm *Spodoptera littoralis* (Boisd.) .  
M.S.c. Thesis , Fac . of Sc . , Cairo Univ.
- Rosenberger, R. and Jones, J.C. (1960):**  
Studies on total blood cell count of the southern armyworm larvae, *Prodenia eridania* (Lepidoptera).  
Ann. Ent. Soc. Am., 53: 351.
- Salama, H.S. and Foda, M.S. (1982):**  
A strain of *Bacillus thuringiensis* var. *entomocidus* with high potential activity on *Spodoptera littoralis*.  
J. Invert. Pathol., 39: 110-111.
- Salama, H.S. and Matter, M.M. (1991):**  
Tolerance level to *Bacillus thuringiensis* Berliner in the cotton leafworm *Spodoptera littoralis* (Boisd.) (Lep. Noctuidae).  
J. Appl. Ent., 111 (3): 225-230.
- Salama, H.S. and Zaki, F.N. (1984):**  
Impact of *Bacillus thuringiensis* Berl. On the predator complex of *Spodoptera littoralis* (Boisd.) in cotton fields.  
Z. Ang. Ent. , 97 (5): 485-490.
- Salama, H.S.; Sharaby, A. and El-Din, M.M. (1993):**  
Mode of action of *Bacillus thuringiensis* and nuclear polyhdrosis virus in the larvae of *Spodoptera littoralis* (Boisd.).  
Insect. Sci. Applic., 14 (4): 537-543.

- Salama, H.S.; Abd El-Baset, M.S. and Ragaei, M. (1992):**  
 Mode of action of chemical additives in enhancing the potency of *Bacillus thuringiensis* against lepidopterous insects.  
 J. Appl. Entomol., 114 (2): 167-173.
- Salama, H.S.; Foda, M.S. and Sharaby, A. (1983):**  
*Bacillus thuringiensis* as a biological control against cotton pests in Egypt.  
 Croydon. U.K. British Crop. Protection Council, 781.
- Salama, H.S.; Zaki, F.N.; Salem, S. and Ragaei, M. (1995):**  
 The use of *Bacillus thuringiensis* to control two lepidopterous insect pests (*Agrotis ypsilon* and *Spodoptera littoralis*).  
 Anz. Schadlingskde., Pflanzenschutz, Umweltschutz, 68: 15-17.
- Salama, H.S.; Foda, M.S.; El-Sharaby, A.; Matter, M. and Khalafallah, M. (1981):**  
 Development of some lepidopterous cotton pests as affected by exposure to sublethal levels of Endotoxins of *Bacillus thuringiensis* for different periods.  
 J. Invert. Pathol., 38: 220-229.
- Sha, C.Y. and Xie, Q.J. (1992):**  
 Changes of haemocyte number, protein and esterase in the haemolymph of *Mythimna separata* (walker) infested with *Bacillus thuringiensis* Berliner.  
 Entomological, Knowledge, 29 (4): 215-217.
- Shalaby, F.F.; El-Khayat, E.F.; Hafez, A.A. and Kares, E.A. (1993):**  
 Impact of chemical and bioinsecticides and their combinations on entomophagous insects in cotton fields and damage by the cotton leafworm and bollworms.  
 Egypt J. Biol. Pest Control, 3(1): 71-84.
- Shams El-Din, A.M (1998a):**  
 Laboratory evaluation of Melanin as ultraviolet screen for *Bacillus thuringiensis* var. *entomocidus* against *Spodoptera littoralis* larvae.  
 Ann. Agric. Sc., Moshtohor, 36 (1): 553-560.

**Shams El-Din, A.M. (1998b):**

Effects of antioxidants and oxidative enzymes on the inactivation of *Bacillus thuringiensis* for cotton leafworm by simulated sunlight-UV.

Ann. Agric. Sc., Moshotohor, 36 (1): 561-568.

**Shapiro, M. and Robertson, I.J. (1990):**

Laboratory evaluation of dyes as ultraviolet screens for the gypsy moth (Lepidoptera: lymantriidae) nuclearpolyhedrosis virus.

J. Econ. Ent., 83: 165-172.

**Sirotna, M.I. (1949) :**

Genesis of the forming elements of blood in healthy and diseased caterpillars and moths of oak silkworm with jowndic .

Reports of Uasknnel , 4 : 22 – 28 .

**Sneh, B. and Gross, S. (1983):**

Biological control of the Egyptian cotton leafworm *Spodoptera littoralis* (Boisd.). (Lep., Noctuidae) in cotton and alfalfa fields using a preparation of *Bacillus thuringiensis* sp. *entomocidus*, supplemented with adjuvants.

Z. Ang. Ent., 95: 418-428.

**Sundaram, A. ; Leung, T.W. ; Devisetty, B.N. ( 1993 ) :**

Formulations and Application ,System , Berger , P. D :  
Devisetty , B. N .; hall , F. R. , Eds ; ASTMSTP 1183 ;  
American Society for Testing and Materials ; Philadelphia , P  
A . Vol . 13 . pp 227 – 241 .

**Sutter, G.R. and Raun, E.S. (1966):**

Histopathology of European corn borer larvae treated with *Bacillus thuringiensis*.

J. Invert. Pathol., 8 (8): 457-460.

**Tonada, Y. (1953):**

Susceptibility of imported cabbage worm of *Bacillus thuringiensis* Berliner.

Proc. Hawaii. Ent. Sco., 15: 159-166.

**Vankova, J. and Leskova, A.J. (1972):**

Changes in the haemolymph of *Galleria mellonella* L. caterpillars infected by the exotoxin of *Bacillus thuringiensis*.  
Acta. Entomol. Bohem., 69 (5): 297-304.

**Wang, Y.Z.; Zhang, L.H. and Huang, C.Y. (1990):**

Pathological modifications of haemocytes of indian meal moth after infection by *Bacillus thuringiensis*.  
Insect. Knowledge, 27 (1): 20-21.

**Woga, H. and Kitano, H. (1985):**

Morphological and functional characterization of the larval haemocytes of the cabbage white butterflies, *Pieris rapae crucivora*.  
Appl. Entomol. Zool., 20 (1): 1-7.

**Yi, S.; Pang, A.S.D. and Van Frankenhuyze, K. (1996):**

Immunocytochemical iacalization of *Bacillius thuringiensis* CryI toxins in the midguts of three forest insects and *Bombyx mori*.  
Can. J. Microbiol., 42: 634-641.