

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	3
I- Studies on the rest stage of pink bollworm	3
I.1- Diapause of pink bollworm <i>Pectinophora gossypiella</i>	3
I.2- Induction, duration and termination of pink bollworm diapause	3
I.3- Factors affected on survival of resting <i>P. gossypiella</i>	5
I.3.1. - Placement of sticks cotton stacks	5
I.3.2- Temperature	5
I.3.3- Ectoparasitic mite (<i>Pyemotes herfsi</i>)	6
I.3.4- Bacteria and virus	7
I.4- Spring emergence of <i>P. gossypiella</i>	7
I.4.1- PBW spring emergence related to weather temperature	7
I.4.2-Relationship between degree-day (or accumulated heat units and pink bollworm spring emergence	8
II- Biological effects of <i>B. thuringiensis</i> on the spiny bollworm	9
III- Histological studies	
III-1 Histopathological effects of <i>B.t.</i> against lepidopterous insects	12
III-2 Pathological effects of δ - eldotoxin	
IV- Control of bollworm	13
IV-1 Chemical control	13
IV.2- Microbial control with <i>Bacillus thuringiensis</i>	16
IV.3-The use of mixture of chemical insecticides with <i>B.</i> <i>thuringiensis</i> and insect growth regulators	18
MATERIALS AND METHODS	
I- Studies on the diapausing larvae of the pink bollworm <i>Pectinophora gossypiella</i>	21
I.1- The effect of ambient conditions on diapausing larvae of <i>P.</i> <i>gossypiella</i>	21
I.2- Forecasting of <i>P. gossypiella</i> spring emergence	22
II- Biological and histological studies about the effect of <i>B.t.</i> var. <i>aizawai</i> (Xentari) on <i>E. insulana</i>	22

	Page
II.1. Biological studies	22
II.1.1- Artificial diet used for rearing spiny bollworm <i>Earias insulana</i>	22
II.1.2- Rearing procedure of <i>Earias insulana</i> (Boisd.)	22
II.1.3- Preparation of different concentrations of <i>B.t.</i> var. <i>aizawai</i> (Xentari) and determination of the LC ₅₀	22
II.1.4- Some biological aspects resulted from feeding the neonate larvae of <i>E. insulana</i> on diet treated with the LC ₅₀ of Xentari	23
III- The histological studies	24
IV- Evaluation of some bollworms control programs	25
IV.1- Insecticides	25
IV.2- Field experiments	27
IV.2.1- Season 1994	27
IV.2.2- Seasons 1995 and 1996	31
IV-3 Efficiency of insecticides	32
IV-4 Statistical analysis	32
IV-5 Determination of the most effective control program	33
RESULTS AND DISCUSSION	34
I- Studies on resting larvae of the pink bollworm (<i>Pectinophora gossypiella</i>)	34
I.1- The excluded resting larvae during 1994/95 and 1995/96 rest period	34
I.2- Follow-up of obtained alive resting larvae during 1994/95 and 1995/96 season	36
I.2.1- Normal death	36
I.2.2- Bacterial infection	36
I.2.3- Virus infection	36
I.2.4- <i>Pyemotes herfsi</i>	39
I.3- Spring emergence of <i>P. gossypiella</i>	40
I.4- Forecasting of timing of PBW spring emergence on the basis of degree-days (dd's) or A.H.U. during 1994/95 and 1995/96 seasons	44
II- Effect of different concentrations of Xentari (<i>B. thuringiensis</i> var. <i>aizawai</i>) against neonate larvae of the spiny bollworm	47

	Page
IV.6- Effect of certain control programs on larval populations of bollworms (<i>P. gossypiella</i> and <i>E. insulana</i>)	103
IV.7- Effect of certain control programs on percentages of infested cotton bolls with <i>P. gossypiella</i>	107
IV.7.1- Season 1994	107
IV.7.2- Season 1995	112
IV.7.3- Season 1996	116
IV.8- Effect of certain control programs on percentages of infested cotton bolls with <i>E. insulana</i>	116
IV.8.1- Season 1994	116
IV.8.2- Season 1995	121
IV.8.3- Season 1996	121
IV.9- Effect of certain control programs on percentages of PBW and SBW infestation	126
IV.9.1- Season 1994	126
IV.9.2- Season 1995	131
IV.9.3- Season 1996	131
IV.10- General Conclusion	135
SUMMARY	141
REFERENCES	145
ARABIC SUMMARY	